

Geochemical patterns in the soils of Zeeland

Nederlandse Geografische Studies 330

Geochemical patterns in the soils of Zeeland

Natural variability versus anthropogenic impact

J. Spijker

Utrecht, 2005

Koninklijk Nederlands Aardkundig Genootschap/
Faculteit Geowetenschappen Universiteit Utrecht

This publication is identical to a thesis submitted in partial fulfilment of the requirements for the degree of Doctor of Philosophy (Ph.D.) at Utrecht University, The Netherlands.

Promotor: Prof. Dr. P.A. Burrough
Faculteit Geowetenschappen, Universiteit Utrecht
Co-promotores: Dr. P.F.M. Van Gaans
Faculteit Geowetenschappen, Universiteit Utrecht
Dr. S.P. Vriend
Faculteit Geowetenschappen, Universiteit Utrecht

This research was financially supported by the Provincie Zeeland

ISBN 90-6809-370-3

Copyright © J. Spijker, p/a Faculteit Geowetenschappen, Universiteit Utrecht, 2005

Niets uit deze uitgave mag worden vermenigvuldigd en/of openbaar gemaakt door middel van druk, fotokopie of op welke andere wijze dan ook zonder voorafgaande schriftelijke toestemming van de uitgevers.

All rights reserved. No part of this publication may be reproduced in any form, by photoprint, microfilm or any other means, without written permission by the publishers.

Printed in the Netherlands by Labor Grafimedia b.v. - Utrecht

Cover: silhouettes of the Zeeland landscape, after a photograph of R. Sluiter.

Suivre l'étoile
Peu m'importent mes chances
Peu m'importe le temps
Ou ma désespérance
Et puis lutter toujours
Sans questions ni repos

La Quête, Jacques Brel

Contents

1	Introduction	15
1.1	History and approaches to soil management in the Netherlands	16
1.2	Soil quality and so called “soil quality maps”	18
1.3	Aim of this thesis	18
1.4	Outline of this thesis	19
2	Geology and pedology of Zeeland	21
2.1	Introduction	21
2.2	Basic geographic information	22
2.3	Geology of Zeeland	23
2.3.1	Pleistocene period: terrestrial phase	24
2.3.2	Forming of the Basal Peat	24
2.3.3	Calais deposits, first inundation	25
2.3.4	Holland Peat	25
2.3.5	Duinkerke deposits	25
2.4	Human activities	27
2.4.1	Dikes and embankments	27
2.4.2	Peat excavation	29
2.4.3	Parceling and land reconstruction	29
2.4.4	Agricultural inputs	30
2.5	Landscape and soil geomorphology	30
2.5.1	Heartlands	32
2.5.2	Channel ridges and creeks	33
2.5.3	Accretion polders	35
2.5.4	Groundwater	36
2.6	Summary and conclusion	38

3	Natural and anthropogenic patterns of covariance and spatial variability of minor and trace elements in agricultural topsoil	41
3.1	Introduction	41
3.2	Materials and methods	42
3.2.1	Area	42
3.2.2	Spatial variance and sampling theory	43
3.2.3	Sampling and chemical analysis	45
3.2.4	Data analysis	46
3.3	Results and discussion	48
3.4	Conclusions	56
4	Sampling and analyses for a regional environmental soil geochemical survey	61
4.1	Introduction	61
4.2	Material and methods	62
4.2.1	Study area	62
4.2.2	General research approach	64
4.2.3	Field sampling	65
4.2.4	Sample preparation	66
4.2.5	Chemical analyses	67
4.2.6	Precision and accuracy	68
4.2.7	Outliers	70
4.2.8	Parametric levelling	70
4.3	Results and interpretation	72
4.3.1	Precision	72
4.3.2	Accuracy	73
4.3.3	Outliers	74
4.3.4	Parametric levelling of data	75
4.3.5	Comparison between XRF and ICP-MS	76
4.3.6	Parameter selection and summary statistics	78
4.4	Discussion and conclusions	82
5	Enrichment and natural variability versus anthropogenic impact	95
5.1	Introduction	95
5.2	The Zeeland geochemical soil data	97
5.3	Concepts and methods	97

5.3.1	Assessing enrichments and baselines in attribute space	97
5.3.2	Spatial representation and spatial anomalies	104
5.4	Results and interpretation	105
5.4.1	Univariate statistics for the Zeeland topsoil data.	105
5.4.2	Univariate comparison with reference data and legal threshold levels	107
5.4.3	Correlation and regression with Al	113
5.4.4	Spatial patterns of topsoil enrichments	118
5.5	General discussion and conclusions	123
6	Regional diffuse geochemical patterns and processes	129
6.1	Introduction	129
6.2	Material and methods	130
6.2.1	Study area and geochemical topsoil data	130
6.2.2	Principal component analysis	131
6.2.3	Fuzzy c-means clustering	133
6.2.4	Statistical software packages	134
6.3	Results and interpretation	134
6.3.1	Principal components analysis	134
6.3.2	Anthropogenic subprocesses	138
6.3.3	Fuzzy clustering	139
6.4	General discussion and conclusions	144
7	Assessment of regional DDT concentrations in the soils of Zeeland	149
7.1	Introduction	149
7.2	DDT, properties and history	150
7.3	Study area	151
7.4	Method	152
7.4.1	Collection of data	152
7.4.2	Basic statistical analysis	153
7.4.3	Analysis of variance	154
7.4.4	Regional variability	154
7.4.5	Comparison with external data and normative values	155
7.4.6	DDT breakdown	155
7.5	Results	156
7.5.1	Data merging and selection	156

7.5.2	Exploratory statistics	157
7.5.3	Local and regional variability	159
7.5.4	Regional overview	161
7.5.5	Comparison with external data and normative values	162
7.5.6	DDT breakdown	166
7.6	Discussion and conclusions	167
8	Synthesis	171
8.1	Methods used	171
8.2	Achievements of this study	172
8.3	Implications	174
	Bibliography	175
	Abstract - Geochemical patterns in the soils of Zeeland, natural variability versus anthropogenic impact	185
	Samenvatting - Geochemische patronen in de bodems van Zeeland, natuurlijke variabiliteit versus antropogene impact	193
	Dankwoord	203
	Curriculum Vitae	205

Figures

2.1	Topography of Zeeland	22
2.2	Pictures of the landscape of Zeeland. Both pictures were taken in Zeeuws-Vlaanderen.	23
2.3	Simplified geological cross section of Zeeland	24
2.4	Simplified geology of the Duinkerke deposits	26
2.5	History of embankment	28
2.6	Historical usage of P and K fertilisers	31
2.7	Landscape typology	32
2.8	Simplified soil map of Zeeland	33
2.9	Digital elevation model of Zeeland	34
2.10	Schematic cross section of the origin of ridge inversion of tidal channels	35
2.11	Excerpt from digital elevation model showing pool areas	36
2.12	Excerpt from digital elevation model showing channel ridges	37
2.13	Excerpt from digital elevation model showing accretion polders	38
3.1	Location map	43
3.2	Stacking of levels of variation design	44
3.3	Stacking of levels of sample design	46
3.4	Contribution of the hierarchical variance components	53
3.5	Summary of cumulative spatial variance patterns	58
3.6	Variance Ratios	59
4.1	Topography of Zeeland	63
4.2	Sample locations per survey	67
4.3	Principal concepts of levelling	71
4.4	Example of levelling of Ga	77
4.5	Comparison between selected elements from the XRF and ICP-MS	81
4.6	Histograms for selected elements and oxides analysed in the topsoil samples	85
5.1	Map of Zeeland with sample locations	98

5.2	Cumulative probability plots of a selection of components	106
5.3	Regional distribution of a selection of components expected not to be influenced by human processes	108
5.4	Regional distribution of a selection of elements expected to be influenced by human processes	109
5.5	Ratio of median values for Zeeland topsoil and the average upper continental crust	110
5.6	Ratio of median values for the Zeeland topsoil and the median of the subsoil	112
5.7	Ratio of median values and legal threshold value	114
5.8	Scatter plots of Al_2O_3 versus selected elements	116
5.9	Box plots of the enrichment ratio e	120
5.10	Spatial pattern of the enrichment for selected elements	121
5.11	Spatial representation of the contamination index	123
5.12	Spatial pattern of the ratio with the S-value for selected elements	124
6.1	Topography of Zeeland and sampling locations	131
6.2	Scree plot of the eigenvalues from the first 8 of 30, unrotated, components	135
6.3	PCA loading plots	137
6.4	Regional distribution of rotated factor scores	139
6.5	FCMC centroids of the scaled data.	141
6.6	Box and Whisker plots of the distribution of the 8 elements for each cluster	142
6.7	Box and Whisker plots of the distribution of selected elements for each cluster . . .	143
6.8	Regional distribution of the FCMC memberships of the sample locations.	144
7.1	Topography of Zeeland	152
7.2	Selected sample locations from the soil information systems of Zeeland	158
7.3	Boxplots of log scaled concentration $\sum DDx$ by survey	160
7.4	Cumulative frequency plot concentration $\sum DDx$	161
7.5	Variance versus approximate scale	163
7.6	Semivariogram of the regional $\sum DDx$ concentrations	164
7.7	Regional overview of $\sum DDx$ concentrations	165
7.8	Relation of organic matter (TOC) and $\sum DDx$	166
7.9	Relation between DDT, its isomers DDE and DDD, and the number of half-times .	168

Tables

3.1	Statistical summary after outlier replacement of the geochemical data for the clayey top soils of Zeeland	48
3.2	Varimax rotated principal component factor loadings	50
3.3	Correlations of concentrations of elements, not included in the PCA-V	50
3.4	Relative percentages of the variance components for each level and their total . . .	51
3.5	Fuzzy cluster memberships	56
3.6	Relation between covariance structure (PCA-V) and the clustered UANOVA variance patterns	57
4.1	Year, region and number of sample locations	66
4.2	Attributes analysed per survey and method	69
4.3	Average precision for ICP-MS	73
4.4	Average precision for XRF	74
4.5	Accuracy of the XRF analyses	74
4.6	Accuracy of the ICP-MS analyses	75
4.7	Number of outliers	76
4.8	Results of the levelling for the XRF data	78
4.9	Results of the levelling for the ICP-MS data	79
4.10	Differences and correlations between elements analysed by both XRF and ICP-MS	80
4.11	Selected elements	80
4.12	Statistical summary of the geochemical data for the topsoil of Zeeland.	86
4.13	Statistical summary of the geochemical data for the subsoil of Zeeland.	90
5.1	Elements selected from the Zeeland dataset	97
5.2	Statistical concepts for assessing enrichments and baselines	99
5.3	<i>S</i> values and regression parameters	104
5.4	Percentiles of Zeeland topsoil	107
5.5	Pearson correlation with topsoil Al ₂ O ₃	115
5.6	Regression constants of the regression of subsoil concentrations	119

5.7	Enrichment ratios in the topsoil	119
5.8	Enrichment differences for the topsoil	120
6.1	Varimax rotated factorloadings and communalities	136
6.2	Robust correlation of other elements with PCA factor scores	138
6.3	Correlation of anthropogenic elements	140
6.4	Median values for each FCMC cluster	145
7.1	Databases and survey area used for merged dataset	157
7.2	Summary values for $\sum DDx$ and total organic carbon	159
7.3	Results ANOVA on the replicated samples	162
7.4	Percentage of sample points exceeding the <i>S</i> -value.	165
7.5	Selected $\sum DDx$ vaues from other areas.	169

1 Introduction

“As knowledge increases, and as the earth becomes more crowded, geochemical information is becoming an increasingly significant factor in decisions affecting the management of the overall environment. Ultimately this information affects human survival.”

Darnley (1995)

It is widely understood that the geochemical environment plays a profound role in the existence of life on earth. A large part of this geochemical environment is the thin layer between the earth crust and the atmosphere. Yet, Bridges and van Baren (1997) already argued that the significance of this thin layer, also called soil, is not always sufficiently appreciated, despite its vital role for human well-being. As they summarise, this natural body of mineral, animal, and plant organic matter forms a critical link between the inanimate rocks and minerals and the living plants and animals.

Soil, especially its biogenic content, can affect the biosphere by its role in global climatic processes. The bidirectional relations between consumption and production of CO₂, CH₄ and N₂O can directly influence climatic changes (Mosier, 1998). More directly, soils also play a major role in human health. Mineral nutrients are, mainly, transferred from soil to humans by plant and animal foods. Deficiencies, excesses, or imbalances in this dietary source can have deleterious influences. Such influences can occur on rather large, even global, scale; for example the atmospheric transport of persistent organic pollutants of soils from moderate climatic areas to colder areas, where the cold conditions cause precipitation of the pollutants and subsequent uptake in the local food chain (Abrahams, 2002). These examples of the close relation between soil, health, and global sustainability confirm that soil plays a critical role as a major interface in our environment and that soil quality can be an important indicator for sustainable environmental management (Doran, 2002).

Soil quality, as defined by Karlen et al. (1997), is “the capacity of a specific kind of soil to function, within natural or managed boundaries, to sustain plant and animal productivity, maintain or enhance water and air quality, and support human health and habitation”. Soil quality is often described in terms of physical (texture, thickness of topsoil layer, water holding capacity), chemical (organic C, Total N, pH, extractable N, P, K) and biological (biomass, soil respiration) parameters (Wienhold et al., 2004). An evaluation of the various quality indicators and their change over time may identify if sustainable management is reached or that soil quality is aggrading/degrading. The type of soil information to be

evaluated depends of course on the function of the soil, be it a natural ecosystem, food production, or just the base for building (Nortcliff, 2002). The framework and criteria for evaluation often depend on policy choices by relevant authorities or organisations.

This thesis is concerned with obtaining an overview of general (geo)chemical soil quality, within the framework of diffuse anthropogenic pollution and sustainable soil management in the Netherlands.

1.1 History and approaches to soil management in the Netherlands

Within the Netherlands soil quality assessment and related monitoring in part follow a thematic approach, as for example soil acidification, effects of manure and fertilizer use, salinization, or other effects of hydrological management. These thematically based activities are divided between regional authorities, and often subject to evolving political interests and influence at this level (Busink & Postma, 2000; Mol et al., 2001). The other, in some ways more consistent and georeferenced approach to obtain soil quality information is rooted in the Dutch soil sanitation and soil management legislation. To place the present study into context, this latter approach and its history are explained below. Most of the following is derived from the good and extensive description by de Roo (2003) of the history of Dutch environmental policy and legislation, and their influence on (spatial) planning.

A major event which instigated environmentalism and seeded the first ideas about soil management in the Netherlands was the discovery of severe soil pollution at Lekkerkerk in 1980. During a new housing development chemical waste was used as building material to level the soil. The dumping of the waste, containing substances such as xylene and toluene causing the area to be uninhabitable, resulted in an unprecedented scandal. This, however, was just a tip of the iceberg, as many cases of soil pollution followed. This increase in cases of soil pollution led in 1983 to the Soil Remediation (interim) Act, which later resulted in the Soil Remediation Guidelines (*Dutch: Leidraad bodemsanering*) (VROM, 1999), and these guidelines defined legal limits which were the foundation of soil remediation until the 1990s. A major concern resulting from the Soil Remediation Guidelines was that the soil should be remediated until the soil was “clean”, i.e. below the threshold values defined in the guidelines. This resulted in a tremendous increase in costs and stagnation of spatial planning as spatial development and building of houses was by law not allowed to proceed until the conditions of the guidelines were met. The soils, after remediation, should be free of contaminants and able to support many functions. This led, for example, into clean soil patches within historically diffusely contaminated urban areas, hence wasting effort and money.

Objections to the guidelines, cost increase and stagnating spatial development, led to a revision of the soil remediation policy at the beginning of the 1990s, resulting in three major developments. The first was the development of ‘active soils management’, which

should provide authorities during their spatial planning and decision making with up-to-date information about the soil condition. Also powers and tasks were decentralised to lower authorities. This should help the remediation efforts, which were until then frustrated by a lack of advance information. Another development was the changeover from the old system of legal limits to a system of limits based on actual risks. With these new limits the urgency of remediation could be assessed and the necessity of remediation was not determined by the nature of the pollution but by its seriousness. With the 'urgency', together with the third development that the government would in principle not pay the bill for clean-up, pressure could be put on the parties responsible for soil remediation. With the new Soil Protection Act of the mid 1990s, which could enforce parties to remediate, soil remediation became a responsibility of society, hence spreading the effort and costs of urgent soil remediation.

Another major implication originated from the second half of the 1990s. The awareness grew that for a more coordinated, instead of consecutive, soil policy, integration of remediation measures and spatial development policies was necessary. Soil management moved towards a more decentralised, integrated, and market dynamic policy. Therefore reliable information about the nature and extent of soil contamination was needed, as this information can be used to make agreements with non governmental parties involved in the spatial planning. This coordination also led to the function oriented approach which implied it was not necessary to remediate the soil until it was "clean" but that the remediation effort could be location specific supporting the future function of the soil (de Roo, 2003).

One of the necessities of the new policy was information about the condition of the soil and the extent of contaminated sites. The government acknowledged this in the third national environmental policy plan (NEPP 3) (VROM, 1997). Their wish was to obtain an overview of the country wide soil quality and to provide of soil management methods. Their goal was to tackle the soil pollution problems, especially the costs and efforts of remediation, within 25 years. In 2002 the stepping stones (*Dutch: Stappenplan Landsdekkend Beeld 2005*) towards this country wide overview were presented and two major tracks were discerned. One track was the inventory of existing polluted sites and the second track aimed at soil management and soil quality. This information should provide a country wide overview of the soil quality to facilitate large scale spatial development. This is where the 'active soil management policy' returns. This policy should lead to the realisation of sustainable soil management and to adequately and efficiently manage existing soil contamination so as to prevent frustration of spatial planning operations (Leenaers et al., 1999).

The purpose of the country wide overview was also to re-evaluate the existing soil background values and how these relate to the existing legal limits. The observation that part of these legal limits were within the range of considered natural background values urged the need for insight in actual soil values. These values should then also provide in a new reference for soil remediation (Leenaers et al., 1999).

1.2 Soil quality and so called “soil quality maps”

One of the major tools in fulfilling the country wide overview of the soil quality and supporting active soil management are the so called “soil quality maps” (*Dutch: bodemkwaliteits kaarten*). The interim guidelines for such maps (*Dutch: Interim-Richtlijn Bodemkwaliteitskaarten*) were presented in 1999 (VROM, 1999; van der Gaast et al., 1998; van Lienen et al., 2000). The starting point of the guidelines was the Building Material Decree (Eikelboom et al., 2001). This came into full operation in 1999, describing the quality criteria for building material, which according to the decree includes soil, used as land fill or site preparation material. This decree should prevent materials being exposed in the surface environment that leach potential contaminating compounds. (Eikelboom et al., 2001).

The aim of the soil quality maps is to indicate if soil, used in terms of the decree as building material, originating from an area was “clean” relative to the legal limits and therefore dispensed to be transported to another area. Without such maps and dispensation, each shipment of soil needs to be extensively surveyed to see if it is “clean”, substantially increasing the costs of spatial development (Anonymous, 1999; van Lienen et al., 2000). When soil is not clean, it is generally not allowed to be used as building material, so spreading of contamination is prevented.

Besides the use as a dispensation tool, from the information of the soil quality maps the country wide overview of contaminated sites and re-evaluation of background levels could also be obtained and these should facilitate active soil management and sustainable soil use as well (Leenaers et al., 1999). However, due to the urgent need for spatial development and increasing soil transport the latter has become less obvious.

The current Dutch soil quality maps, based on the building material decree, are aimed to indicate or predict when “soil quality” exceeds certain legal limits. Since these limits indicate the risk of soil pollution they are actually “soil pollution risk maps” (van der Gaast et al., 1998; Swartjes, 1999). Also, according to the guideline, soils are grouped based on their soil concentrations of environmental priority compounds relative to the legal limits (VROM, 1999). This reduces the soil quality, usually not including other biological, physical and chemical soil indicators, to a black and white concept: legal limits are exceeded or not. Therefore, in my opinion, the current soil quality maps are neither the means of providing the new background values as wished by the NEPP 3 nor do they facilitate sustainable soil management. However, they are still suitable for fulfilling the country wide overview of contaminated sites and facilitate enforcement of the building material decree.

1.3 Aim of this thesis

Despite the changeover from curative measurements and remediation to prevention of contamination, the evolution from soil quality maps to soil pollution risk maps is not surprising. The aim of Dutch soil policy is still to reduce remediation costs and prevent stagnation of spatial development. The next step towards sustainability seems, yet, a small one. An

overview of soil background values as reference is a basic need as a starting point for sustainable management. However, to provide the overview of soil background values and assess the impact of anthropogenic processes on the natural soil composition, another approach is needed.

This thesis aims at assessing patterns in geochemical soil composition and distinguishing natural variability from anthropogenic impact. The variability is assessed in geographical space, where the spatial interaction of soil components takes place, and in attribute space, where interaction between soil constituents is exposed. Patterns from both spaces can then be related to processes influencing the soil composition.

Based on these patterns, and related processes, anthropogenic influence can be distinguished from natural variability. This can provide tools and information to support the wide overview as required by the NEPP 3.

The chosen study area for this research is the Province of Zeeland, in the south west of the Netherlands. The large rural area of young Holocene deposits with a rich human history makes it a suitable area for testing the main hypothesis that human influence leaves a distinct and quantifiable pattern, based on variability within the geographical and attribute space of the soil composition.

1.4 Outline of this thesis

This thesis is addressed in 6 chapters and a synthesis. Each chapter is written to stand on its own and can be read more or less independently of the others.

To understand soil geochemistry in an area it is necessary to understand the factors that determine the geochemical variability. In the next chapter a description is given of the research area, the rural part of the province of Zeeland. It is shown how both geological and pedological processes, and human activities have influenced the soils and shaped the landscape. This information is the basis on which a regional geochemical survey can be realized. In such a survey interest focuses on regional features and this is only useful when small scale variability does not dominate the observed regional patterns. In chapter 3 it is hypothesised that distinct spatial patterns of variability exists for groups of data related to anthropogenic or geochemical processes. This requires that the sampling strategy should anticipate those different patterns in variability by using composite or single samples.

In chapter 4 the process of obtaining a province wide geochemical dataset is described. This dataset should contain information on pristine soil composition and information how this composition is altered by human processes. The noise level in such a dataset should be as low as possible so the large variety of different processes can be discerned at a adequate level of significance. Also, sources of variance and their magnitude should be determined. While chapter 3 focusses on field scale variability, chapter 4 examines analytical sources of variance and bias. The final dataset presented in chapter 4 should provide a true reference dataset, which is suitable for environmental and geochemical assessment within a regional context.

For environmental legislation and soil management policies it is essential to have a good overview of the present day soil composition. Although the above dataset presents a reference of actual soil values it is assumed that these values are imprinted by human activities. In chapter 5 it is hypothesised that this anthropogenic imprint can be established using a geo-chemical baseline that comprises the natural variability in soil composition. Since no standard approach exists, this chapter shows different approaches which differ in degree of complexity and efficacy. The assessment started in chapter 5 continues in chapter 6, where the question as to what extent the human contribution determines the regional soil composition, is discussed. It is supposed that the anthropogenic imprint on the soils results in distinguishable regions where specific human processes have a relevant contribution to the soil composition. If such regions exist then they can be important for the zoning of soil pollution risk maps.

The first six chapters focus on inorganic geochemical soil composition based on a specially collected dataset. From a practical point of view it is also interesting to obtain some knowledge about levels of organic pollutants in soil, such as persistent organic pesticides, as they impose certain environmental risks and are a concern of the local authorities. In chapter 7 the occurrence of DDT in the Zeeland soils is assessed based on data derived from soil information systems associated with soil quality maps. The extent of the contamination by DDT residues, relative to legal limits and values obtained from other areas, and variability are studied. Besides insight into the DDT residue concentrations, this chapter also demonstrates the level of suitability of data from soil information systems for a regional and environmental assessment.

2 Geology and pedology of Zeeland

2.1 Introduction

Zeeland (Sea-land) is probably the most intriguing Dutch province within the Delta of the rivers Rijn, Maas, and Schelde*. Its name testifies of the sea as the primary element in its formation and history, which had full play up to the end of the 20th century. However, it was not only nature that formed the province, people also had a substantial influence on the landscape. Their biggest struggle was to keep what was threatened to be taken by the sea, a continuing battle against the rising water. The final victory, more or less, was considered to be gained with the completion of the “Delta works” in 1989. Due to the mix of the works of Mother Nature and Man, the province is unique in the Netherlands and Europe. The intriguing part is the complex interaction between natural and human processes together with a rich human history.

A description of the research area of this study, that aims to unravel the factors that determine the Zeeland soil geochemical variability, should therefore address both the natural and human history of the province. Geology is undoubtedly a key natural factor in soil formation and soil composition. Variation in parent material is expected to be responsible for the larger part of the geochemical variability. With respect to human activities, interest is in the local processes that led to the current landscape, soil morphology, and possibly soil chemistry. Understanding the parent materials and human activities provides insight into the patterns of variation of Zeeland soils. Given this prior information, assumptions can then be made on which to base the realization of the geochemical soil survey and hypotheses can be formed to explain the features encountered in the resulting data.

It is thus the aim of this chapter to give an overview of both the geological/pedological history and the human activities in the province, and to summarize the major landscape units resulting from their interaction. Since the geochemical soil survey concentrates on the rural area of Zeeland, this overview will have the same focus. The information for this overview has been taken from soil studies which were performed as part of soil mapping (STIBOKA, 1964, 1967, 1980; van der Sluis et al., 1965; Bazen, 1987; Pleijter & Wallenburg, 1994) and the Dutch Geological Survey (Zagwijn, 1991; Vos & van Heeringen, 1997). No new research was done to append or validate the already available data.

*Rhine, Meuse and Scheldt respectively, in this thesis the Dutch names are used




Figure 2.1. Topography of Zeeland, showing names (italic) of the peninsula's and areas, and some cities (bold)

2.2 Basic geographic information

The location and topography of Zeeland are given in figure 2.1. Zeeland is located in the south-west of the Netherlands, along the coast of the North Sea and bordering Belgium. Besides the main land of Zeeuws-Vlaanderen it consists of several islands and peninsulas. The general landscape of Zeeland is a relatively flat and open country of polders with dikes and villages on the horizon. Illustrations of this landscape are shown in figure 2.2. The province has a maritime-climate resulting in moderate summers and winters. Annual



Figure 2.2. Pictures of the landscape of Zeeland. Both pictures were taken in Zeeuws-Vlaanderen.

rainfall varies from 750-800 mm, which is close to the annual rainfall for the Netherlands as a whole. Evaporation is about 600-615 mm a year. The average temperature is around 10.1-10.4°C, which is the highest for the Netherlands (Heijboer & Nellestijn, 2002). The mild temperature and the fact that the province has the largest number of yearly sun hours makes it popular with tourists

The total area of Zeeland, about 2930 km², is divided into 1440 km² for agriculture, 120 km² for nature of which 30 km² are forest, and 240 km² for other purposes such as buildings, recreation and industry. Almost 1140 km² is water. This roughly means that one third of the province is water and 80% of the remainder is agricultural area. The larger part of the agricultural area (980 km²) is arable land, while only 150 km² are meadows. The main crops are corn, root- and tuberous plants. Livestock is usually sheep and to a lesser extent cattle. Despite its large areal extension and subsequent important role in shaping the landscape, the economic role of agriculture is marginal. Only 4% of the added value is earned from agriculture and fisheries and under 7% of the labour is in the agricultural sector (<http://www.zeeland.nl/zeeland/>).

2.3 Geology of Zeeland

A simplified geological cross section through Zeeland is depicted in figure 2.3. The entire section consists of non-consolidated sediments, mainly of Holocene marine origin (Westland Formation). A chronological description of the deposits is given below.




Figure 2.3. Simplified geological cross section of Zeeland. Time is ^{14}C years before present (BP). After Bazen (1987).

2.3.1 Pleistocene period: terrestrial phase

During the Pleistocene, in the late Weichselian period, large eolian cover sands with some fluvial Schelde fresh water sediments (i.e. gyttjas and organic clays) were deposited (Twente Formation). Large parts of the current North Sea were dry, leaving Zeeland out of the marine realm. The cover sands of the Twente Formation only surface in the south of Zeeuws-Vlaanderen, bordering Belgium. The formation dips downwards towards the north and is the basis of the Westland Formation for Zeeland.

2.3.2 Forming of the Basal Peat

At the beginning of the Holocene, due to the melting of the land ice, the sea level rose sharply, more than 75 cm per 100 year. This rise dominated the sea-land interactions during the major part of the Holocene. When the transgression reached Zeeland, the area transformed into a tidal basin. This basin was formed in the paleo Schelde valley that was bordered by two large sand ridges (escarpments) at the east and the south (Brabantse Zoom and Rilland Ridge). The rising sea water level and the increasing water supply from the river Schelde also resulted in higher groundwater levels. In combination with the warmer climate, peat developed on the marine margins, called Basal Peat. With the inland movement of the shoreline the peat moved along. At the borders of the paleo valley the peat transits into the Holland Peat (see below). Through compaction and subsequent burial, the Basal Peat has been reduced to a layer with a thickness of just a few decimetres found at depths of 5 m at Walcheren to 20 m at Schouwen-Duiveland. In Zeeland the Basal Peat

does not outcrop.

2.3.3 Calais deposits, first inundation

Around 6000-3000 BC tidal channels emerged where most of the peat was removed by erosion, these were filled in with clay deposits, the Calais deposits. During this period the influence of the sea diminished and a coastal barrier system developed. Behind this barrier, (lagunal) basins formed in which clayey sediments deposited, while the tidal channels consisted of more sandy sediments. The decline in sea level tipped the balance between sea level rise and filling of the basin. As a result the tidal area filled up, the tidal channels decreased in size and the coastal barrier could expand laterally until only two major tidal inlets were left, positioned around the current openings of the Westerschelde and Oosterschelde. At present, outcrops of the Calais deposits are only found at Schouwen-Duiveland in an area called the “Prunje” east of Serooskerke (see figure 2.1). It otherwise is usually found at depths of 1.5-4.5 m. In Zeeuws-Vlaanderen, due to the higher elevation, the Calais deposit is absent.





2.3.4 Holland Peat

Due to the attenuation of the sea level rise and the shrinking of the tidal channels, drainage conditions deteriorated, and behind the now almost closed coastal barrier, only some openings were left by rivers, a new peat landscape emerged. It gradually changed from brackish to freshwater, due to the fresh water supply of the Schelde. The quantities of nutrients diminished though, and the peat flora changed from eutrophic to oligotrophic vegetation. Only along the Schelde did the peat stay eutrophic. The peat landscape was crossed by several streams of which the Oosterschelde was the largest one. These streams kept active openings in the coastal barrier through which the sea could gain influence on the land again. Holland Peat is found almost everywhere in Zeeland, unless it is excavated or eroded, in a layer with a thickness ranging from 0.5-2.0 m usually within a few metres from the surface. During excavations in the Middle Ages most of the peat was removed on Schouwen-Duiveland and Walcheren.

2.3.5 Duinkerke deposits

Figure 2.4 shows the geological map of the Duinkerke layer after Vos & van Heeringen (1997). The Duinkerke deposits are the youngest deposits in Zeeland and are formed in a similar tide-influenced environment as the Calais deposits. The basal layer of the Duinkerke deposits often contains peat detritus, even lumps, resulting from the erosion of the Holland Peat layer. The clayey and silty nature of the deposits has been gradually influenced by subsequent dike building, which took the tidal areas out of the marine realm and consequently ended the sedimentation process.

Geology

-  Channel deposits >1250 AD
-  Channel deposits <1250 AD
-  No deposits, Pleistocene sands
-  Cover layer, clay/sand




Figure 2.4. Simplified geology of the Duinkerke deposits, after Zagwijn (1991); Vos & van Heeringen (1997)

During the earliest embankments (1000-1200 AD) heavy clays were deposited due to the quiet marine environment, resulting in the so called Heartland areas (Dutch: Oud- en Middelland). The quiet environment and slow sedimentation rates are presumably also the cause of the observed decalcified nature of the sediments. Moreover, these areas, so called “schorren” or “poelen” were often overgrown by vegetation. This source of organic material led to a reducing environment in the sediments.

Areas embanked after 1200 AD are called Newlands; these are areas much closer to the tidal channels in which more fine sand, silt, and shell debris were deposited due to the

more turbulent environment. The type of embanked environment also gradually changed. First, mostly the accretions alongside the already existing dikes were embanked; later on also the large tidal channels were closed. These were usually areas with barely any, or no, vegetation. Decalcification is much less than in the Heartlands due to the higher shell content and younger age.

A former geological model, describing the transgressive and regressive phases and resulting in a subdivision of the Duinkerke deposits as 0 to IIIb layers, is nowadays considered invalid. Reasons for this, as given by Vos & van Heeringen (1997), are that the described transgressive phases are not synchronous in space, are insufficiently proved, and sometimes are incorrectly dated. However, this model is frequently referenced in pedologic descriptions of Zeeland that originate from before the invalidation of the model and it is therefore mentioned here for completeness.

2.4 Human activities

Indications of the first human presence in Zeeland date back to Late Neolithic Ages (3100-2100 BC), while during the Middle Roman Age (circa 100 BC-100 AD) the area was already densely populated. However, these early inhabitants of Zeeland were barely responsible for the formation of the present day landscape. The most dramatic changes occurred with the advancement of industry and technology. From the studied literature it appears that there have been three major processes since the Middle Ages: 1) the dike building and reclamation of the area, 2) the excavation of the Holland and Basal peat, and 3) the reparceling of the agricultural land and Modern Time land reconstructions after the most recent floods (STIBOKA, 1964, 1967, 1980; van der Sluis et al., 1965; Bazen, 1987; Pleijter & Wallenburg, 1994; Vos & van Heeringen, 1997). I further consider that the soils are influenced by input of metals and organic compounds from agricultural activities and atmospheric deposition. Although the latter is not often recognised from the literature describing the landscape and soil geomorphology, it is generally accepted that such influences are ubiquitously present in soils.

2.4.1 Dikes and embankments

Around 1000 AD the first dikes were raised but it was not until the 12th century AD when the inhabitants started the systematic embankment of large areas such as the island of Beveland (see figure 2.5 Rijkswaterstaat (1971)). The early dikes primarily defended the Heartlands against the continuous threat of the sea and the regular storm surges during these times. These dikes were relatively low since the elevation of the salt marshes was about 1.5 m above sea level. From the 13th century onwards the dike building becomes more offensive. The salt marshes outside the first embankments, sometimes newly formed, were also reclaimed. These dikes were also motivated by the need for agricultural land and the growing prosperity due to the trading with Vlaanderen and England. It was this prosperity and the rise of abbeys that resulted in more political influence and money to invest in




Figure 2.5. History of embankment

more systematic land reclamation on a larger scale. Often new mud flats/tidal flats silted up against the new dikes, which were successively reclaimed. These are the so called “open aan-waspolders” which can be loosely translated as accretion polders. Also dams were built in large tidal channels that were subsequently reclaimed. The reclamation of Zeeland was not an easy task, the sea showed its character during several storm surges by breaking through the dikes. This resulted in large losses of valuable land that, in particular in the early ages with primitive equipment, were very difficult to reclaim again.

The reclamation of land continued until far into the 20th century as did the storm surges and floods. But the people of Zeeland kept continuing and improving the reclamation and protection of their land. In the 17th century these improvements led to a change of the device of the province to “luctor et emergo” (I struggle and overcome), which sounded far more heroic than the old device “Domine, serva nos, perimus” (Lord, save us, we perish).

The last two floods, one induced by the allied forces during World War II in Walcheren and the other the great flood of 1953, resulted in large land reconstruction works. The dramatic flood of 1953, when 1835 people perished, forced the Dutch government to start the Delta Works, an impressive engineering project in which the dikes were raised and the large open sea arms were closed with dams (Goemans & Visser, 1987). With the completion of the Delta Works in 1988 Zeeland reached its present form.

2.4.2 Peat excavation

The exploitation of peat, mainly the Holland peat, started in the Roman Age but it is the excavation during the Late Middle Ages that most influenced the present landscape. The areas from which the peat was excavated were mostly located in the at present pool areas.

While the peat was used for fuel during the Roman Age, the Holland Peat became more salty and covered by a thin layer of marine sediments due to the frequent floodings and inundations of the peat landscape. The peat layers were then exploited for their salt content. Exploitation of the peat (“darinckdelven” or “moernereren” and refinement of the salt (“selnering”) took place until prohibition in the 15th century. The process involved removing the cover sediment layer and digging out the peat. By burning the peat and mixing the ashes with sea water, and subsequent refining of the salt in lead, copper and iron pans, the valuable salt was obtained. The excavations, both within and outside reclaimed areas, resulted in a lowering of the surface level by about 1 m, which endangered such areas by making them even more vulnerable to floods. This was one of the main reasons for the prohibition. Moreover, the exploited areas were left behind in a bad state, as the surface level lowering was highly variable at short distances. The resulting “hollebollig” (concave/convex) landscape has short scale fluctuating soil moisture, and soil moisture salt content, and very poor drainage.

2.4.3 Parceling and land reconstruction

The allotment of the (agricultural) land among the inhabitants differed between the Heartlands and the Newlands. The early farmers of the Heartlands started on the grounds of the sandy channel ridges, which were easier to rework with their primitive equipment, resulting in irregular and patchy patterns of parcels. In the Newlands the parceling was more rational. A cooperative group of people would buy the salt marshes and reclaim the area. The new polder was then divided into the mainland, high quality grounds, and in the “volgerland”, lower quality ground. Each participant in the cooperative group then obtained parcels on both kinds of ground, or was paid money if no mainland grounds could be obtained. This resulted in a more regular, block wise, parceling. By inheritance, sales, and other changes over time, the parcels became divided later into smaller and smaller areas.

The last floods during World War II and 1953 caused severe destruction. This led to large scale land reconstruction works that re-parceled the agricultural land, leading to a better organised landscape. New, straight roads were built and ditches were replaced. Also the

“hollebollig” landscape, with its low agricultural value, was reworked. The cover layer was removed and the underlying layers leveled. The cover layer was then placed back. In large areas, mainly those where peat excavations had occurred, this disturbed the soil profile of the subsoil but kept the upper layer intact.

2.4.4 Agricultural inputs

The intensification of agriculture, facilitated by the large scale land reconstructions, initially also led to an increase in the usage of fertilisers and pesticides as compared to pre-war levels. This increase was subsequently reduced due to more environmental awareness and governmental regulations in the last part of the 20th century (RIVM, 2004). Based on historical statistics figure 2.6 shows an example of the usage of P and K inorganic fertilisers for the Netherlands.

Some fertilisers are known to contain substantial amounts of heavy metals like Cd, Pb, Cu, and Zn (de López Camelo et al., 1997; Gimeno-García et al., 1996; Nash et al., 2003). The accumulation of these heavy metals in soils has also been recognised in the Netherlands and for some parts of Zeeland (Groot et al., 2001; van Drecht et al., 1996; RIVM, 2004). Residues of persistent organochlorine pesticides, such as DDT, γ -HCH, and Dieldrin, are found as a heritage from the past. It is not uncommon that the values of organochlorine pesticides in Dutch soil exceed legal permissible limits (Groot et al., 2001; RIVM, 2004). Besides local input by agricultural practice, atmospheric deposition from sometimes remote sources may also contribute to the deposition of organochlorine pesticide (Rovinsky et al., 1995; Villa et al., 2003) and heavy metals (Koeleman et al., 1999). Although these processes are not specific to the Zeeland region, in fact some of them are global, they are expected to exert a major influence on the (geo)chemical soil composition of Zeeland.

2.5 Landscape and soil geomorphology

The final geomorphology of Zeeland is a result of both the natural and human processes as described above. While varied and sometimes complex, they resulted in only a few major landscape types and associated soil types that will be outlined in some detail below: the Heartlands, channel ridges, and accretion polders (“open aanwasolders”) Dunes, beach sands, the small area of Pleistocene sands in the south of Zeeland, and areas outside the dikes are left out of this description since they are not within the focus of the geochemical research. A brief description of the major groundwater systematics will be given to complement this section.

The relevant major landscape types are depicted in figure 2.7 (Halfwerk, 1996). While the classification differs from that based on the invalidated Duinkerke 0 to IIIb regression model (see §2.3.5) the major features are the same. The oldest pool areas are mainly formed by the Heartlands the Newlands are subdivided into polders and large tidal channels.

The soils of Zeeland have developed on marine clay deposits and according to the classi-









Figure 2.6. Historical usage of P and K fertilisers. Source: CBS statline database (<http://www.cbs.nl/>)

fication of the FAO they are referenced as fluvisols (FAO, 1990). A simplified soil map is shown in figure 2.8. As the various soil types on the official soil map differ mainly in clay and sand content, this simplified map depicts the main groups: sands, silts, and clays. Secondary features as calcite content, soil profile differences, and structure properties are not shown. The similarity between the soil map and the land type map of figure 2.7 is as expected. This shows the close interaction between landscape development (by reclaiming land) and soil development.

As a further illustration a detailed digital elevation model is shown in figure 2.9. It shows the moderate elevations ranging mainly from a few metres below to a few metres above mean sea level. With this model the various features of the Zeeland landscape can be made

Typology

-  Dunes
-  Pool areas, Heartlands
-  Old tidal channels, Newlands
-  Other polders, Newlands
-  Pleistocene cover sands




Figure 2.7. Landscape typology, for further explanation, see text.

visible.

2.5.1 Heartlands

The soils of the Heartlands can be divided into the old polders, and the so called pool areas (Dutch “veenpoelen”), that were formed by excavation of peat. Although these areas are considered the oldest, the cover layer can be significantly younger due to later floods and inundations. In general the soils consists of heavy clays with very often peat detritus in the soil profile. As a result of their sedimentary environment and age, these soils show progressive decalcification resulting in the lowest calcite contents.




Figure 2.8. Simplified soilmap of Zeeland, see text.

The excavation of the pool areas and the later land reconstruction works resulted in a disturbed soil profile with a high local variability in the subsoil. On the soil maps these soil types are given as associations of more than one soil type.

2.5.2 Channel ridges and creeks

One of the most remarkable features of the Zeeland landscape is the relief inversion of the tidal channels. These geomorphological features appear as ridges (channel ridges) in the landscape. The process of ridge inversion is depicted in figure 2.10. The inversion is caused by differences in compaction between the sandy channel deposits and the more




Figure 2.9. Digital elevation model of Zeeland acquired with laser altimetry. Boxes indicate the excerpts as shown in figures 2.11 (1), 2.12 (2), and 2.13 (3). (5x5 m ground resolution, 5 cm altitude resolution, more info at <http://www.ahn.nl/>)

compactable clay and peat deposits adjacent to the channels. The soils of channel ridges are generally more silty and sandy than the old polders and pool soils. Their subsoil often contains shell detritus.

Relief inversion is most pronounced for the Heartland channels, due to their initially larger depth (see figure 2.11). In areas that were reclaimed as accretion polders or from embanked tidal channels the inversion is usually much less or did not occur at all, due to the absence of the eroded peat. The ridge and creek patterns of these areas can be seen in figure 2.12, showing that large soil variability can occur over short distances due to the relatively abrupt transition from mud flat sediments to the more sandy creeks. During land reconstruction




Figure 2.10. Schematic cross section of the origin of ridge inversion of tidal channels. After Vos & van Heeringen (1997).

some of the elevated ridges were leveled with their surroundings and the material was used elsewhere to level the “hollebollig” land. Especially the leveling of the Heartlands of Schouwen-Duiveland was very extensive which can be seen in figure 2.9.

2.5.3 Accretion polders

The soil types of the accretion polders vary from sandy to heavy clays, depending on the marine and tidal environment before the reclamation. The soils often contain shell detritus and are not yet decalcified. In figure 2.13 the 18th century accretion polders in the western part of Noord-Beveland are shown (Rijkswaterstaat, 1971). A feature often encountered in such polders is the transition from more heavy clays close to the oldest dike towards less heavy clays near the newer dike. This, of course, is due to the slightly less turbulent marine sedimentary environment present at the time near the oldest dike. The small variability in sedimentary conditions resulted in gradual changes in clay content that are often not




Figure 2.11. Excerpt from digital elevation model showing pool areas

reflected in the soil map but only noticed during fieldwork.

2.5.4 Groundwater

According to the provincial water management plan (*provinciaal waterhuishoudplan*) three important shallow groundwater systems can be discerned; a salt/brackish system, a thick fresh water system, and a thin fresh water system (Nierop, 2000). The salt/brackish system and the thin freshwater system are dominated by upward seepage due to the low surface elevation relative to sea and fresh water surface levels, and are concentrated around the pool and other low areas. The thick freshwater system concentrates around tidal ridges and




Figure 2.12. Excerpt from digital elevation model showing channel ridges

higher elevated areas, the main direction of flow is determined by infiltration. The excess of water due to seepage and rainfall is removed towards the sea by a system of ditches and channels using pumping-stations. The phreatic level varies from close to the surface down to a depth of 2 m (Rijkswaterstaat, 1971). Regulation of the groundwater level is important for agriculture. As the input of freshwater cannot be controlled by fresh water inlets because of the generally brackish surface water, water levels are kept higher in winter to provide a buffer against drier periods in summer.




Figure 2.13. Excerpt from digital elevation model showing accretion polders

2.6 Summary and conclusion

Zeeland is located in a marine delta of the rivers Rijn, Maas and Schelde. It was the sea level rise following the end of the Weichselian ice age, that brought Zeeland within the marine realm. The soils of Zeeland can be regarded as relatively homogeneous and young. They were moulded in a continuous interaction of natural processes and human endeavour.

During the last depositional age, the Duinkerke period, three activities dominated soil formation: dike building and land reclamation, peat excavation, and large scale land reconstruction. The embankment of large areas removed these areas from the marine realm and fixed their condition. The division between the two major landscape types, Heartlands and

Newlands, is based on the embankment history. The relatively elevated Heartlands were mainly embanked in the Middle Ages as defensive measures against the sea, while the Newlands were reclaimed in later periods to acquire new agricultural land. As a consequence the soils of the Heartlands are mainly fairly heavy clays, while most of the Newlands are more sandy and silty. Especially in the Heartlands, large areas of peat were excavated, leaving behind an area of low agricultural value with an uneven surface and short range fluctuating moisture content. This, amongst other factors, led to major land reconstruction works, initiated after the two last catastrophic inundations of 1944 and 1953, in which the topsoil was removed and the subsoil leveled, after which the topsoil was replaced. This resulted in a variable subsoil with a relatively homogeneous topsoil.

Considering the fact that soils of Zeeland are in general developed in from marine clay deposits, it can be expected that a main source of geochemical variability is the varying clay content. The natural pattern of creek ridges and pool areas already creates relatively abrupt transitions. As local homogeneity or gradual variation may be further disturbed by the extensive human works, variability in soil composition is expected to be still higher than is directly evident from the soil map. Each (embanked) area with its own history, both sedimentary and human, may have its own pattern of soil variability. Given the fact that 80% of terrestrial Zeeland is used for agriculture, nearly 70% of which is arable land, human processes related to fertilisation and pesticide use are further expected to have influenced soil composition. This will result in elevated concentrations of so called “heavy metals” (Cd, Cu, Pb, and Zn) and persistent organochlorine pesticide residues. Finally, atmospheric inputs should also be considered as contributing to soil geochemistry.

3 Natural and anthropogenic patterns of covariance and spatial variability of minor and trace elements in agricultural topsoil

3.1 Introduction

Soil contamination is one of the major environmental issues within the Netherlands. The government, in its Third National Environmental Policy Plan (VROM, 1997), has called for a nationwide assessment of soil quality before the year 2005. The two tracks of the assessment include: 1; the stock of contaminated sites that need remediation, a job intended to be carried out to completion within the coming two decades; and 2; the general or “diffuse” soil quality. This second track of the assessment concentrates on making accessible and integrating available data on soil quality in support of soil protection policy and spatial planning.

As a consequence of stricter environmental legislation regarding building materials (VROM, 1999), municipal and provincial authorities recently have been putting much effort into the draft of so called soil pollution risk maps (in Dutch: BodemKwaliteitsKaarten or BKKs)(van der Gaast et al., 1998; van Lienen et al., 2000). These maps show the levels of priority chemicals relative to their legal thresholds in soil. A legally ascertained map allows dispensation of some of the clauses in the legislation regarding the effort needed to certify that soil that is transported to and from the area is legally “clean”. The data collected within the BKK-scope will also provide an important input into track 2 of the nationwide soil quality assessment.

In view of the large commitment of financial and human resources dedicated to soil quality assessment, the need was felt for a more scientific evaluation of soil quality and the benefit of soil quality maps, alongside the governmental tracks. One of the issues requiring further attention is the quality of soil quality maps, in respect of spatial variability and sampling procedures.

In print as: Spijker, J., Vriend, S.P., Van Gaans, P.F.M., 2005, “Natural and anthropogenic patterns of covariance and spatial variability of minor and trace elements in agricultural topsoil”, Geoderma.

In the Netherlands environmental soil surveys are usually based on sampling designs that use composite samples. The Dutch standard for soil sampling (NNI, 1999b) details the required procedure of such a design. This standard is based in part on agricultural practice and requirements for soil remediation research, and as such aims at a local rather than a regional scale. For reasons of comparability (the very first aim of a standard) and efficient use of existing data, the same design is also used in the BKK procedure and other more regional studies. For example in the geochemical soil survey for the province of Zeeland on which most of this thesis will be based, samples consisted of a composite of 15-20 subsamples taken from a field area of about 100 m · 100 m, at a density of approximately one sample per 3 to 9 km². The aim of this general Zeeland study is the (multivariate) characterization of the inorganic soil composition within the rural areas, both for naturally occurring and for anthropogenically enriched elements, like the so called “heavy metals” (Cd, Cu, Pb, Sb, Sn, and Zn). Given this context, expectations regarding the advantages of error reduction through composite sampling, based on experience and expert judgment in daily operation of remediation surveys, might be false.

The benefits of the reduction in variance through compositing depend on the spatial variation pattern and the spatial scale of interest. If interest is mainly in regional patterns compositing will only be useful when local variability is large, including in relation to analytical variance (i.e. variance associated with chemical analysis). If local features are of interest, compositing may be generally more relevant. The aim of this chapter is to estimate the variability related to spatial scale and sampling procedure for a wide range of elements in the topsoil of Zeeland. The hypothesis is that distinct spatial patterns of variability exist for groups of elements that are geochemically or anthropogenically related and whose variability depends on common factors and processes. The relative benefit of compositing may then be different for different groups of elements, which indicate that the quality of soil quality maps may have to be viewed differently depending on the desired application.

3.2 Materials and methods

3.2.1 Area

The chosen study area, the peninsula of Walcheren/Zuid Beveland, is located in the province of Zeeland, in the south-west of the Netherlands, see figure 3.1. The geological processes and human activities responsible for soil variability in this area are representative for Zeeland, and probably for similar deltaic areas around the world. The polder-landscape of the study area mainly consists of marine clay deposits (Duinkerke deposits) which are part of the Holocene Westland formation (Vos & van Heeringen, 1997). The Holocene alternation of marine clay deposits and peat deposits was caused by a sequence of transgressive and minor regressive events since the end of the last glacial period of the Pleistocene (see also chapter 2). The peninsula has a history of flooding and land-reclamation that continued until the 20th century. The area is divided into two major marine clay land-types based on their age of reclamation and relative altitude: Heartlands and Newlands.




Figure 3.1. Location map of the study area Walcheren/Beveland in Zeeland, the Netherlands, with sample sites.

The Heartlands originated as an alternation between salt-marshes and peat in the period between 500-1200 AD. During the various floods large tidal inlets formed that were subsequently filled with marine sand. The peat was excavated for the production of salt, resulting in pools which were later filled with heavy marine clays. These pool-clays are not very well suited for cultivation, but drainage and re-allotment improved this situation. A landscape developed with a variable, sometimes disturbed, soil profile of heavy clay and sometimes sparse peat fragments, cut by the sandy inlets. Due to the settling of the peat layers an altitude inversion occurred and the sand ridges are now about 1-1.5 m above the clay areas. The Newlands are of later origin than the Heartlands and in general were not used for peat excavation. The soil profile is less disturbed and consists of sandy marine clay deposits (Bazen, 1987). Soils of both land-types can be classified as fluvisols and are mainly used for farming.

3.2.2 Spatial variance and sampling theory

The total observed variance of a soil characteristic is in principal a summation of variance components that each can be attributed to a specific source. For example the variability in lime content in soil can, amongst others, be attributed to variation in the parent material and to variation in the extent of leaching with fresh water (Sposito, 1989). For the

heavy metals in topsoil an anthropogenic source of variance can be expected. Of course an unavoidable additional source is analytical error. In the simplest form of Analysis of Variance (ANOVA) the “within” variance obtained through replicate sampling within e.g; one type of parent material is considered as noise. If, from an F-test, the variance “between” different parent materials is significantly larger than the pooled “within” variance, parent material is concluded to be an additional source of variability.

In soil science, or in geological sciences in general, the total variance can often be viewed as being composed of spatial components. The well-known semivariogram displays the cumulative variance as a function of increasing distance (e.g. Journel & Huijbregts (1981)). A discrete version of this spatial variance pattern can be obtained through an ANOVA based on a hierarchical nested sampling design with different distances as subclasses or levels (Webster, 1985; Miesch, 1975).

The model-based semivariogram approach is most useful in cases where there is a continuous, gradual increase in variance with spatial scale, and the aim is to provide a geographically continuous assessment of the precision of interpolated maps. The design based ANOVA approach is most suitable in cases where a more stepwise increase in variance is expected, associated with spatial entities of a certain (approximate) size, such as agricultural fields, once flooded areas, reclaimed polders, etc. In this situation hierarchical spatial scales can be predetermined (de Gruijter & ter Braak, 1990). Besides, the ANOVA approach is more concise (Youden & Mehlich, 1937). Especially when using a so called unbalanced nested design (see figure 3.2 and Miesch (1976)) the number of samples required to assess the spatial variance structure of an area is reduced (Garrett, 1983). This design, that is often used and for which the data analysis is extensively described in the literature (Nortcliff, 1978; Oliver & Webster, 1986; Webster & Oliver, 1990; Oliver & Khayrat, 2001), was also chosen for the present study.




Figure 3.2. Stacking of levels of variation design in balanced (upper scheme) and unbalanced (lower scheme) designs (after Garrett (1983))

While the classical hierarchical unbalanced ANOVA (UANOVA) method has been proven useful in geochemical applications (Garrett & Goss, 1979; Garrett, 1983), there are at

present many other methods for estimating variance components for an unbalanced design, such as maximum likelihood, restricted maximum likelihood and a principal components based method (Searle et al., 1992; Khatree et al., 1997). The main disadvantage of the classical UANOVA, as seen by most authors, is the possibility of negative estimates for one or more of the variance components. Some go as far as calling this feature “awkward and embarrassing” (Searle et al., 1992), and therefore disapprove of the method. The advantages of (U)ANOVA, however, are clarity, simplicity, and robustness. More advanced methods such as Restricted Maximum Likelihood may be highly sensitive to whether or not an additional level of spatial scale is discerned (as in our case for example the division in Heartland and Newland) or not.

The possibility of negative estimates for variance components was not considered an insurmountable problem in this study. Negative estimates occur when the estimate for the between group variance is smaller than the estimate for the within group variance or, in spatial terms, when the calculated variance at a certain scale between units of a finer scale is smaller than the calculated variance within these finer scaled units. While in principle impossible (except for zonal features), a negative estimate can always occur if the real value is close to zero. Negative estimates thus can be substituted by zero. This method is generally accepted and performs well, also compared to more advanced methods (Pettitt & McBratney, 1993; Khatree et al., 1997).

The aim of this study is not to determine absolute values for the individual variance components of single elements, but to distinguish general patterns of spatial variance for groups of related elements. The simplicity and robustness of the UANOVA are therefore preferred over the sophistication, but sensitivity, of the more advanced methods.

3.2.3 Sampling and chemical analysis

For both Heartlands and Newlands eight locations were randomly selected in the marine clay deposits of the rural area of the peninsula resulting in 16 locations (see figure 3.1). Five local and two regional levels of variance were chosen. At each sample location single soil samples were collected at five specific distances, in a random direction from a base sample. The seven levels/distances thus investigated matched respectively the variability for a duplicate sample (1 m), at short distance (30 m), within the sample field (100 m), between neighbouring fields (300 m), at kilometre level (1000 m), between locations (regional), and Heartlands vs. Newlands (landtype). This sampling scheme is illustrated in figure 3.3. Heartland sand ridges were avoided to exclude large differences in soil type within one location.

The resulting 96 samples were taken with a hand auger from the plough layer. If this layer could not be distinguished, the samples were taken from the depth range 5-20 cm. After drying at 40 °C, grinding down to approximately 50 μm , and a hot aqua-regia digestion, the samples were analyzed by ICP-MS, together with 13 randomly selected duplicates at the analytical (within-sample) level. The following elements were determined: As, Ag, Au, B, Ba, Be, Bi, Ca, Cd, Ce, Cr, Cs, Cu, Dy, Er, Eu, Fe, Ga, Gd, Hf, Ho, La, Li, Lu,




Figure 3.3. Stacking of levels of the sample design for this research. For each landtype the design of one location is shown

Mg, Mn, Mo, Nb, Nd, Ni, Pb, Pr, Pt, Rb, Re, Sb, Sc, Se, Sm, Sn, Sr, Ta, Tb, Th, Ti, Tl, Tm, U, Y, Yb, Zn, Zr.

As pointed out by among others Walsh et al. (1997) in case of an acid leach such as aqua regia there are few recognized reference materials available that can be used for method validation. Moreover, the focus of this chapter is on variance and covariance, for which precision rather than accuracy is the priority issue. Accuracy was therefore assumed to be just as good as that attained in the general Zeeland study, for which part overlap with XRF data and comparison with ISE standards was available (van Dijk & Houba, 2000).

3.2.4 Data analysis

The data were first examined for analytical errors. From the duplicate analysis the relative error was calculated using

$$v = \frac{s}{\bar{x}} \cdot 100\% \quad (3.1)$$

Where v is the error relative to the overall mean \bar{x} and the standard deviation s . The latter calculated with

$$s = \sqrt{\frac{\sum_{i=1}^n (x_{i1} - x_{i2})^2}{2n}} \quad (3.2)$$

where the standard deviation is calculated from n duplicates with values x_{i1} and x_{i2} . Elements with large relative error were excluded from further statistical analysis.

Outliers are defined as values larger than the 3rd quartile plus 2 times the interquartile range, This procedure is equivalent to calculating the fences for box and whiskers plots according to Tukey (1977), except that for the inner fences the value of 2 was chosen

above the original value of 1.5 to compensate for possible positive skewness. This kind of Winsoring of the extremes was applied to prevent them from having undue influence on the UANOVA analysis. Outliers were replaced by a random sample from the upper (4th) quartile from the same distribution of the same element without outliers since the UANOVA poorly handles missing values. Listwise deletion obviously is not desired. The random replacement instead of a fixed maximum value ensures remaining variance also between outliers. We prefer this procedure over a logarithmic or similar transformation since the character of the statistical distribution of the variance components may also be scale dependent. In the case of just the sporadic outlier we do not assume that this replacement method adversely influences the distribution of the data.

For the UANOVA, the program of Garrett & Goss (1980) was used, to identify for each element which spatial scale/level contributes most to the overall variance and to show the relative contribution of each level. To identify groups of elements with similar spatial variability patterns a fuzzy c-means clustering (Bezdek et al., 1984; Vriend et al., 1988; Urbat et al., 1997) was performed on the UANOVA results. Due to the random selection of the analytical duplicates, independent of the UANOVA scheme, analytical variance is not incorporated in the UANOVA model. Moreover, it was felt that the analytical variance should not play an explicit role in the fuzzy clustering of the UANOVA results.

The assumed reduction of variance of a composite sample design, as used in the general Zeeland study, in respect to a single sample design as used here, was assessed with an F-test. Thirty-three samples from the general Zeeland study that cover the same area as that of the single sample design, although with a different sampling pattern (see figure 3.1), were selected and their overall variance was calculated. From the single sample design only the base sample and the 1000 m sample were selected, resulting in 32 samples in total. Using the 1000 m sample as well, the number and hence the spatial spread of locations, match those of the general Zeeland study.

Since the aim of this study is to distinguish general patterns of spatial variances of groups of related elements, it is also interesting to consider their covariance patterns. Of course these may also vary with spatial scale. Interest here is focussed on regional features so specifically the covariance at the overall (regional) level was considered. To this end a principal components analysis with varimax rotation (PCA-V; Davis (2002); Reimann et al. (2002)) was applied to the dataset. To provide sufficient dimensionality, defined as the ratio between the number of cases and variables (i.e. elements) used, a selection of representative elements was made for the PCA. Due to collinearity, it is not necessary to include all elements that are expected to be mainly related to one factor such as parent material. For instance, one or two rare earth elements (REE) will suffice instead of all REE, increasing the dimensionality. To check the assumption that the elements that are not included in the PCA-V are still explained by the factor model, the correlation with the PCA-V factorscores was studied. For the same reason of sufficient dimensionality, robust versions of PCA, that use only a subset of the cases, were not deemed adequate here. It was assumed that the Winsorising/replacement of the outliers sufficed to obtain a stable factor model.

3.3 Results and discussion

The analytical duplicates indicated that the elements Ag, Au, B, Mo, Nb, Pt, Re, Se, Ta, Tl, and U had much larger relative error ($v \gg 20\%$) than the other elements, indicating insufficient reproducibility. This was probably related to low concentrations in combination with a nugget type small scale variance. For Zr, with a relative error of 10.6%, also accuracy was reported by the laboratory to be lacking. All elements mentioned above were therefore not included in further analysis.

Trimming of outliers occurred in 2-4 cases for As, Cd, Sb, Sn, Zn, and respectively 7 and 10 cases for Cu and Pb. It was assumed, based on field experience, that at least part of these outliers result from atypical contamination by for example metal scrap, paint chips and leadshot, rather than being representative for the general soil geochemistry.

Reproducibility (v , see eq. 3.1) and summary statistics, after outlier replacement, are given in table 3.3. All concentrations of the reported elements are within detectable range for the ICP-MS, so no values below the detection limit occur.

Table 3.1. Statistical summary after outlier replacement of the geochemical data for the clayey top soils of Zeeland (n=96), sd = standard deviation, Q1= first quartile, Q3=third quartile, v relative error (see eq. 4.2). Ca, Fe, Mg, Ti, are given in to wt-%, other elements are in ppm.

	min	Q1	mean	median	Q3	max	sd	v
Ca	0.31	0.91	1.76	1.43	2.55	5.19	1.02	8.91
Fe	1.30	2.25	2.70	2.73	3.17	4.62	0.67	14.88
Mg	0.29	0.48	0.60	0.60	0.71	0.98	0.15	6.82
Ti	0.06	0.08	0.09	0.10	0.11	0.12	0.02	6.60
As	7.59	14.33	17.42	17.31	19.93	30.73	4.24	4.41
Ba	50.69	81.85	102.3	104.7	122.1	172.5	27.20	8.43
Be	0.67	1.05	1.30	1.33	1.54	2.02	0.33	5.36
Bi	0.09	0.17	0.21	0.21	0.25	0.38	0.06	3.08
Cd	0.15	0.29	0.35	0.34	0.40	0.60	0.09	5.13
Ce	32.40	46.33	53.06	53.88	60.05	70.91	9.53	4.35
Cr	31.01	48.12	58.41	58.84	70.83	87.69	13.78	8.81
Cs	2.04	3.65	4.94	5.06	6.20	8.04	1.56	4.84
Cu	7.62	11.66	16.45	15.04	19.49	33.23	5.72	7.10
Dy	1.71	2.49	2.87	2.93	3.25	3.80	0.52	3.40
Er	0.84	1.22	1.41	1.45	1.62	1.89	0.26	3.54
Eu	0.49	0.72	0.82	0.84	0.92	1.11	0.16	2.98
Ga	2.07	3.96	5.02	5.18	5.98	8.14	1.46	9.09

Continued on next page

Table 3.1 – continued from previous page

	min	Q1	mean	median	Q3	max	sd	ν
Gd	2.35	3.32	3.86	3.95	4.38	5.11	0.71	3.20
Hf	0.41	0.58	0.70	0.71	0.82	0.99	0.15	5.78
Ho	0.31	0.46	0.53	0.54	0.60	0.72	0.10	7.63
La	14.35	21.95	25.20	25.91	28.84	34.45	4.75	3.82
Li	12.85	21.98	29.38	30.55	36.04	50.52	9.01	5.34
Lu	0.10	0.15	0.17	0.17	0.19	0.23	0.03	5.81
Mn	141.6	233.1	308.2	291.6	370.8	542.0	95.19	8.24
Nd	14.16	19.92	22.91	23.26	26.02	30.44	4.14	4.09
Ni	7.60	15.19	18.62	19.31	22.61	29.83	5.26	4.31
Pb	11.54	21.51	27.03	26.96	30.70	50.19	7.61	3.07
Pr	3.63	5.29	6.05	6.12	6.88	8.09	1.11	5.46
Rb	32.45	50.97	66.36	66.00	79.90	105.4	18.56	5.50
Sb	0.27	0.35	0.42	0.42	0.47	0.65	0.08	9.89
Sc	3.70	5.98	7.50	7.62	8.90	11.53	1.91	8.51
Sm	2.76	3.90	4.46	4.53	5.08	5.89	0.81	3.88
Sn	1.16	1.74	2.19	2.07	2.50	3.95	0.61	4.41
Sr	37.12	53.23	69.86	65.25	87.95	124.3	21.19	3.51
Tb	0.33	0.47	0.54	0.55	0.61	0.73	0.10	3.48
Th	3.73	5.55	6.53	6.65	7.53	8.76	1.29	4.56
Tm	0.11	0.17	0.19	0.19	0.21	0.25	0.03	5.53
Y	8.32	12.13	13.86	14.19	15.76	18.76	2.59	5.09
Yb	0.73	1.08	1.22	1.27	1.40	1.66	0.23	5.48
Zn	34.86	61.62	74.26	73.12	86.32	131.6	20.53	5.21

Twenty representative elements were selected for the PCA-V, resulting in a dimensionality of 4.8, and the results are shown in table 3.2. For the elements not included in the PCA-V, correlations with the PCA-V factorscores are given in table 3.3 for the first factor. Correlations with scores of other factors were substantially lower, all less than 0.3. This confirms that the elements not included in the PCA-V are indeed highly correlated with one of the factors. The first rotated factor accounts for 47.6% of the total variance within the model. Elements with high factor loadings on this factor are mostly clay related natural elements, i.e. not influenced by human interference. The second factor accounts for 14% of the factor model variance and has high loadings for Ca and Sr and moderate loadings for As, Cu, Mg, and Mn. This factor represents the primarily natural variation in carbonate content of the soils. The elements As and Mn are possibly partly related to pH-dependent redox processes. The third factor has high loadings for the possibly anthropogenically influenced elements Cd, Pb, Sb, Sn, and to lesser extent Zn, Bi, and Cu. The factors have been

labeled clay, calcite, and anthropogenic respectively.

Table 3.2. Varimax rotated principal component factor loadings. Factor 1 (clay) explains 47.6% of the variance, factor 2 (lime) 14% and factor 3 (anthropogenic) 14.6%. Loadings between -0.25 and 0.25 are not shown

	Factor 1	Factor 2	Factor 3		Factor 1	Factor 2	Factor 3
As	0.40	0.46		Mg	0.87	0.44	
Bi	0.85		0.43	Mn	0.57	0.38	
Ca		0.90		Ni	0.88		0.32
Cd	0.26		0.74	Pb	0.33		0.67
Cu		0.43	0.40	Rb	0.96		
Fe	0.87			Sb	0.45		0.65
Ga	0.94			Sn			0.66
La	0.92			Sr		0.95	
Li	0.90	0.28		Ti	0.92		
Lu	0.91		0.27	Zn	0.68		0.50

Table 3.3. Correlations of concentrations of elements, not included in the PCA-V, with the the PCA-V scores on factor 1

	<i>r</i>		<i>r</i>		<i>r</i>
Ba	0.93	Ho	0.92	Sm	0.93
Be	0.93	Ce	0.93	Tb	0.92
Cs	0.94	Gd	0.93	Th	0.92
Dy	0.92	Nd	0.93	Tl	0.92
Er	0.92	Pr	0.93	Tm	0.91
Eu	0.94	Cr	0.95	Y	0.91
Hf	0.92	Sc	0.94	Yb	0.91

Figure 3.3 shows the results of the UANOVA graphically for some selected elements. These plots show the percentage by which each of the variance components contributes to the total variance. The percentages for all elements and levels are given in table 3.3. For most elements, the 1 m variance component is virtually equal to the analytical variance. Only for As, Bi, Pb, and Sn the analytical variance is about a factor 10 smaller.

Table 3.4. Relative percentages of the variance components for each level and their total. Negative variance components are denoted as 0%, region is regional component and N-H is landtype component, TVC is the total of variance components in (mg/kg)².

	1m	30m	100m	300m	1km	region	N-H	TVC
As	46.0	6.9	8.5	9.3	8.6	13.8	6.9	1.83·10 ¹
Ba	5.4	3.7	2.6	29.3	13.9	33.4	11.6	8.09·10 ²
Be	5.6	2.3	13.1	20.8	21.2	32.1	4.9	1.11·10 ⁻¹
Bi	7.7	0.3	3.7	30.4	24.8	2.2	30.8	4.00·10 ⁻³
Ca	1.2	11.1	0.2	41.9	0	10.7	34.8	1.46·10 ⁸
Cd	3.2	30.6	25.3	23.7	0	17.2	0	8.70·10 ⁻³
Ce	5.0	3.4	5.9	30.3	6.5	33.8	15.0	9.79·10 ¹
Cr	5.8	6.5	5.6	27.9	2.6	33.4	18.2	2.11·10 ²
Cs	3.7	3.4	4.7	37.4	11.9	27.2	11.8	2.70·10 ⁰
Cu	2.4	25.4	30.0	17.5	0	14.9	9.9	3.47·10 ¹
Dy	3.4	6.8	5.8	22.2	16.5	27.6	17.8	2.90·10 ⁻¹
Er	3.6	6.1	3.7	24.3	16.7	25.3	20.3	7.44·10 ⁻²
Eu	1.8	4.0	8.0	21.6	24.2	27.8	12.7	2.66·10 ⁻²
Fe	27.8	2.6	0	37.9	0	26.7	5.0	4.87·10 ⁷
Ga	9.8	0.6	15.4	17.2	8.2	32.4	16.4	2.31·10 ⁰
Gd	2.7	4.5	5.8	24.7	16.4	29.1	16.7	5.32·10 ⁻¹
Hf	11.2	3.5	4.9	25.0	17.7	20.3	17.5	2.26·10 ⁻²
Ho	3.9	5.1	6.7	21.5	19.7	25.3	17.8	1.03·10 ⁻²
La	6.5	1.4	4.0	32.1	6.3	34.8	14.8	2.32·10 ¹
Li	5.9	3.4	15.4	20.0	21.3	34.0	0	7.86·10 ¹
Lu	7.1	5.0	8.0	19.3	23.7	17.9	19.1	9.98·10 ⁻⁴
Mg	6.3	5.2	7.0	28.4	9.8	43.3	0	2.40·10 ⁶
Mn	14.1	4.5	0	57.3	0	24.1	0	9.61·10 ³
Nd	3.7	4.0	6.6	25.2	11.3	34.4	15.0	1.83·10 ¹
Ni	4.2	6.7	14.5	24.1	22.1	27.0	1.5	2.84·10 ¹
Pb	13.8	3.1	5.0	39.5	7.1	0	31.6	7.32·10 ¹
Pr	4.1	4.0	5.1	28.8	7.6	34.4	16.0	1.36·10 ⁰
Rb	4.7	4.6	3.2	41.5	4.9	28.1	13.1	3.81·10 ²
Sb	22.2	4.0	15.4	20.2	38.2	0	0	6.58·10 ⁻³
Sc	7.2	2.0	7.7	18.4	22.4	32.2	10.1	3.80·10 ⁰
Sm	2.2	4.9	6.6	24.0	13.6	30.9	17.8	7.13·10 ⁻¹
Sn	28.2	12.1	7.2	47.8	0	4.6	0.1	3.91·10 ⁻¹
Sr	1.8	14.7	3.1	33.8	0	18.1	28.5	5.67·10 ²

Continued on next page

Table 3.4 – continued from previous page

	1m	30m	100m	300m	1km	region	N-H	TVC
Tb	3.0	3.3	8.0	20.6	21.3	26.2	17.7	$1.06 \cdot 10^{-2}$
Th	4.0	2.4	6.0	30.4	12.4	27.2	17.5	$1.82 \cdot 10^0$
Ti	8.2	6.4	4.8	15.8	4.0	34.1	26.7	$3.03 \cdot 10^4$
Tm	4.9	5.8	5.9	20.1	21.6	23.2	18.5	$1.25 \cdot 10^{-3}$
Y	3.7	6.1	8.6	18.6	16.8	26.6	19.6	$6.99 \cdot 10^0$
Yb	8.2	2.3	6.9	18.8	24.3	21.2	18.5	$5.42 \cdot 10^{-2}$
Zn	3.3	8.2	5.7	37.9	8.2	6.1	30.7	$5.13 \cdot 10^2$

After fuzzy c-means clustering with varying numbers of clusters and visual inspection of all the plots, the output of the UANOVA was divided into 3 groups. The plots shown in figure 3.3 are representatives of each group. The first group (figure 3.3, Eu) shows a gradual increase of the variance with distance, with about equal contributions of the four highest levels. The second group (figure 3.3, Mg) shows a similar increase except that the contributions at the 1000 m level and the landtype level are lower than the 300 m and regional level. Both groups have in common that the major increase in variance is at the last four levels, i.e. distances larger than the field scale (300m). The third group (figure 3.3, Cd) shows a more irregular pattern which is different for each element within the group. They have in common that the largest increase occurs at the first three levels, i.e. within the field scale. The clusters are respectively labeled gradual 1, gradual 2, and irregular. The cluster memberships are given in table 3.5. The gradual 1 cluster contains the heavy REE, Sc and Y whereas gradual 2 contains the light REE and clay/feldspar related elements. The irregular cluster combines Ca and Sr, heavy metals, and redox sensitive elements.

Figure 3.5 gives an overview of all the measured elements and their variance patterns. For clarity only the cumulative variance is plotted against the distance. From this figure the difference between the three types of variation patterns becomes clear. In general the curves of gradual 2 are less smooth as those of gradual 1, with a clear inflection at the 300 m level. The elements of the irregular cluster show a steep ascending curve for short distances which flattens towards larger distances. This underlines that the major part of the total variance for these elements occurs at the field scale.

Table 3.6 shows the relation between the covariance structure, derived from the PCA-V and the three spatial variance patterns. In general the elements with high loadings on the clay PCA-V factor, or that are highly correlated with this factor, are grouped in the first two clusters (gradual 1 & 2). For these elements the contribution of the small scale variance to the total variance is relatively small. Thus observed regional variance, whether or not composite samples are collected, will indeed be largely caused by regional features. It is remarkable that while both light and heavy REE have high correlations with the clay factor in the PCA-V, they cluster separately based on their spatial variance patterns.

All the elements with high loadings on the anthropogenic and redox/lime factor of the




Figure 3.4. Contribution of the hierarchical variance components for Eu to the total variance (bars) and cumulative variance (line). The levels/distances represent the variability for a duplicate sample (1m), at short distance (30m), within the sample field (100m), between neighbouring fields (300m), at kilometre level (1000m), between locations (regional), and Heartlands vs. Newlands (landtype). Eu represents an element for gradual 1. *continued on next page*

PCA-V have an irregular variance pattern. For these elements small scale variation is of more importance than regional variation. The irregularity of the variation patterns for the anthropogenically influenced elements is attributed to the more erratic effect of human processes on soil geochemistry, in this area mostly related to farming, compared to the processes related to the deposition of marine clays.

The irregularity of the elements with high loadings on the redox/lime factor may be related to the geohydrological situation in Zeeland. Large parts of the area are below sea level and subject to localized upward seepage of groundwater. It can be assumed that due to redox




Figure 3.4. *continued* Mg represents an element for gradual 2. *continued on next page*

processes at the phreatic level As and Mn become locally enriched. The elements Ca and Sr have a high correlation coefficient of 0.9. These elements, together with Bi, Pb, and Zn, have a noticeable difference in concentration between Heartlands and Newlands (figure 3.1). Their local variation pattern might be influenced by soil-water interactions, similar to the redox sensitive elements. Post depositional leaching of Sr bearing calcite from the marine deposits results in irregular depletion. It is also known that alternating oxidizing/reducing conditions enhance calcite leaching (van den Berg & Loch, 2000).

Figure 3.6 depicts the ratio of overall variance as compared between the composite samples from the general Zeeland study and the single base and 1000 m samples from this variance oriented study. It shows a relatively symmetric distribution ranging between 0.22 and 0.93 around a median ratio of 0.57. This corresponds to the observed range in cumulative field-scale contribution to the total variance (figure 3.5). Indeed the anthropogenically influenced




Figure 3.4. *continued* Cd represents an element for irregular.

elements with the highest local variance components show the largest variance reduction due to compositing. For these elements compositing will be necessary for local estimates of concentration levels. A disadvantage of compositing is that with the reduction of variance covariances are also attenuated. For regional studies compositing is only worthwhile when local variance is large compared to regional variance, and the number of subsamples needed for adequate reduction is highly dependent on this ratio of local to regional variability. We conclude that for surveying anthropogenic influence a larger number of subsamples is needed than for determining natural backgrounds, for which single samples may often be adequate.

Table 3.5. Fuzzy cluster memberships. Elements are sorted according to dominant cluster and membership.

	gradual 1	gradual 2	irregular		gradual 1	gradual 2	irregular
Ho	0.91	0.07	0.03	Cs	0.16	0.64	0.21
Tb	0.89	0.08	0.03	Mg	0.20	0.63	0.17
Y	0.81	0.14	0.05	Th	0.32	0.57	0.11
Dy	0.81	0.15	0.04	Ga	0.30	0.54	0.16
Tm	0.79	0.14	0.07	Rb	0.13	0.52	0.35
Eu	0.77	0.17	0.06	Sm	0.41	0.52	0.07
Yb	0.74	0.16	0.10	Ti	0.30	0.48	0.21
Er	0.72	0.19	0.08	Li	0.37	0.45	0.17
Lu	0.71	0.17	0.12	Pb	0.15	0.18	0.67
Gd	0.68	0.27	0.05	Zn	0.16	0.19	0.66
Hf	0.64	0.22	0.14	Sn	0.14	0.23	0.63
Sc	0.53	0.36	0.10	Ca	0.16	0.23	0.60
Ni	0.47	0.36	0.17	Mn	0.14	0.32	0.54
Be	0.43	0.43	0.13	Sr	0.20	0.30	0.49
Pr	0.09	0.85	0.05	As	0.25	0.32	0.44
Nd	0.11	0.85	0.04	Fe	0.19	0.39	0.42
Ce	0.09	0.84	0.06	Bi	0.35	0.25	0.41
La	0.12	0.79	0.09	Sb	0.32	0.29	0.39
Ba	0.18	0.75	0.07	Cu	0.27	0.34	0.39
Cr	0.15	0.72	0.13	Cd	0.26	0.35	0.39

3.4 Conclusions

The covariance pattern as revealed with a varimax principal component analysis (PCA-V), groups the trace and minor elements into a clay, anthropogenic, and calcite factor.

An unbalanced analysis of variance (UANOVA), in combination with a fuzzy c-means clustering of the variance components, reveals three general patterns of spatial variability that clearly group geochemically related elements. The third, irregular pattern, is characterised by a relatively large contribution of local variance components to the overall variance. This cluster is limited to the elements on the anthropogenic and calcite factors and Fe, and is interpreted as being the result of human and/or geohydrological processes.

Composite samples reduce local variance. However, for most elements the regional variance predominates. In regional studies the extra effort of taking composite samples should be carefully weighed against the relatively small amount of noise caused by local variance. For the so called “heavy metals” (Cd, Cu, Pb, Sb, Sn, and Zn) small scale variability is

Table 3.6. Relation between covariance structure (PCA-V) and the clustered UANOVA variance patterns

PCA-V factor	Spatial variance types (clusters)		
	gradual 1	gradual 2	irregular
Clay	Be, Dy, Er, Eu Gd, Hf, Ho, Lu Ni, Sc, Tb, Tm Y, Yb	Ba, Ce, Cr, Cs Ga, La, Li, Mg Nd, Pr, Rb, Sm Th, Ti	Bi, Fe, (Mn), (Zn)
Anthropogenic			Cd, Cu, Pb, Sb Sn, Zn
Calcite			As, Ca, (Cu), Mn Sr, (Zn)

an important source of the total variation. For these elements moderate compositing, i.e. a small number of subsamples, may be useful in regional studies. The 15-20 subsamples as used in the general geochemical survey (see next chapters) appear to be adequate for the “irregular” elements, and more than adequate for the other elements.




Figure 3.5. Summary of cumulative spatial variance patterns as estimated with UANOVA. The cumulative variance as percentage of the total variance on the y-axis is plotted against the distance on the x-axis. For x-axis scale see figure 3.3. The distances of the regional and landtype levels were in these figures arbitrary set at 3000 m and 5000 m. The order of the plot is from high memberships in the gradual 1 cluster (Ho to Be), via high memberships in the gradual 2 cluster (Pr to Li) to high memberships in the irregular cluster (Pb to Cd).




Figure 3.6. Variance ratios between composite samples and single samples (F-statistic; circle), and its 95 % confidence interval (line). 33 samples selected from the general Zeeland study employing composite sampling, and 32 samples from the single sample design, base and 1000 m sample.

4 Sampling and analyses for a regional environmental soil geochemical survey

4.1 Introduction

To manage and safeguard the environment one needs primary knowledge about its status and condition. For terrestrial life, including the human species, the biosphere is strongly dependent upon the chemistry of the earth's surface layer. Soils form a fundamental component of this layer, hence knowledge of the distribution of major and trace elements in soils, and their mutual relations in spatial as well as attribute space, are of major concern for understanding our environment. The basic factors that determine the composition of soils are: 1) geogenic/pedogenic origin, 2) the vertical redistribution through soil forming processes, and 3) the ubiquitous spread of anthropogenic substances as the consequence of diffuse entry into the soil. In geochemistry the original composition is often assessed by means of a baseline, while divergence from this baseline may then be related to human influence.

For the Dutch situation this baseline is expressed by in legal background threshold values (Dutch: "Streefwaarden") that form the backbone of the Dutch legal framework of permissible limits for soil quality and soil remediation (see also chapter 1). These legal values were derived in a search for true soil background values in a survey of natural areas, under the assumption that these were not significantly influenced by man and thus could be considered as relatively clean. Van der Meent et al. (1990) calls this method of acquiring background values disputable since it is hard to establish criteria for a "relatively clean area". Their suggestion is to use preanthropogenic deposits for determining background concentrations or use scientific deduction. They also consider that natural background concentrations for the Dutch situation do not exist. However, Bölviken et al. (1996) pointed out that pristine, preanthropogenic material can be found and sampled everywhere. A geochemical survey with an appropriate sampling design can provide both influenced and pristine soil material. A comparison between the two then offers a diagnosis tool on the state of a major environmental quality: the soil geochemistry.

This chapter focuses on how to obtain a consistent, high quality, and province-wide dataset

that contains both information about pristine soil composition, i.e. the natural composition, and information on how the soil is influenced by human processes. From such a dataset geochemical baselines can be derived as well as a quantification of the imprint by man. The use of the dataset for a baseline, hence as a reference dataset, demands that the methods are extensively described so a proper evaluation of their efficacy and a comparison with other data can be made. Of course, accuracy and precision are of the utmost importance. To discern a large variety of different processes at an adequate level of significance, through relations between soil attributes and in geographical space, it is necessary to keep the level of noise as low as possible. One of the major concerns is to correct for bias caused by having different surveys spaced in time. Due to necessary re-calibrations and unavoidable changes in laboratory practice and apparatus, such biases are virtually unavoidable. To anticipate to these changes a new method of parametric levelling is used based on sets of between survey duplicates (BSD) and linear regression.

From the complete dataset with all measured soil attributes a -still sizable- set of soil chemical attributes was selected, based on their affirmed high accuracy and precision, and no or only random missing values. Thus the desired georeferenced, consistent, and high quality dataset for the entire area studied was created. This dataset will be optimal for statistical analysis and environmental assessment for legal authorities. The remainder of the attributes measured may, when used with prudence and competence, still provide important additional insight into the soil geochemical patterns and processes.

4.2 Material and methods

4.2.1 Study area

This research focuses on defining and distinguishing natural and anthropogenic geochemical patterns in soil. The province of Zeeland was chosen for this study because it is an agricultural area with a relatively small variability in soil types that are of a young, Holocene, age. Because of the low permeability, high adsorption capacity of the clayey soils, and their relatively young age, vertical transport through natural soil forming processes or anthropogenic input is expected to be minimal. Thus, the presence of a pristine soil layer at depth may be reasonably assumed.

The province of Zeeland belongs largely to the Holocene estuary at the outlet of the rivers Rijn, Maas and Schelde*, see figure 4.1. It consists of the mainland of Zeeuws-Vlaanderen, bordering Belgium, and several islands/peninsulas that have a long and rich history of land reclamation and inundation. Inundation of reclaimed land was partly related to the rise of average sea-level in the first half of the last millennium. However, both natural and human induced flooding have occurred until recent times. Examples are the strategic flooding of Walcheren near the end of the Second World War, and the great storm flood of 1953 that inundated large parts of Zeeland. The history of land reclamation through dike building is

*In this thesis Dutch names are used for the rivers Rhine, Meuse and Scheldt respectively




Figure 4.1. Topography of Zeeland, showing names of peninsula's and areas (italic), and some cities (bold)

complicated, resulting in a geographically complex polder-landscape.

The Holocene geology and pedology of Zeeland have been extensively described in chapter 2 of this thesis. In summary the major soil types of Zeeland are immature marine clay deposits ranging from heavy clays (clay fraction $>25\%$) to sandy tidal inlet and tidal flats (clay fraction $<7\%$). The Zeeland landscape is divided into Heartlands, the oldest and first reclaimed areas with often very heavy pool-clays, and Newlands, younger areas with more sandy clay soils. The elevation of the field level, relative to the Dutch average sea level NAP, ranges from about 1.5 m for channel ridges to -1.3 m for the pool clay areas. Land use is mainly agricultural and in the Newlands is characterised by its larger scale and intensive

artificial drainage.

4.2.2 General research approach

Examples of geochemical baseline studies are common in modern geochemistry (see for example de Vos et al. (1996); Gregorauskiene & Kadunas (1997); Salminen & Tarvainen (1995); Burenkov et al. (1999); Cocker (1999); Rapant et al. (1999); Ferreira et al. (2001); Singh et al. (2001)), chiefly based on practices of exploration geochemistry. As Hale & Kühnel (1997) explain, the classical purpose of exploration geochemistry, finding metal-liferous ore bodies, can be transferred to the inventory of contaminated soils. Yet, there are important differences, e.g. looking for a single element stationary ore body as compared to a possibly mobile multi-element contamination of soil and the huge difference in time scales of both phenomena. Also for natural geochemical anomalies the underlying process or cause is often known or can be deduced from mineral relationships, while this is much less the case for contaminated soils. The causes of contamination, the interaction with soil constituents, and the relation between contaminating elements are often not well understood. Still the basic geochemical techniques are the same and approaches known in geochemistry are now considered very useful in the environmental field (Nikkarinen et al., 1996; Selinus & Esbensen, 1995).

Examples of environmental geochemical studies include the assessment of a background by using overbank samples (Swennen & van de Sluys, 2002). Here older, deeper, fluvial deposits are used as pristine material, and the relatively modern top sediments to assess the anthropogenic influence. This approach of using deeper sediments as pristine material has also been successfully applied in other studies and different soil types (Blaser et al., 2000; Wilcke et al., 1998; Sterckeman et al., 2004). The specific choices for sampling and analytical procedures to be used in this study are based on several considerations.

Most of the earlier mentioned geochemical baseline and/or environmental studies are generally limited to a small group of trace elements, notably the so called heavy metals (Cd, Cr, Cu, Ni, Pb, Zn) and As. Although those usually constitute the environmental priority elements, broadening the analytical focus with major components like Al_2O_3 , Fe_2O_3 , CaO, and trace elements such as Rb, Ga, Sr, and the rare earth elements (REE), will result in a better understanding not only of the human effect but of the natural composition and relations in the soil as well.

To make efficient use of the multi-element approach, detection limits for all elements should be sufficiently low so as to stringently reduce the number of measurements below the detection limit, since they will weaken and complicate data interpretation and statistical analysis.

To serve as a reference dataset, sampling and analytical methods should be at least partly in line with the current Dutch practice in assessments for soil sanitation permits and soil quality statements. This chiefly implies that composite samples are collected and that chemical analysis is preferably based on an aqua regia leach.

Current state of the art analytical techniques with high throughput that allow measurement

of a large number of major, minor, and trace elements including the REE, are Inductively Coupled Plasma-Mass Spectrometry (ICP-MS) and X-ray Fluorescence (XRF). Especially ICP-MS ensures low detection limits for a wide range of trace elements. While XRF provides the total content of an element in a sample, ICP-MS is always based on (partial) sample decomposition. The efficacy of the decomposition or leach depends on the chosen medium and exact procedure followed, and on the binding of the specific element in the sample matrix. Hot aqua-regia sample decomposition was chosen to comply with to the desired comparability with Dutch standard procedures. A known limitation of this method is the underestimation of concentrations of some metals due to resistance to the acid attack and resulting in a partial leach (Scancar et al., 2000; Tsolakidou et al., 2002). The overlap in elements between XRF and ICP-MS can be used as an supplementary check on analytical quality, and for some elements the difference between results may give some indication on their mineralogy.

One of the difficulties in soil analysis is sample inhomogeneity, which can result in low reproducibility and high variability of the measured values. Therefore homogenisation of the samples needs special attention, both at the field sampling stage and when subsampling for chemical analysis, in order to reduce this source of variability as much as possible. Variance due to inhomogeneity is dependent on grain size and mineralogy, and thus varies from element to element. Especially trace elements that form their own minerals, such as Zr, may suffer from the so called nugget effect (see also Chapter 3), which can only be reduced by increasing the weight of the analytical sample or reducing the grain size by milling.

As analytical variance and minor inaccuracies are inevitable properties of measured data, knowledge of the size of these errors is needed to allow any geochemical data interpretation. Therefore, for both the XRF and ICP-MS methods several duplicate samples and standards were analysed together with the samples. These included the between survey duplicates (BSD) for the purpose of levelling, i.e. eliminating the bias caused by separately analysing the different surveys that were widely spaced in time. Moreover, data were screened for anomalous values at all stages of the analytical procedure.

4.2.3 Field sampling

The samples were collected in 4 field surveys between 1996 and 2003. Sampling followed a geographically and pedologically stratified random procedure, with a density varying from 1 location per 1 km² in the first survey, up to 1 per 10 km² in the later surveys. The variation in density was based on improved knowledge about local and regional variation (see chapter 3) and a change from a local to a more regional focus with later surveys. The exact sample location within a randomly selected target site depended on permission of the landowner.

Focus was on rural areas, so built up zones and the dunes were excluded. Obvious potential point sources of pollution (industrial sites, junk yards, grave yards, orchards) were avoided as well. Soil types covering only a small area of the province were also disregarded.

The sample locations are indicated on figure 4.2 and detailed in table 4.1. At each location a visually homogeneous area of about $0.1 \cdot 0.1 \text{ km}^2$, two composite samples were collected: one from the topsoil, i.e. the 1-25 cm depth interval (topsoil sample), and one from a vertically homogeneous soil layer, a layer visibly consisting of one lithological unit, at greater depth (generally about 35-80 cm - subsoil sample). The subsoil samples were assumed to represent the pristine soil composition. The top layer usually comprised the plough layer which is vertically homogenised by agricultural practice. It was assumed that human activity primarily influenced this soil layer. Soil formation, which is still in incipient stages, is assumed not yet to have chemically altered the topsoil significantly, so its natural bulk chemical composition can be compared to the subsoil layer. Where no clear top layer existed, like on meadows, the top sample was taken from the 1-25 cm interval.

Table 4.1. Year, region and number of sample locations.

Survey	Year	region	locations
1	1996	Noord Beveland, Tholen St. Philipsland	68
2	1999	Walcheren, Zuid Beveland	60
3	2000/2001	Schouwen Duiveland Zeeuws Vlaanderen	107
4	2003	Tholen, Zuid Beveland	35
total			270

Drillings were made at 15-20 random sites within the location using a hand auger. Three to four spoonfuls of soil were collected from each drilling at the desired depth interval, to create two composite samples. Thus with an equal volume of sample material from each drilling a manageable yet representative composite sample of about 300 g was assured that could be easily homogenised after drying. Based on the results of chapter 3, suggesting that just a small number of subsamples may be preferred over a single sample for anthropogenic influenced elements, the chosen number of subsamples appeared more than sufficient.

For each location a brief description was made of the soil profile, the surrounding area, and any detail that could be of relevance to interpreting the analytical results. The location of the sampled field was recorded to a 100 m accuracy for georeferencing.

4.2.4 Sample preparation

The samples were dried at 40-45 °C to constant weight. The clayey nature of the samples required disintegration of the dried samples with a porcelain pestle and mortar to about 5 mm grain size. A Jones splitter was used to obtain representative subsamples for further analysis. For each sample about one third was used for analyses and the remainder was stored in a sample archive. This archive was later used for the levelling of the data (see




Figure 4.2. Sample locations per survey

§4.2.8) and is intended as a reference for future research.

4.2.5 Chemical analyses

One aliquot, from the one-third sample, of about 30g was pulverised to $<50\mu\text{m}$, using a Herzog HSM-F36 Tungsten-Carbide disk-mill, for homogenisation and subsequent ICP-MS and XRF analyses. For each technique a separate quantity of material was taken out of the aliquot. The use of the tungsten-carbide disk mill contaminated the samples with elements like W, and Co.

The samples for the ICP-MS were digested with aqua-regia according to a standardised

method of the Dutch Geological Survey. A subsample of 250 mg of the milled 30 g aliquot was leached with 6 ml ultra-pure hydrochloric acid and 2 ml ultra-pure nitric acid and digested in a closed Teflon vessel in a microwave oven (Milestone 1200 Mega) for 13 min at 600 W and 17 min at 300 W. After a cooling period of 10 to 20 minutes another 2 ml of nitric acid was added and the hydrochloric acid was removed through evaporation. For this part, due to changes in the laboratory, an evaporator (survey 1 and 2), a sand bath (survey 3), and an aluminum block (survey 4) were used as heat sources in the different surveys. The solution was quantitatively transferred into polyethylene flasks and diluted to 50 ml with demineralised water. From this volume an aliquot was 5 times diluted, together with an internal standard (In) and submitted for ICP-MS analysis.

For the XRF method 10 g of the milled sample was mixed with wax tablets and pressed in a Herzog HTP-S and analysed, using α -correction for major elements and Compton backscatter correction for trace elements. A new XRF apparatus was installed between surveys 2 and 3.

Table 4.2 lists the elements that were determined by each method. Due to enhancements in the laboratory and the possibility to analyse more elements the set of parameters was extended after the first survey, and to a smaller extent after the third survey.

4.2.6 Precision and accuracy

During all stages of the chemical analysis data integrity was closely watched by means of comparison between XRF and ICP-MS results, duplicate comparison, drift plots, etc. Given the large number of attributes determined, these a priori attempts to eliminate gross analytical errors focused on the key attributes like the major oxides and the priority heavy metals.

Precision of the analytical results within a survey was estimated through duplicate analysis of some 15% of the ICP-MS samples, and about 4% of the XRF samples as this method is known to be more precise. For each ICP duplicate a new subsample was decomposed by aqua regia, for each XRF duplicate a new subsample was ground and pressed into a tablet.

The precision was calculated with

$$s = \sqrt{\frac{\sum_{i=1}^n (x_{i1} - x_{i2})^2}{2n}} \quad (4.1)$$

$$v = \frac{s}{\bar{x}} \cdot 100\% \quad (4.2)$$

where v is the error relative to the mean (\bar{x}) of the duplicates and s is the standard deviation with $(x_{i1} - x_{i2})$ the difference between the n sets of duplicates.

Reference ISE samples (van Dijk & Houba, 2000), that were treated in the same way as the samples, were used to check on accuracy. Comparison of laboratory performance with the recommended values of the international standards for XRF results is relatively straightforward, as these represent total concentrations. Since for some elements the aqua regia

Table 4.2. Attributes analysed per survey and method, numbers indicate the different surveys

Element	XRF	ICP-MS	Element	XRF	ICP-MS	Element	XRF	ICP-MS
Ag		2,3,4	La	2,3,4	all	Ta		2,3,4
Al	all	4	Li		2,3,4	Tb		2,3,4
As	all	all	Lu		2,3,4	Th	all	2,3,4
Au		all	Mg	all	2,3,4	Ti	all	2,3,4
B		2,3,4	Mn	all	2,3,4	Tl		2,3,4
Ba	all	all	Mo		all	Tm		2,3,4
Be		2,3,4	Na	all	4	U	all	2,3,4
Bi		2,3,4	Nb	all	2,3,4	V	all	4
Ca	all	2,3,4	Nd	2,3,4	2,3,4	Y	all	all
Cd		all	Ni	all	all	Yb		2,3,4
Ce		all	P	all	4	Zn	all	all
Cr	all	all	Pb	all	all	Zr	all	2,3,4
Cs		all	Pr		2,3,4			
Cu	all	all	Pt		all			
Dy		2,3,4	Rb	all	all			
Er		2,3,4	S	all				
Eu		2,3,4	Sb		all			
Fe	all	2,3,4	Sc	2,3,4	2,3,4			
Ga	all	all	Si	all				
Gd		2,3,4	Se		all			
Hf		2,3,4	Sm		2,3,4			
Ho		2,3,4	Sn	2,3,4	all			
K	all	4	Sr	all	all			

ICP-MS data represent only a partial leach, this should be taken into consideration when evaluating their accuracy. As pointed out by a.o. Walsh et al. (1997) in this case there are few recognized reference materials available that can be used for method validation. Where possible the overlap between XRF and ICP-MS will be used to assess the typical element-specific recovery.

Accuracy a was determined as the bias b between average measured value \bar{x} and the reference value relative to r using

$$b = \bar{x} - r \qquad a = \frac{b}{r} \qquad (4.3)$$

4.2.7 Outliers

Outliers can occur due to an anomalous sample or due to analytical error. In the first case the sample may originate from a location with high concentrations of a certain element or elements. These elevated concentrations, either caused by natural or human processes, can denote a local soil feature, another possibility is the incidence of small, non visible, amounts of allochthonous material like flakes of paint, pieces of leadshot, metal scraps, etc. in a subsample (van Gaans & Vriend, 1996). The grinding and homogenisation of the sample may then contaminate the entire sample. Outliers caused by analytical error can have various causes that are usually hard to account for. A repeated analysis can disclose such outliers but for many elements the overlap between the ICP-MS and XRF can be used. If an ICP-MS analysis shows an outlier for a certain element while the XRF does not, it is very likely that this outlier is an analytical effect.

Outliers were defined as values larger than the 3rd quartile value plus several times the interquartile range according to

$$x > P_{75} + f \cdot (P_{75} - P_{25}) \quad (4.4)$$

This procedure is equivalent to calculating the fences for box and whiskers plots according to Tukey (1977). In this formula P denotes a percentile and x is the measured value. The factor f , usually set at 1.5 for inner fences, was set to 2, making allowance for a possible skewness. If equation 4.4, with $f = 2$, is applied to the normal distribution virtually no outliers exist, while for a log normal distribution data about 5% will be identified as outliers caused by the large tail of a theoretic log normal distribution. Outliers were not removed from the dataset but marked as such so they can be treated separately.

4.2.8 Parametric levelling

Since the time span between the 4 surveys was several years (see table 4.1) it is inevitable that small, but systematic shifts will have occurred between the different data subsets. These shifts are caused by changes in procedures, fluctuations in calibration and/or performance of the analytical instruments etc., and are unavoidable in laboratories that keep improving analytical performance. Instead of consolidating the method for years it is better to anticipate these changes so expected bias can be compensated. The shifts, although small and within acceptable analytical precision for individual parameters, can have a negative influence on multivariate relations. For example, covariability may shift from one subset to another, thereby weakening the relation in the overall dataset. Artificial groups can also result from these artifacts (Rutten, 2004).

To level, or eliminate, these small differences so called *Between Survey Duplicates* (BSD) were used. One survey was chosen as the reference and about 15 to 20 samples from this reference survey were reanalysed in the analytical batches during the other surveys. The other surveys were then corrected by means of shift and multiplier values based on an




Figure 4.3. Principal concepts of levelling. The circles show the deviation from the $y = x$ line, as a result of shifting. The triangles show the same data after correction

ordinary least square linear regression equation derived from the set of BSD.

This method was chosen above well known methods like regression of quantiles (Daneshfar & Cameron, 1998; Darnley et al., 1995) or correction based on the reference samples used. The advantage of this method is that the levelling is based on a, for the area, representative range of measured values.

Figure 4.3 shows the procedure of the BSD levelling for an example dataset. Due to a systematic shift the data measured in a second batch are shifted away from the $y = x$ line. If the outcome of the analyses only differs due to analytical variability then the points would

be equally spread around this line. From the shifted data cloud a equation can be calculated using simple linear regression in which the newly measured concentration is expressed as a function of the referenced sample. Using the regression slope a and intercept b the unlevelled data can then be corrected with

$$C_{levelled} = \frac{C_{unlevelled} - b}{a} \quad (4.5)$$

in which $C_{levelled}$ denotes the corrected value based on the original $C_{unlevelled}$ value.

Differences between the duplicates are the summation of random analytical noise and the bias caused by the analytical shift. To assess the validity of the parametric levelling the correlation between all the BSD pairs after levelling is calculated. A low value for r indicates a large contribution of random analytical noise and thus a less accurate bias correction.

This parametric levelling does not remove analytical bias completely but reduces different biases towards one bias, which must still be checked against (international) reference samples. The correction procedure can influence the precision of the separate surveys only when the regression coefficient a is much larger or smaller than 1, as this inflates or deflates overall variance within the corrected survey data.

4.3 Results and interpretation

4.3.1 Precision

After analysis of each batch, the data were first visually inspected by various kinds of graphs like box-whisker plots, scatter plots, drift plots, histograms and where possible scatter plots of elements both determined by XRF and ICP-MS. These were to reveal drift and differences between (subparts of) batches, outliers and erroneously measured samples. If necessary the samples were re-analysed and erroneous data records were corrected.

Results for the average analytical precision, based on within batch analytical duplicates, are shown in tables 4.3 (ICP-MS) and 4.4 (XRF). As an additional check on the representativity of the duplicates, the mean of all the duplicates was compared to the overall mean for all the data; no relevant differences were found.

Most of the elements measured by ICP-MS have precision below 10% while B, Bi, Cu, Sn, Ti, and Zn have a slightly higher error (table 4.3). The elements Ag, Au, Hf, Nb, Pt, Se, Ta, Tl, and Zr are not within acceptable range. These latter are mostly elements at low concentrations, that do not occur in the alumino-silicate mineral matrix but only in their own trace minerals (nugget effect), some of which are also incompletely digested by aqua regia. Considering the aqua regia microwave digestion method, these precision results are within the ranges expected.

The XRF, due to the principles of the method, has much lower analytical error, in general

Table 4.3. Average precision for ICP-MS, based on within batch analytical duplicates. The precision of the 51-58 duplicates (except Au, Ta, Tl, Pt, which have 40-42 duplicates) is given as standard deviation v relative to the mean of the duplicates \bar{x} .

	\bar{x} [ppm]	v [%]		\bar{x} [ppm]	v [%]		\bar{x} [ppm]	v [%]
Ag	0.06	15.7	Ga	4.49	7.8	Sb	0.27	8.2
As	11.88	4.8	Gd	3.17	4.6	Sc	6.52	9.5
Au	0.04	117.2	Hf	0.33	19.8	Se	0.66	27.4
B	40.72	12.1	Ho	0.40	5.2	Sm	3.74	4.7
Ba	70.66	9.4	La	21.29	5	Sn	1.33	13.0
Be	0.97	7.5	Li	22.76	7.8	Sr	71.17	4.8
Bi	0.15	11.0	Lu	0.13	8.2	Ta	0.03	48.9
Ca	18055	7.7	Mg	4775	6.9	Tb	0.44	5.2
Cd	0.21	10.0	Mn	206.79	6.9	Th	5.49	7.1
Ce	42.86	5.1	Mo	0.48	8.9	Ti	630.32	11.5
Cr	40.56	6.9	Nb	0.99	21.3	Tl	0.34	23.1
Cs	3.53	4.4	Nd	19.46	4.9	Tm	0.14	7.0
Cu	9.91	10.7	Ni	15.76	5.2	U	0.93	6.5
Dy	2.27	5.2	Pb	17.15	6.8	Y	10.73	6.3
Er	1.06	6.3	Pr	5.11	4.6	Yb	0.90	7.4
Eu	0.65	4.7	Pt	0.01	57.1	Zn	53.77	12.2
Fe	19458	6.4	Rb	45.97	5.8	Zr	12.85	17.3

less than 2% (table 4.4). Precision for S, Cu, La, Nd and Sc is relatively poorer, but still in line with the precision levels found for the ICP-MS. The results for Sn, Th, and U are not within an acceptable range. These are trace elements that also primarily form their own trace minerals but are partly incorporated in alumino-silicates as well. The trace minerals dominate the overall concentration and cause the observed nugget effect for the XRF results. The aqua-regia extractable alumino-silicate part results in much lower, but more precise concentration levels as obtained by ICP-MS. The in general high quality of the XRF analysis as obtained here, points out that the homogenisation and splitting steps were adequate to remove sample inhomogeneity.

4.3.2 Accuracy

Accuracy is given in tables 4.5 for the XRF and 4.6 for the ICP-MS. The overall negative bias and much lower accuracy for the ICP-MS, as compared to the XRF, are as expected. The ICP-MS accuracy for the elements Cd, Tl, and Zn are surprisingly good. For the XRF, accuracy is in general sufficient but Nd, Sn, and U show high values indicating poor accuracy and a positive bias.

Table 4.4. Average precision for XRF, based on within batch analytical duplicates. The precision of the 8-9 duplicates is given as standard deviation v relative to the mean of the duplicates. The average concentration of the duplicates \bar{x} is given in [wt-%] for oxides and S and [ppm] for other elements.

	\bar{x}	v [%]		\bar{x}	v [%]		\bar{x}	v [%]
Al ₂ O ₃	8.21	0.83	As	13.37	1.88	Sc	8.48	9.96
CaO	2.83	3.78	Ba	278.74	1.35	Sn	4.61	46.19
Fe ₂ O ₃	3.67	0.57	Cr	75.41	3.55	Sr	97.26	2.17
K ₂ O	1.85	0.48	Cu	10.02	9.36	Th	7.14	27.78
MgO	0.85	1.84	Ga	9.40	3.05	U	2.03	25.73
MnO	0.04	1.37	La	25.91	6.23	V	66.08	1.98
Na ₂ O	0.76	0.97	Nb	10.42	2.77	Y	20.65	2.80
P ₂ O ₅	0.17	1.67	Nd	25.52	7.44	Zn	51.34	1.99
S	0.08	9.77	Ni	17.82	3.45	Zr	328.09	1.92
SiO ₂	71.03	0.57	Pb	22.65	5.28			
TiO ₂	0.47	0.70	Rb	76.53	1.16			

Table 4.5. Accuracy of the XRF analyses. For calculation of accuracy a , see equation 4.3

	a		a		a
Al ₂ O ₃	0.06	As	-0.01	Sc	0.19
CaO	-0.03	Ba	0.04	Sn	0.46
Fe ₂ O ₃	0.04	Cr	0.04	Sr	0.04
K ₂ O	0.01	Cu	0.01	Th	-0.20
MgO	-0.09	Ga	-0.01	U	0.82
MnO	0.04	La	-0.22	V	-0.02
Na ₂ O	0.19	Nb	0.07	Y	0.16
P ₂ O ₅	-0.02	Nd	0.49	Zn	-0.07
SiO ₂	-0.07	Ni	-0.01	Zr	0.01
TiO ₂	-0.06	Pb	-0.05		

4.3.3 Outliers

The number of outliers for the complete dataset of top and subsoil, as defined by equation 4.4 for both the ICP-MS and XRF method are given in table 4.7.

The elements with the highest number of outliers are Pb, Sb, and Zn for ICP-MS and Pb, U, and S for XRF. Only for Cu, Mn, Pb, and Zn in the topsoil, and for Ca and Sr in the subsoil common outliers recur. Many others appear to be method, rather than sample-

Table 4.6. Accuracy of the ICP-MS analyses. For calculation of accuracy a , see equation 4.3

	a		a		a
Ag	-0.60	Ga	-0.61	Sb	-0.55
Al	-0.73	K	-0.81	Sc	-0.42
As	-0.14	La	-0.33	Se	-0.96
B	-0.73	Li	-0.15	Sn	-0.63
Ba	-0.85	Mg	-0.17	Sr	-0.42
Be	-0.53	Mn	-0.21	Th	-0.22
Bi	-0.98	Mo	-0.77	Tl	0.05
Ca	-0.33	Na	-0.97	U	-0.37
Cd	0.03	Nb	-0.84	V	-0.38
Ce	-0.21	Nd	0.11	Y	-0.52
Cr	-0.50	Ni	-0.09	Zn	0.05
Cu	-0.27	P	-0.16	Zr	-0.97
Fe	-0.27	Pb	-0.22		

specific, indicating analytical error.

The elements Pb, Zn, and Cu are known contaminants and the presence of samples with anomalous high concentrations is a first indication of anthropogenic altering of the topsoil. As no comparison between ICP-MS and XRF is possible for Cd, P, and S this could be true for these elements as well. The anomalies for CaO and Sr in the subsoil may be related to a concentration of shell fragments which were regularly found in the subsoil during sampling.

In view of the size of the total dataset and the regional objectives of the overall survey, these outliers are negligible and can be simply omitted in the regional characterisation.

4.3.4 Parametric levelling of data

Survey 3 was chosen as the reference survey for the BSD because in this sampling round the largest number of samples were analysed. Hence, the smallest possible number of samples needed to be corrected. Moreover, the new XRF apparatus was already in use for this survey. The choice for survey 3 therefore also facilitates comparison with future data. An example of the levelling is given in figure 4.4 in which Ga, as measured on the XRF, is levelled towards the reference.

The overall correlations between the BSD after the parametric levelling are given in tables 4.8 and 4.9. From these tables it is clear that for the XRF the correlations are higher than for the ICP-MS. For the XRF the elements Sn, Th, and U have low correlations, which is in accordance with their low precision (table 4.4). In general the correlations are between

Table 4.7. Number of outliers, only elements with at least 2 outliers are shown. Common outliers are samples which are both outlier for the XRF and ICP-MS. The overall summation shows how many samples exist with outliers for at least 2 elements.

ICP-MS			XRF			Common	
element	top	sub	element	top	sub	top	sub
Ca	6	4	CaO	0	2	0	2
Cd	6	3	–			–	–
Cu	3	3	Cu	3	3	2	0
Mn	3	1	MnO	6	1	3	0
–			Na ₂ O	1	3	–	–
Nb	1	4	Nb	0	0	0	0
–			P ₂ O ₅	9	5	–	–
Pb	10	0	Pb	8	0	4	0
Pr	2	0	–			–	–
Sb	7	8	–			–	–
Sn	5	2	Sn	1	0	0	0
Sr	1	5	Sr	1	5	1	4
Ta	3	2	–			–	–
Tl	4	3	–			–	–
U	3	4	U	11	2	1	1
Zn	9	4	Zn	3	0	3	0
–			S	12	6	–	–
overall	8	6		6	2		

0.8 (Na₂O , S, and Ba) and 1. The ICP-MS has overall lower correlations between 0.6-0.9, which are also in general accordance with the precision data (table 4.3).

The high correlations for most of the elements show that the use of ordinary least squared linear regression is acceptable as regression method. For each element plots like figure 4.4 were used to check the regression and confirm that they were first order. No exceptions for the latter were found. Tables 4.8 and 4.9 also show that the regression slopes are generally close to 1 and for those elements with a relatively low slope (e.g. Nb, Ta, and U for the ICP-MS and Sn, and Th for the XRF) also the correlation is low. These elements have also a low precision (table 4.3 and 4.4).

4.3.5 Comparison between XRF and ICP-MS

As expected the use of an aqua-regia digestion often resulted in apparently lower concentrations and also lower precision for some elements for the ICP-MS due to their poor leach-

Levelling XRF Ga




Figure 4.4. Example of levelling of Ga for the XRF. The circles show the data after correction while the other symbols show the unlevelled data for the different batches. The solid line is the $y = x$ line while the dashed lines are the regression lines for each batch.

ability in the medium. Table 4.10 shows the correlation and absolute differences between the techniques. To illustrate the relations between both techniques figure 4.5 shows scatter plots of a selection of elements. From this figure the great difference and low correlations of Ba and Zr are apparent. Elements like Cr and Ga are, despite their good reproducibility of respectively 6.9% and 7.8% by ICP-MS, apparently systematically incompletely leached from the soil matrix but still show a good correlation.

The comparison also gives indications of the typical mineralogy, especially for presumed heavy minerals like barite, cassiterite, zircon, garnet and more general amphiboles and

Table 4.8. Results of the levelling for the XRF data. r is the correlation coefficient and a is the slope of the linear regression

Element	r	a	Element	r	a
Al ₂ O ₃	1	1.02	La	0.97	1.13
CaO	0.97	0.98	Nb	0.96	0.96
Fe ₂ O ₃	1	1.01	Nd	0.93	0.86
K ₂ O	1	0.99	Ni	0.98	0.98
MgO	0.96	0.93	Pb	0.96	0.98
MnO	0.89	0.82	Rb	1	1
Na ₂ O	0.75	0.61	Sc	0.95	1
P ₂ O ₅	0.99	0.99	Sn	0.19	0.28
S	0.82	0.59	Sr	0.91	0.83
SiO ₂	0.90	0.92	Th	0.46	0.64
TiO ₂	0.97	1	U	0.51	0.81
As	0.98	0.98	V	0.99	1.01
Ba	0.84	0.83	Y	0.96	0.97
Cr	0.93	0.91	Zn	0.97	0.91
Cu	0.99	0.94	Zr	0.96	1
Ga	0.99	0.98			

pyroxenes. Figure 4.5 shows that heavy mineral associated elements like Ba, Zr, Ga, Cr, Sc, and to a lesser extent Ni, have lower values by ICP-MS than on the XRF. Apparently, they only partially leach in an aqua regia solution (Scancar et al., 2000; Tsolakidou et al., 2002). For Ba and Zr poor correlations exist while for Cr, Ga, and Ni they are better. The main distinction is that Ba and Zr are very poorly leached by aqua regia (respectively about 25% and 3%), and mainly occur outside the alumino-silicate matrix. Their total concentration as measured by XRF thus will not be correlated to Al₂O₃, in contrast to their ICP measured aqua regia leached part.

A good example of the benefits of comparing two methods is Zn. This shows unexpected overall enrichment in the ICP-MS extract compared to the XRF together with several ICP-MS outliers. After consultation with the laboratory staff it was inferred that the extensive reconstruction activities in the building over the past years apparently have generated a Zn enriched fine dust that continues to contaminate the air quality in the laboratory.

4.3.6 Parameter selection and summary statistics

Selection of parameters for a final, optimal dataset was based on the following criteria: no, or only sporadic missing values, satisfactory precision and accuracy, and in case of

Table 4.9. Results of the levelling for the ICP-MS data, r is the correlation coefficient and a is the slope of the linear regression

Element	r	a	Element	r	a
As	0.86	0.94	Mn	0.78	0.80
Ba	0.74	0.70	Mo	0.46	0.35
Be	0.89	0.90	Nb	0.42	0.40
B	0.67	0.81	Nd	0.79	0.76
Bi	0.85	0.82	Ni	0.90	0.90
Ca	0.88	1.04	Pb	0.76	0.88
Cd	0.83	0.90	Pr	0.80	0.83
Ce	0.82	0.86	Rb	0.90	0.90
Cr	0.91	0.92	Sb	0.77	0.81
Cs	0.85	0.90	Sc	0.72	0.70
Cu	0.64	0.77	Sm	0.81	0.82
Dy	0.76	0.72	Sn	0.64	0.75
Er	0.76	0.72	Sr	0.92	0.94
Eu	0.68	0.73	Tb	0.77	0.72
Fe	0.87	0.89	Th	0.76	0.81
Ga	0.87	0.88	Ti	0.68	0.64
Gd	0.77	0.73	Tm	0.73	0.68
Ho	0.77	0.73	U	0.66	0.63
La	0.84	0.85	Y	0.84	0.85
Li	0.87	0.86	Yb	0.72	0.64
Lu	0.62	0.53	Zn	0.67	0.72
Mg	0.89	0.92			

overlap between XRF and ICP-MS: preferably optimal precision and accuracy, but also compatibility with the standard Dutch practice of aqua-regia extraction.

The first criterion results in rejection of a number of parameters that were not analysed in the first survey: Ag, B, Be, Bi, Li, Sc, Ta, Tl, and most of the REE. Of these Ag, Ta and Tl also suffer from poor precision. Other elements that are rejected because of unsatisfactory precision/accuracy are Au, Pt, and Se. In the comparison between XRF and ICP-MS for optimal precision and accuracy, XRF is favoured for most elements, except for Sn, which has a higher reproducibility by ICP-MS. For Ba which has a large difference between ICP-MS and XRF (table 4.10), the ICP-MS analyses were chosen to be more compatible with other Dutch research. For most other elements that have satisfactory precision for both XRF and ICP-MS, method bias is relatively small (4.10).

Table 4.11 summarises the 31 parameters selected for the optimal dataset ($n=270$). The

Table 4.10. Differences and correlations between analysed elements both on XRF and ICP-MS. The difference is calculated as median XRF minus median ICP-MS. When elements are measured as oxides on the XRF, the ICP-MS values are recalculated to oxides [wt-%] and marked with a *.

	difference median [ppm]	correlation		difference median[ppm]	correlation
As	1.84	0.95	Ni	2	0.94
Ba	214.10	0.60	Pb	3.51	0.86
Ca*	0.99	0.95	Rb	31.82	0.96
Cr	39.56	0.92	Sc	2	0.85
Cu	0.67	0.87	Sn	2.14	0.33
Fe*	0.87	0.96	Sr	26.72	0.93
Ga	4.13	0.95	Th	2.67	0.64
La	3.73	0.86	Ti*	0.36	0.87
Mg*	0.06	0.93	U	1.25	0.42
Mn*	0.01	0.93	Y	9.77	0.87
Nb	9.22	0.27	Zn	-5.38	0.90
Nd	5.79	0.85	Zr	316.09	-0.16

main advantages of this rigorous selection, maximising precision and minimizing the number of missing values, is that the dataset is optimally suited for multivariate statistical analysis. When necessary, the complete dataset can be used for cross reference. This will especially be relevant for B, Be, Bi, Li, Sc, and the REE that only failed the first criterion.

Table 4.11. Selected elements

XRF	Al ₂ O ₃	CaO	Fe ₂ O ₃	K ₂ O	MgO	MnO	Na ₂ O
	P ₂ O ₅	S	SiO ₂	TiO ₂			
	As	Cr	Cu	Ga	Nb	Ni	Pb
	Rb	Sr	V	Y	Zn	Zr	
ICP-MS	Ba	Cd	Ce	Cs	La	Sn	Sb

A statistical summary of the dataset is given in tables 4.12 and 4.13 for respectively top and subsoil. Outliers as identified in section 4.3.3 were not included in the calculations, as they are considered not to represent the regional population at which the Zeeland soil survey aims, or to be analytical error. The tables provide the minimum (0 percentile), 25 percentile, median (50 percentile), 75 percentile and maximum (100 percentile) from which the distribution of the data can be derived. Also the mean and standard deviation are given.

Figure 4.6 gives an illustration of the type of distributions found, in the form of histograms




Figure 4.5. Comparison between selected elements from the XRF and ICP-MS, concentrations in ppm.

for some selected elements from topsoil samples. In geochemistry it is often mentioned that data, especially for minor and trace elements, show log-normal tendencies but this is not always the case (Reimann & Filzmoser, 1999; Allegré & Lewin, 1995). For the Zeeland data neither straightforward normality nor log normality should be assumed. Calculated skewness indeed tend to be positive for most minor and trace elements, and their histograms show long tailed distributions, but they are not identical to true lognormal distributions. The inverse analogue holds for the major elements like Al_2O_3 and Na_2O that generally tend to show more normal distributions.




Figure 4.5. *continued* Comparison between selected elements from the XRF and ICP-MS, concentrations are in ppm

4.4 Discussion and conclusions

Soils form a major part of our environment and assessing the degree to which they are influenced by man is of great importance. When assessing at individual concentrations of elements it is often difficult to distinguish anthropogenic effects from natural variation. Establishing a bivariate or multivariate geochemical baseline, that incorporates a major part of natural variability, is an accepted method to derive background concentrations. Divergence from this background might then be caused by human disturbance. To quantitatively assess which components are affected by human activities, a consistent regional dataset is

needed where special attention is paid to quantifying data quality. This chapter documents the construction of such a dataset for the province of Zeeland.

The sampling design was aimed at sampling present-day soil and pristine soil concentrations by sampling an influenced topsoil layer and an assumed pre-anthropogenic deeper subsoil layer. Based on earlier results (see chapter 3) the minimum sampling density of 1 sample per 1-10 km² was regarded sufficient for obtaining a regional overview of the soil chemical composition in Zeeland. The analytical methods used (ICP-MS with aqua regia digestion and XRF on pressed powder tablets) and the broad group of elements measured (major elements like Al₂O₃ and Fe₂O₃, anthropogenically affected elements like Cd, Pb, Sb, trace elements like the REE) resulted in a data set of 87, partly overlapping, parameters. Each group of elements is thought to provide specific information on natural variation, human influence, or their interaction, as will the comparison between top and subsoil.

The comparison allowed by the overlap in measured parameters between ICP-MS and XRF shows a low extraction efficiency by aqua regia especially for elements like Ba, Zr, Ga, Cr, Sc, and Ni, which probably are partly related to the presence of heavy minerals containing these elements to various degrees. The parameter overlap can also be used to determine if outliers are true anomalous samples or caused by analytical error. The elements Cu, Mn, Pb, and Zn display outliers in the topsoil data common to both techniques, which is an indication that these are (anthropogenic) anomalies.

Precision was estimated by analysing about 15% of the ICP-MS samples and about 4% of the XRF samples in duplicate. The smaller number of duplicates for the XRF was considered sufficient since this method is known to be far more reproducible than the aqua-regia ICP-MS method. The better XRF reproducibility is indeed confirmed by the results for analytical variance, which is in general lower for the XRF (0.5-10% relative to the mean) than for the ICP-MS (5-13% relative). Exceptions are Th, U, and Sn as measured by XRF and Ag, Au, Nb, Pt, Se, Ta, Tl, and Zr for ICP-MS; their low precision is ascribed to nugget effects. The analytical variability is generally smaller than the local sampling variance as found in chapter 3, and therefore adequate for the desired regional evaluation of soil concentrations. The number of outliers found in the data also appears quite acceptable, as from a total of 270 samples a maximum of 10 outliers for Pb is found.

In this study the levelling of the data based on Between Survey Duplicates (BSD) proved to be a good method to eliminate subsurvey biases. The fact that BSD levelling uses a set of samples that has the full range of element composition and concentration present, and is thus representative of the total survey, makes this method preferable to corrections solely based on analytical standards. The levelling uses ordinary linear regression, which was confirmed to be adequate in view of high correlations between the BSD duplicate pairs and a visual check on the linearity of the relation. The higher precision for the XRF also leads to higher correlations for the parametric levelling.

Based on the precision and accuracy a selection was made of 31 parameters that were measured in all surveys, and hence for virtually all locations and soil depths. This selection is thus optimised for multivariate statistical analysis by maximizing data quality and minimizing the number of missing values. This optimal dataset is limited compared to all the

data analysed, but still comprises a sufficient subset of major, minor, and trace elements for environmental assessment of the soils of Zeeland. It is intended to form the core for all multivariate statistical and geochemical analysis, from which cross reference to the complete data set can still be made.

Summary statistics show that for most of the parameters neither a normal nor a lognormal distribution should be assumed. Although most major elements tend to a more or less normal distribution, while most minor and trace elements are positively skewed, deviations from the standard type distributions are apparent. Suspected contaminants like Cd, Sb, As, Pb and Zn are more skewed than other trace elements.

The data collection exercise documented in this chapter shows that, the use of state of the art analytical methods and appropriate use of international standard samples and within survey, as well as between surveys, sample duplicates are all prerequisites for obtaining a true reference dataset suitable for environmental and geochemical assessment of soil within a regional context.




Figure 4.6. Histograms for selected elements and oxides analysed in the topsoil samples. Oxides in [wt-%], other elements in [ppm]

Table 4.12. Statistical summary, with outliers omitted, of the geochemical data for the topsoil of Zeeland. Q1=25 percentile, Q3=75 percentile, sd=standarddeviation. Components in the selected dataset are shown with a dot. The exclude field gives the main reason not to include the component in the selected dataset: i=incomplete/not analysed in all surveys, p=unsatisfactory precision, l=unsatisfactory leveling, o=other method preferred (see text).

	method	min	Q1	median	Q3	max	mean	sd	exclude	unit
•	Al ₂ O ₃	3.41	6.95	8.23	9.59	12.53	8.31	1.85	-	[wt-%]
•	CaO	0.37	1.48	2.50	4.40	8.92	3.08	1.97	-	[wt-%]
	Ca	2678	8639	62600	22480	62600	17219	12525	i	[ppm]
	as CaO	0.37	1.19	1.71	3.09	8.62	2.37	1.72	-	[wt-%]
•	Fe ₂ O ₃	0.91	2.80	3.62	4.41	6.54	3.67	1.12	-	[wt-%]
	Fe	5188	14446	19003	23677	37369	19531	6556	i	[ppm]
	as Fe ₂ O ₃	0.73	2.03	2.67	3.33	5.25	2.75	0.92	-	[wt-%]
•	K ₂ O	1.24	1.72	1.88	2	2.33	1.87	0.21	-	[wt-%]
•	MgO	0.03	0.61	0.84	1.05	1.69	0.84	0.31	-	[wt-%]
	Mg	1122	3700	4676	5500	8566	4709	1375	i	[ppm]
	as MgO	0.18	0.60	0.76	0.89	1.39	0.76	0.22	-	[wt-%]
•	MnO	0.01	0.03	0.04	0.04	0.07	0.04	0.01	-	[wt-%]
	Mn	96.43	177.54	220.10	265.71	402.57	226.05	57.79	i	[ppm]
	as MnO	0.01	0.02	0.03	0.03	0.05	0.03	0.01	-	[wt-%]
•	Na ₂ O	0.48	0.67	0.74	0.81	1.04	0.74	0.11	-	[wt-%]
•	P ₂ O ₅	0.11	0.16	0.18	0.20	0.35	0.18	0.04	-	[wt-%]
•	S	0.01	0.08	0.09	0.11	0.17	0.09	0.03	-	[wt-%]
•	SiO ₂	55.43	68.65	73.48	77.23	92.17	72.84	6.70	-	[wt-%]

Continued on next page

Table 4.12 – continued from previous page

	method	min	Q1	median	Q3	max	mean	sd	exclude	unit	
•	TiO ₂	xrf	0.24	0.41	0.48	0.55	0.71	0.48	0.09	-	[wt-%]
	Ti	icp-ms	357.81	577.24	653.32	721.81	939.14	643.01	108.47	i	[ppm]
	as TiO ₂	icp-ms	0.06	0.09	0.11	0.12	0.15	0.10	0.02	-	[wt-%]
	Ag	icp-ms	0.04	0.06	0.07	0.08	0.13	0.07	0.02	i	[ppm]
•	As	xrf	6.95	12.62	15.86	18.35	30.11	15.70	3.83	-	[ppm]
	As	icp-ms	5.45	10.98	13.98	16.22	25.60	13.89	3.59	o	[ppm]
	Au	icp-ms	0	0.01	0.02	0.05	0.15	0.03	0.03	p	[ppm]
	B	icp-ms	11.70	28.78	39.31	47.05	69.30	38.94	12.35	i	[ppm]
	Ba	xrf	221.80	274.73	291.32	304.74	334.70	289.78	20.95	o	[ppm]
•	Ba	icp-ms	27.91	57.02	72.69	85.91	126.32	72.56	19.61	-	[ppm]
	Be	icp-ms	0.31	0.76	0.96	1.14	1.73	0.97	0.28	i	[ppm]
	Bi	icp-ms	0.07	0.14	0.17	0.22	0.36	0.18	0.06	i	[ppm]
•	Cd	icp-ms	0.18	0.26	0.29	0.33	0.56	0.30	0.06	-	[ppm]
•	Ce	icp-ms	24.07	36.71	41.45	46.87	56.86	41.46	6.70	-	[ppm]
•	Cr	xrf	49.60	69.30	78.10	87.73	103.50	78.14	12.60	-	[ppm]
	Cr	icp-ms	14.12	31.37	37.37	45.27	66.22	38.38	10	o	[ppm]
•	Cs	icp-ms	0.81	2.86	3.59	4.47	7.92	3.75	1.28	-	[ppm]
•	Cu	xrf	5.16	10.36	12.93	16.67	27.44	13.87	4.76	-	[ppm]
	Cu	icp-ms	5.10	9.41	11.35	14.02	23.14	11.95	3.64	o	[ppm]
	Dy	icp-ms	1.23	2.02	2.37	2.66	3.45	2.34	0.44	i	[ppm]
	Er	icp-ms	0.56	0.93	1.11	1.24	1.59	1.09	0.20	i	[ppm]
	Eu	icp-ms	0.34	0.60	0.70	0.81	1.05	0.70	0.14	i	[ppm]
•	Ga	xrf	3.92	8.03	9.39	11.03	14.71	9.55	2.14	-	[ppm]
	Ga	icp-ms	1.49	4.07	5.19	6.31	9.78	5.31	1.60	o	[ppm]

Continued on next page

Table 4.12 – continued from previous page

	method	min	Q1	median	Q3	max	mean	sd	exclude	unit
	Gd	1.72	2.83	3.27	3.75	4.76	3.27	0.60	i	[ppm]
	Hf	0.05	0.17	0.25	0.36	0.64	0.28	0.14	i	[ppm]
	Ho	0.21	0.36	0.42	0.47	0.61	0.42	0.08	i	[ppm]
	La	14.24	22.30	25.84	29.58	42.41	26.01	5.49	i	[ppm]
•	La	11.32	18.34	20.99	23.67	30.20	20.92	3.54	-	[ppm]
	Li	5.71	16.73	22.64	27.08	47.89	22.75	8.05	i	[ppm]
	Lu	0.08	0.12	0.15	0.16	0.21	0.14	0.03	i	[ppm]
	Mo	0.10	0.28	0.37	0.56	1.35	0.45	0.24	l	[ppm]
•	Nb	6.41	9.18	10.58	11.67	13.83	10.46	1.72	-	[ppm]
	Nb	0.33	0.58	0.91	1.40	2.59	1.02	0.51	i	[ppm]
	Nd	18.20	22.88	26.20	28.03	34.10	25.73	3.61	i	[ppm]
	Nd	10.74	17.09	19.90	22.58	27.01	19.68	3.43	i	[ppm]
•	Ni	5.70	14.46	18.34	22.38	35.87	18.90	5.92	-	[ppm]
	Ni	5.11	12.97	16.27	19.66	31.16	16.81	4.86	o	[ppm]
•	Pb	14.73	21.95	25.77	30.57	47.88	26.82	6.52	-	[ppm]
	Pb	10.34	17.62	20.67	25.09	37.42	21.68	5.28	o	[ppm]
	Pr	2.78	4.41	5.11	5.67	8.23	5.04	0.91	i	[ppm]
	Pt	0	0.01	0.01	0.01	0.03	0.01	0.01	p	[ppm]
•	Rb	42.37	65.62	75.03	87.13	116.24	77.33	14.83	-	[ppm]
	Rb	11.99	34.93	43.97	55.43	83.64	45.56	14.09	o	[ppm]
•	Sb	0.19	0.28	0.32	0.36	0.60	0.33	0.07	-	[ppm]
	Sc	3.50	6.48	8.05	10.20	14.80	8.25	2.27	i	[ppm]
	Sc	2.21	4.90	6.14	7.40	11.34	6.20	1.75	i	[ppm]
	Se	0.10	0.30	0.70	1.09	1.89	0.72	0.46	p	[ppm]

Continued on next page

Table 4.12 – continued from previous page

	method	min	Q1	median	Q3	max	mean	sd	exclude	unit
	Sm	2.14	3.27	3.75	4.18	5.10	3.71	0.63	i	[ppm]
	Sn	0	2.60	3.94	5.82	9.63	4.15	2.18	p	[ppm]
•	Sn	0.78	1.51	1.80	2.05	3.46	1.79	0.45	-	[ppm]
•	Sr	48.67	78.37	92.40	119.31	191.14	101.68	30.17	-	[ppm]
	Sr	23.06	48.35	65.40	100.03	176.58	76.18	35.76	o	[ppm]
	Ta	0.01	0.02	0.03	0.05	0.13	0.04	0.03	i	[ppm]
	Tb	0.24	0.39	0.46	0.52	0.69	0.46	0.09	i	[ppm]
	Th	0	6.69	8.02	8.84	12.48	7.72	1.88	p	[ppm]
	Th	2.92	4.72	5.44	6.10	7.87	5.38	1.02	l	[ppm]
	Tl	0.11	0.24	0.32	0.41	0.75	0.34	0.13	i	[ppm]
	Tm	0.08	0.13	0.15	0.17	0.22	0.15	0.03	i	[ppm]
	U	0.49	2.03	2.18	2.48	3.40	2.21	0.51	p	[ppm]
	U	0.57	0.82	0.95	1.07	1.55	0.97	0.20	i	[ppm]
•	V	21.70	51.82	64.97	79.73	137.35	67.20	20.06	-	[ppm]
•	Y	11.47	18.34	20.68	23.05	29.05	20.69	3.24	-	[ppm]
	Y	5.68	9.43	10.96	12.32	15.67	10.85	1.91	o	[ppm]
	Yb	0.48	0.82	0.95	1.06	1.40	0.94	0.17	i	[ppm]
•	Zn	20.70	44.08	53.19	65.42	99.35	54.87	14.51	-	[ppm]
	Zn	31.84	50.82	58.36	66.83	106.83	59.69	12.84	o	[ppm]
•	Zr	178.12	296.08	325.53	366.48	500.13	329.30	54.45	-	[ppm]
	Zr	3.56	8.32	10.87	13.77	23.91	11.38	3.97	o	[ppm]

Table 4.13. Statistical summary, with outliers omitted, of the geochemical data for the subsoil of Zeeland. Q1=25 percentile, Q3=75 percentile, sd=standarddeviation. Components in the selected dataset are shown with a dot. The exclude field gives the main reason not to include the component in the selected dataset: i=incomplete/not analysed in all surveys, p=unsatisfactory precision, l=unsatisfactory leveling, o=other method preferred (see text).

	method	min	25-prec.	median	75-perc.	max	mean	sd	exclude	unit
• Al ₂ O ₃	xrf	2.58	5.54	7.23	9.30	13.29	7.49	2.51	-	[wt-%]
• CaO	xrf	0.52	2.80	3.94	5.27	10.70	4.05	1.90	-	[wt-%]
• Ca	icp-ms	2961	15277	22772	30486	63723	24390	12716	i	[ppm]
as CaO		0.41	2.10	3.14	4.20	8.77	3.36	1.75	-	[wt-%]
• Fe ₂ O ₃	xrf	0.51	2.08	2.99	4.13	7.39	3.20	1.40	-	[wt-%]
• Fe	icp-ms	2123	11304	16094	23571	43678	17680	8425	i	[ppm]
as Fe ₂ O ₃		0.30	1.59	2.26	3.31	6.14	2.49	1.18	-	[wt-%]
• K ₂ O	xrf	0.96	1.54	1.69	1.94	2.61	1.75	0.31	-	[wt-%]
• MgO	xrf	0.04	0.46	0.77	1.06	1.67	0.80	0.40	-	[wt-%]
• Mg	icp-ms	1022	3332	4565	5803	8615	4571	1742	i	[ppm]
as MgO		0.17	0.54	0.74	0.94	1.40	0.74	0.28	-	[wt-%]
• MnO	xrf	0.01	0.02	0.03	0.04	0.07	0.03	0.01	-	[wt-%]
• Mn	icp-ms	69.21	133.20	174.54	210.69	387.79	177.79	58.39	i	[ppm]
as MnO		0.01	0.02	0.02	0.03	0.05	0.02	0.01	-	[wt-%]
• Na ₂ O	xrf	0.50	0.69	0.75	0.81	1.02	0.75	0.10	-	[wt-%]
• P ₂ O ₅	xrf	0.01	0.07	0.09	0.11	0.32	0.09	0.04	-	[wt-%]
• S	xrf	0	0.05	0.06	0.08	0.17	0.06	0.03	-	[wt-%]
• SiO ₂	xrf	54.93	68.96	73.26	77.96	89.31	73.56	6.59	-	[wt-%]

Continued on next page

Table 4.13 – continued from previous page

	method	min	25-prec.	median	75-prec.	max	mean	sd	exclude	unit
•	TiO ₂	xrf	0.14	0.34	0.43	0.53	0.43	0.13	-	[wt-%]
	Ti	icp-ms	297.47	509.76	616.04	726.53	607.97	138.84	i	[ppm]
	as TiO ₂		0.05	0.08	0.10	0.12	0.10	0.02	-	[wt-%]
	Ag	icp-ms	0.02	0.04	0.05	0.06	0.05	0.02	i	[ppm]
•	As	xrf	3.66	8.25	10.94	13.52	11.35	4.17	-	[ppm]
	As	icp-ms	1.86	6.82	9.01	11.53	9.57	3.80	o	[ppm]
	Au	icp-ms	0	0.01	0.03	0.06	0.04	0.04	p	[ppm]
	B	icp-ms	7.39	27.69	37	52.62	40.24	16.45	i	[ppm]
	Ba	xrf	194.46	248.20	264.25	288.69	351.30	268	o	[ppm]
•	Ba	icp-ms	21.86	43.27	59.32	76.17	62.01	23.73	-	[ppm]
	Be	icp-ms	0.24	0.61	0.84	1.18	0.91	0.39	i	[ppm]
	Bi	icp-ms	0.02	0.08	0.13	0.19	0.14	0.07	i	[ppm]
•	Cd	icp-ms	0.04	0.09	0.12	0.15	0.12	0.04	-	[ppm]
•	Ce	icp-ms	18.56	30.16	36.58	44.06	37.24	9.29	-	[ppm]
•	Cr	xrf	32.60	60.64	70.62	81.72	71.82	17.65	-	[ppm]
	Cr	icp-ms	10.70	24.17	31.62	41.31	33.68	12.66	o	[ppm]
•	Cs	icp-ms	0.50	2.03	2.95	4.20	3.18	1.50	-	[ppm]
•	Cu	xrf	0.32	4.51	6.21	8.36	6.41	2.87	-	[ppm]
	Cu	icp-ms	1.25	4.25	5.68	7.84	6.15	2.69	o	[ppm]
	Dy	icp-ms	0.94	1.65	2.11	2.59	2.14	0.64	i	[ppm]
	Er	icp-ms	0.45	0.77	0.98	1.21	1	0.30	i	[ppm]
	Eu	icp-ms	0.25	0.50	0.62	0.76	0.64	0.19	i	[ppm]
•	Ga	xrf	3.61	6.58	8.22	10.46	8.66	2.79	-	[ppm]
	Ga	icp-ms	1.15	3.10	4.21	5.80	4.53	1.97	o	[ppm]

Continued on next page

Table 4.13 – continued from previous page

	method	min	25-prec.	median	75-prec.	max	mean	sd	exclude	unit
	Gd	1.32	2.40	3.02	3.59	4.73	2.99	0.83	i	[ppm]
	Hf	0.07	0.17	0.24	0.37	0.74	0.28	0.15	i	[ppm]
	Ho	0.17	0.29	0.37	0.46	0.61	0.38	0.11	i	[ppm]
	La	10.50	17.82	22.04	26.25	37.42	21.98	5.99	i	[ppm]
•	La	8.52	14.68	18.16	21.78	28.05	18.39	4.69	-	[ppm]
	Li	4.17	13.35	19.44	29.23	46.61	21.37	10.22	i	[ppm]
	Lu	0.06	0.11	0.13	0.16	0.22	0.13	0.04	i	[ppm]
	Mo	0.07	0.29	0.43	0.69	1.34	0.50	0.29	l	[ppm]
•	Nb	3.50	7.78	9.31	11.34	15.82	9.51	2.53	-	[ppm]
	Nb	0.17	0.39	0.58	0.90	2.51	0.69	0.39	i	[ppm]
	Nd	12.40	20.08	23.30	26.78	32.60	23.36	4.58	i	[ppm]
	Nd	8.19	14.69	17.91	21.78	27.16	17.98	4.73	i	[ppm]
•	Ni	2.51	10.63	15.57	21.84	35.33	16.38	7.56	-	[ppm]
	Ni	4.01	10.06	13.77	18.52	30.54	14.70	5.82	o	[ppm]
•	Pb	6.53	12.88	15.95	20.26	34.83	16.70	5.16	-	[ppm]
	Pb	2.69	9.23	14	17.57	33.32	13.60	5.29	o	[ppm]
	Pr	2.12	3.58	4.44	5.54	6.78	4.52	1.20	i	[ppm]
	Pt	0	0.01	0.01	0.02	0.02	0.01	0.01	p	[ppm]
•	Rb	34.24	54.02	64.69	81.79	126.77	69.41	20.12	-	[ppm]
	Rb	9.93	25.28	33.92	48.36	99.50	37.95	17.34	o	[ppm]
•	Sb	0.07	0.16	0.19	0.23	0.53	0.20	0.06	-	[ppm]
	Sc	2.90	6.23	7.70	9.38	13.80	7.79	2.38	i	[ppm]
	Sc	1.93	4.26	5.49	7.49	10.51	5.88	2.15	i	[ppm]
	Se	0	0.19	0.56	0.96	1.47	0.60	0.41	p	[ppm]

Continued on next page

Table 4.13 – continued from previous page

	method	min	25-prec.	median	75-prec.	max	mean	sd	exclude	unit
Sm	icp-ms	1.53	2.71	3.30	4.09	5.23	3.37	0.91	i	[ppm]
Sn	xrf	0	1.80	3.21	4.48	8.21	3.24	1.94	p	[ppm]
• Sn	icp-ms	0.18	0.75	1	1.28	2.48	1.02	0.39	-	[ppm]
• Sr	xrf	66.44	96.37	116.58	141.10	221.08	122.18	32.47	-	[ppm]
Sr	icp-ms	26.80	70.91	89.55	116.01	213.02	96.69	36.04	o	[ppm]
Ta	icp-ms	0	0.02	0.03	0.06	0.11	0.04	0.03	i	[ppm]
Tb	icp-ms	0.18	0.32	0.41	0.50	0.68	0.41	0.12	i	[ppm]
Th	xrf	0.84	6.28	7.94	8.85	12.15	7.52	2.29	p	[ppm]
Th	icp-ms	2.17	3.86	4.94	6.32	8.37	5.08	1.49	l	[ppm]
Tl	icp-ms	0.07	0.16	0.25	0.37	0.75	0.27	0.14	i	[ppm]
Tm	icp-ms	0.06	0.10	0.13	0.16	0.23	0.14	0.04	i	[ppm]
U	xrf	0.02	1.88	2.13	2.47	3.20	2.06	0.57	p	[ppm]
U	icp-ms	0.39	0.67	0.81	0.97	1.61	0.83	0.23	i	[ppm]
• V	xrf	16.10	40.11	54.70	75.77	136.29	59.74	27.07	-	[ppm]
• Y	xrf	6.67	15.88	18.69	22.35	31.57	19.08	4.88	-	[ppm]
Y	icp-ms	4.32	7.48	9.20	11.40	16.09	9.56	2.74	o	[ppm]
Yb	icp-ms	0.40	0.66	0.85	1.04	1.39	0.86	0.25	i	[ppm]
• Zn	xrf	6.30	23.54	33.60	47.56	74.15	35.35	15.59	-	[ppm]
Zn	icp-ms	8.92	31.34	40	53.30	89	41.54	15.48	o	[ppm]
• Zr	xrf	122.86	298.91	330.97	365.79	499.01	334.36	51.59	-	[ppm]
Zr	icp-ms	4.13	7.45	9.89	13.59	24.81	10.98	4.47	o	[ppm]

5 Enrichment and natural variability versus anthropogenic impact

5.1 Introduction

For the Netherlands it is often claimed that natural soils, i.e. soils without anthropogenic influence, do not exist anymore (e.g. van der Meent et al. (1990)) . It is believed that over time almost every part of the country has been subjected to the activities of people (see also chapter 2). Despite this extensive influence, Dutch soils are statutorily divided into clean (i.e. non-polluted) and polluted. The National Environmental Policy Plan (VROM, 1997) states that the inventory of polluted and non-polluted soils must be completed by 2005. The background values in Dutch soils will then be re-evaluated based on the set of non-polluted soils. However, this approach has some serious drawbacks. With rigid legal thresholds defining the difference between non-polluted and polluted soil samples, the line between the two categories of samples is extremely narrow. Moreover, the causal link may be missing completely. Furthermore, in view of the widespread anthropogenic influence, the set of non-polluted soil samples will undoubtedly be affected to some extent.

For the Netherlands, soil pollution is defined in environmental legislation. The basis of these normative values is taken from a study of Edelman (1984) that established background values based on soil samples from relatively clean areas. This study was later criticised by van der Meent et al. (1990) because of the poor definition of “relatively clean areas”. In the Netherlands one can, or even must, speak of soil pollution when a measured value exceeds the legal threshold. Even a minor difference in measured concentration can flip the result from non polluted to polluted. This comparison of rigid values does not incorporate any information about variability and uncertainty in both measured and normative value, reducing the outcome of the test to a boolean result. The latter has indeed been evidenced by some soil sanitation projects in residential areas, where individual front and back yards were assessed for the need of sanitation. One garden might be considered clean while the neighbouring garden could be polluted, resulting in a highly patchy pattern of polluted and non-polluted yards (pers. comm.). Although the rigid approach is apparently effective for addressing legal issues, it does not necessarily reflect reality, which is that most soils are altered to some degree by human influence.

To solve the problem, altered elevated concentrations levels should be recognised regard-

less of whether the high values are caused by pollution or natural variation. One way to do this is to compare the data with an unaltered, pristine, reference. In the field of geochemistry this often comprises a geochemical baseline that defines the natural background. (Salminen & Tarvainen, 1997; Matschullat et al., 2000). Salminen & Gregorauskiene (2000) provide an overview of different definitions for such a baseline. Their main points are that a geochemical baseline should describe the natural variation of an element in a particular natural material and should not be considered as a single value but as a fluctuating reference surface over a wider area. They also noted that finer material, such as clay, has a different range of element abundances than other material. Hence a geochemical baseline should incorporate the differences in parent material and underlying geogenic (and biogeochemical) processes.

If deviations from a geochemical background can be coherently discerned, the question still to be answered is if they should be called “pollution”. Pollution, in a more zealous context, is the destroying of purity or sanctity, while another meaning is simple “making dirty”. But when is the purity of the soil destroyed or has someone made it dirty? Is disturbing the soil, but leaving the bulk chemistry intact, also pollution? And what about the use of fertiliser and other enhancements? This “disturbs” the chemical composition of the soil, i.e. making it dirty, but is usually considered as an improvement instead of a destruction. Considering the multitude of human induced alterations to the soil that may lead to increased concentration levels, and considering the problems of defining pollution or its synonym contamination, it is probably better to speak of enrichment. This term, which in fact describes what actually happened, will therefore be used in this thesis.

For the province of Zeeland (south-west Netherlands), the question of how to distinguish between natural and human induced geochemical patterns, was the reason to initiate the present geochemical soil survey. This study aims at establishing the anthropogenic imprint on the soil using a geochemical baseline that comprises the natural variability in soil composition. Results should give a more precise insight in the state of enrichment (pollution), of the soils, and form a firm base for the soil and environmental protection policies of the province.

The use of a natural baseline for background values may seem much more obvious than the use of just a legal threshold but the question remains how to derive such a baseline from present-day, thus influenced, soil data. While chapter 4 described the sampling and analytical requirements, here focus is on data analysis and interpretation. Several (statistical) techniques that vary in complexity and accuracy are available to fulfill the aim of this study. As, among others, Salminen & Gregorauskiene (2000) have shown no best method exists for establishing a baseline, no commonly agreed method for calculating human influence exists either. From the point of view of the end user, the local government, a simple method without the requirement for many (extra) data is preferred. From a scientific point of view, explained variability is of more importance than simplicity. The method must supply a realistic, accurate background value, and a more complex method, using relevant additional data, might be required. This chapter illustrates several techniques that permit the determination of which components are enriched and how enrichments may be quantified. The differences, degree of complexity, and efficacy of these methods will be discussed.

5.2 The Zeeland geochemical soil data

The geological/pedological background of the Zeeland area was given in chapter 2. The geochemical soil survey focuses on the clayey soils developed in the Holocene, marine sediments in the rural, agricultural, part of Zeeland. Figure 5.1 shows the 270 sampling locations. From each location 15 to 20 subsamples were collected from the upper soil layer (plough furrow, ca. 0-25 cm depth) and combined into one composite sample. This sample was subsequently analysed for major, minor and trace elements using X-ray fluorescence (XRF) and inductively coupled plasma-mass spectrometry (ICP-MS) following an aqua regia digestion. The selection of parameters chosen for further data analysis, based on accuracy and precision (see chapter 4) is shown in table 5.1. On average precision and accuracy for the ICP-MS data are slightly less than for the XRF (see chapter 4). Since SiO_2 is, due to closure, inversely proportional to the major components, and also geochemically the least interesting, it will not be treated in this chapter.

In addition to the topsoil samples also subsoil samples (ca. 35-80 cm depth) were collected to serve as a reference, based on the hypothesis that these subsoils are not, or are less, influenced by human processes. For details about the sampling, analytical procedure, accuracy, precision and outlier treatment, see chapter 4.

Table 5.1. Elements selected from the Zeeland dataset

XRF	Al_2O_3	CaO	Fe_2O_3	K_2O	MgO	MnO	Na_2O
	P_2O_5	S	(SiO_2)	TiO_2			
	As	Cr	Cu	Ga	Nb	Ni	Pb
	Rb	Sr	V	Y	Zn	Zr	
ICP-MS	Ba	Cd	Ce	Cs	La	Sn	Sb

5.3 Concepts and methods

5.3.1 Assessing enrichments and baselines in attribute space

There are several statistical methods for assessing patterns of enrichments in geochemical data. Table 5.2 schematically shows the essential differences between these methods. They can be divided into univariate, bivariate and multivariate techniques applied to the studied data set, with or without an external or reference data set. Using this division a methodical approach towards assessing the Zeeland data is presented.

A further division is made between values and (legal) limits. Legal limits can also be regarded as external data to which the soil values can be compared, but these are usually to a certain extent artificial. For instance these limits can be based on a combination of




Figure 5.1. Map of Zeeland with sample locations, see also figure 2.1, p. 22.

negligible risk levels for all life forms and common soil background concentrations. When interest is in the soil geochemical patterns it is obvious that legal limits are not the first choice to serve as a reference. However, they are the basis of the Dutch normative system and soil data obtained from various studies are generally compared with them. Therefore these legal limits are also considered in this study. The univariate and bivariate techniques mentioned in Table 5.2 are detailed below while the multivariate approaches are treated in the next chapter.

Table 5.2. Statistical concepts for assessing enrichments and baselines, for further explanation see text.

	Within dataset	Between datasets		
		reference/external data	legal limits	
Univariate	Frequency plots	Ratios, Spider diagrams	Threshold values, Standard soils	1
Bivariate	Covariance, Correlation	Linear regression of baseline	Normalisation to standard soil	2
Multivariate	Principal components analyses, Clustering	–	–	3

Values	Limits
a	b
	c

Univariate distribution based thresholds (1a)

The advantage of calculating a background based on univariate statistics is that only the measured values for the single element in question are needed. Ideally based on the statistical distribution a threshold can be derived below which the measured values are considered “background” while above the values are considered as “enriched”. In geochemical exploration a threshold value is often used to divide the distribution into background data and enriched (or anomalous) data, the latter hopefully indicating an economic worthwhile mineralization (Rose et al., 1979). These exploration methods can also be applied in environmental soil inventories as explained by Hale & Kühnel (1997) and several methods to calculate background thresholds in environmental studies has been summarised by Matschullat et al. (2000). An often overlooked disadvantage of such a background threshold is that it is based on the overall data distribution which might be shifted towards higher values due to enrichments. In the case of diffuse pollution, enrichments below the threshold are not recognised. However, background thresholds generally do give an indication if a value is within the natural range or not.

One of the proposed methods of Matschullat et al. (2000) is based on the cumulative frequency plotted against the measured value. By using a probability scale for the cumulative frequency axis this method is equal to probability graphs which are known from the geochemical exploration field and are extensively described in Sinclair (1976) (see figure 5.2 as an example). The advantage of such a plot is that a normal distribution plots as a straight line, thus deviances from normality are easily recognised. A clear bend in the curve may indicate a skewed distribution. The presence of higher and more deviating values, is often typical for human influenced components. From the inflection point a background threshold, as concentration or percentile, can be graphically determined. More bends in a curve

may indicate multi-modality. Determining the bends in the plots graphically, results in some subjectivity towards the chosen percentile. In contrast, an arbitrary choice, although objective, might not reflect the optimal threshold. For example, in the Netherlands the 90-percentile, or P_{90} , is an arbitrary level which is often used (van der Gaast et al., 1998).

Comparison with standard data or other reference data (1b)

An often used method to judge if measured values are anomalous or enriched is to compare them with global average values like that for the average upper continental crust (UCC) (Wedepohl, 1995). Despite the differences between average crust and soil material, the UCC is a good reference for element composition, especially for immature soils such as found in Zeeland. A ratio of about one between measured and global UCC value indicates no deviation from the global average while ratios much larger or smaller than one indicate an anomaly. Anomalies with ratios larger than one can be considered as enriched towards the global average, irrespective of the fact whether this enrichment is caused by natural processes (like the enrichment of more mobile elements in weathering products) or caused by enrichment due to human input.

Another approach is to use reference values from pristine soils of the same area. The great advantage is that the reference soils have then been subjected to the same natural processes as the soils of interest, yet not subjected to human influence (Facchinelli et al., 2001; Shotyk et al., 2001). In this case a ratio different than one might indicate anthropogenic influence. The data collected for this research (see chapter 4) therefore incorporate subsoil samples, under the assumption that these are chemically less or not disturbed by man. The means or medians of this subsoil data set can then be used as a local reference value comprising the geochemical background.

Using global averages or local references has as drawback that the actual field values are compared with a single value only, neglecting natural and sampling variance. A single ratio has limited value without any information about variability. Therefore also the ratios of the 2nd and 4th quartile of the data will be calculated to provide insight in the variability.

Legal thresholds (1c)

The same method, as with the “real” global averages or local backgrounds, of calculating ratios can be applied to legal limits like the Dutch legal threshold values (Dutch: “Streefwaarden” or *S*) (VROM, 1994; Swartjes, 1999). For the Dutch situation disadvantages of this technique are that legal threshold limits are defined, besides organic compounds, for only 16 elements (As, Ba, Be, Cd, Cr, Co, Cu, Hg, Pb, Mo, Ni, Sb, Se, Ta, V, Zn), and that their derivation is not well documented in international literature. For Sn, a legal limit is lacking, only an indicative value can be derived from regression parameters (see below). Besides the value for *S*, an intervention value is defined that signals the level of severe pollution (Dutch: “Interventiewaarde” or *I*). This level is based on risk rather than on background soil composition, and it will not be further considered in this study (Swartjes, 1999).

Bivariate distribution and explained variance (2a)

Clays are in essence Al bearing phyllosilicates which, especially for typically clayey soils such as found in Zeeland, have a strong correlation with Al_2O_3 (Tack et al., 1997; Wilcke et al., 1998; Sharma et al., 2000). Clay content is usually also considered to be a proxy for natural concentrations of many other elements and as such is currently used in normalising Dutch legal limits towards the “standard soil”. Many scientist believe that the strong correlation of many elements with clay content is caused by substitution and solid solution of these elements in the phyllosilicate matrix, resulting in an even more direct relation with Al_2O_3 than with clay (van Gaans et al., 1995; Mol, 2002; Tebbens et al., 2000; Bianchini et al., 2002; Huisman et al., 1997), the latter being based on grain size and not on mineralogy. Consequently Al_2O_3 content can be used as a proxy for pristine soil concentrations in relation to parent material.

If elements show a high correlation with Al_2O_3 then the variability of such an element is for a large part explained by the natural variability of Al_2O_3 . If no such strong correlation exists, then it is very likely that other processes determine the element variability, either natural or anthropogenic. For example, elements related to heavy minerals, like Zr, or related to biogenic processes, like CaO, are expected to have no or a low correlation with Al_2O_3 , emphasising that their mineralogy is not akin to the clay phyllosilicates.

Calculating the correlation with Al_2O_3 is a rather simple technique but it requires a relatively large number of data points for confidence. The advantage though is that no external reference samples are needed, so the method can be used irrespective of the presence of a reference data set.

The use of correlations with Al_2O_3 does not directly result in a baseline but shows for which elements the major part of their variability can be explained by natural variation in aluminosilicate content. For these elements it can then be assumed that their concentration is not influenced by external processes and can thus be regarded as representing the geochemical baseline.

Bivariate comparison with reference data (2b)

Instead of looking for a distinction between natural and anthropogenic relations within the data set, it is also possible to use a reference data set for which it is assumed that the compositional relations are primarily the result of natural processes, without significant anthropogenic variability. As elaborated before, here the subsoil samples are considered as the natural reference.

Some researchers prefer to take just the ratio between the top and subsoil sample collected at one location (Facchinelli et al., 2001). This procedure is not suitable for Zeeland since it is possible that the two soil layers have a different sedimentary history, thus form distinct different layers with different composition (see chapter 2). Other researchers try to overcome this problem by first normalising this ratio, r , with a conservative element (Blaser et al., 2000) using

$$r = \frac{\left(\frac{x}{y}\right)_{top}}{\left(\frac{x}{y}\right)_{sub}} \quad (5.1)$$

in which x is the element of interest normalised to the conservative element y . This method always requires 4 measurements for calculating an enrichment. Furthermore the covariance between x, y for both soil depths must be studied closely to prevent spurious results (van der Weijden, 2002).

Another method uses linear relations with clay or Al_2O_3 content, or any other soil or sediment parameter that is known not to be influenced, or conservative (Bauer et al., 1992; Bauer & Bor, 1995). As does the calculation of the direct within data set correlation described above, this method requires a sufficiently large number of reference data points for confidence, and of course a good correlation between the element considered and the predictor variable used. In this thesis the subsoil data set with the outliers omitted is used as undisturbed reference. Sulfur, due to the relatively large number of outliers, was excluded. Linear equations were derived in which the estimated concentration (C_{est}) of each element in the topsoil was expressed as a function of Al_2O_3 content using

$$C_{est} = Al_2O_3 \cdot a + b \quad (5.2)$$

where a and b are the slope and intercept from the regression analysis based on the subsoil data, and Al_2O_3 is the measured concentration in the topsoil sample.

A significant deviation of the actually measured concentration from the calculated value is evidence of an enrichment caused by processes other than natural variability in clay/phylosilicate content, which is calculated as

$$e = \frac{C_{top}}{C_{est}} \quad (5.3)$$

where C_{top} is the analysed topsoil sample concentration of the element considered. Compared to equation 5.1, this method only requires 2 components analysed once the regression parameters are established.

It was decided to use ordinary least squares (OLS) regression. Although some researchers consider “better” regression techniques in geochemistry, e.g. Szava-Kovats (2002), these techniques are usually trying to minimise the influence of outliers and to compensate for the error in the independent variable. Since the outliers had already been omitted from the subsoil data set and considering the analytical error of only 0.8% (see table 4.4, p. 74) for Al_2O_3 I presume that OLS regression is adequate.

Although theoretically the possibility of depletion of anthropogenically influenced components does exist, it is not very likely in young marine clayey areas such as Zeeland. If

depletion occurred, for example due to acidification of the soil, the situation would easily be recognised. Depletion due to weathering is not considered, except for calcite related components. Because of the very young age of the Zeeland soils the differences between top and subsoil caused by weathering are expected to be negligible, and within the bounds of sampling and analytical noise.

From the enrichments e , and after establishing that the enrichments are of anthropogenic origin, it is possible to calculate a contamination degree d (Rapant et al., 1999). This degree indicates the amount of contamination caused by a group of n elements by calculating the average enrichment using

$$d = \frac{1}{n} \sum_{i=1}^n e_i \quad (5.4)$$

Soil type dependent legal threshold (2c)

According to the normative framework, Dutch legal threshold values are dependent on the soil type using clay content (grain size $<2\mu\text{m}$) and organic matter content as the primary distinctive soil properties for all Dutch soil types (VROM, 1994). Threshold values are given as maximum permissible levels for sustainable soil quality (S) and are based on a so called standard soil containing 25% clay and 10% organic matter. Using the dictated equation

$$s = S \cdot \frac{\alpha + C\beta + O\gamma}{\alpha + 25\beta + 10\gamma} \quad (5.5)$$

the given value S can be recalculated to the soil dependent value s using the measured organic matter (% O) and clay (% C) content of that soil (VROM, 1994). The regression parameters α , β , and γ , are also provided by the normative framework. In principle the S values are valid only for concentrations determined in aqua-regia extracts, while here most elements are analysed by XRF, giving total concentrations. For most elements the discrepancy is negligible except for Ba and V. For Ba, therefore, ICP-MS data were selected but for V precision with ICP-MS was poor. Thus V is omitted from the comparison with S .

As noted earlier, only for 16 inorganic elements are legal threshold available. Most of the Dutch soil surveys within the scope of environmental law or practice, are standardised towards them. Data are often expressed as either the ratio of the measured values and the soil dependent threshold limit or as the equivalent concentration in standard soil, as it is believed that such a standardisation facilitates comparison between areas and surveys.

Table 5.3. S values and regression (α , β , γ) parameters (see equation 5.5) for relevant elements. S -value for Sn derived from regression parameters, no official value given. For Sb no regression parameters are given (VROM, 1994).

	α	β	γ	S value
As	15	0.4	0.4	29
Cd	0.4	0.007	0.021	0.8
Cr	50	2	0	100
Cu	15	0.6	0.6	36
Ni	10	1	0	35
Pb	50	1	1	85
Sb	-	-	-	3
Sn	4	0.6	0	19
Zn	50	3	1.5	140

5.3.2 Spatial representation and spatial anomalies

In addition to discerning geochemical baselines and enrichments in attribute space, plotting geochemical soil data in geographical space might reveal regional patterns or anomalies. Regional patterns are usually visible as spatial groups of points with similar values indicating regional features of the soil composition. Anomalies would be single points with a divergent value compared with other surrounding points, denoting either a depletion or, as expected for Zeeland, an enrichment compared to the surroundings. These anomalies might not occur as such in attribute space because the data point is not at the extreme of the distribution but only in comparison with spatially neighbouring samples.

If no mathematical or geostatistical spatial model for each point in geographical space is assumed, only values related to the sampled locations can be plotted. To make the value graphically visible symbols will be used for which the size varies with the magnitude of the attribute concerned. Examples of such symbol maps already exist (Reimann et al., 2002; Bounessah & Atkin, 2003; McGrath & Zhang, 2003). The size function should preferably be continuous, suitable to differentiate between high and low values, and should adjust to the observed frequency distribution of the attribute. Therefore the plotting method of the Geological Survey of Finland was adopted (Gustavson et al., 1997), which is based on bubble plots for which the bubble size is an exponential function which monotonically increases with the cumulative distribution. This bubble size function is designed for geochemical variables which follow a positively skewed or log normal distribution, emphasizing the upper values. This function can be regarded as a variant of the univariate cumulative frequency distribution, where the bubble size shows the position of the data point within the distribution, small sizes corresponding with values in the lower part and large sizes corresponding with values in the upper tail.

5.4 Results and interpretation

5.4.1 Univariate statistics for the Zeeland topsoil data.

Cumulative frequencies

Figure 5.2 shows the cumulative frequency distributions of 12 selected components together with the 90-percentile (P_{90}). Some components like Al_2O_3 and Rb have straight, somewhat S-shaped, curves while other components like Cd, P_2O_5 , and Sb have a clear bend, and a steeper slope, at the top. This suggests that the latter have a deviation towards higher values and might be locally enriched. The components La and Na_2O have clear outliers around the P_{99} showing that this technique can be used also to detect outliers. From the almost straight line of components like Al_2O_3 and Fe_2O_3 it can be concluded that their distribution follows the normal distribution and that it is likely that they are uninfluenced by anthropogenic processes.

The component P_2O_5 shows that the bend in the top of the curve starts around the P_{90} point suggesting that the P_{90} value is a reasonable boundary for distinguishing enriched, influenced soil for this component. For Cd, Cu and Sb the P_{80} or P_{85} seem more appropriate (see table 5.4). For the assumed non influenced elements, e.g. Al_2O_3 , Fe_2O_3 , and Rb, the P_{90} definitely is an underestimation of the maximum background level, which theoretically is the P_{100} , the P_{99} is used instead. This shows that the P_{90} , as used in Dutch practice, is partly useful but only if the corresponding cumulative frequency plot is studied as well. Preferably the percentile should be adjusted in accordance with the plot. Using an arbitrary value, there is a clear risk of over- or underestimating the background threshold.

The P_{90} , P_{95} , and P_{99} values for all the elements are given in table 5.4. The optimal threshold is determined by choosing the optimal percentile based on the bends in the cumulative frequency plots (plots are not shown). Despite the subjectivity of this method, within the 5% resolution adopted, the results obtained are adequate.

The use of the cumulative plots is helpful in indicating which elements show anomalies relative to a normal distribution. While this method is very simple and easy to apply, it gives no indication if a component is anthropogenically enriched or not. Also it does not recognise enrichments which occur within the range of the normal distributed uninfluenced data, e.g. some concentrations on the lower end of the distribution can be substantially enriched without changing the relation in the cumulative frequency plots. If all the samples are enriched, e.g. by diffuse contamination, then this is not revealed in the CDF plots as well.

Spatial representation

The regional distributions of the same elements as selected for figure 5.2, are illustrated in figure 5.3 and 5.4. Figure 5.3 depicts elements that are expected not to be influenced by human processes and figure 5.4 shows the elements that are probably influenced by man. The




Figure 5.2. Cumulative probability plots of a selection of components. Percentiles are plotted along the X axis using a normal distributed probability scale, concentrations are plotted along the Y axis (oxides in [wt-%], others on [ppm]). The dashed lines depict the P_{90} for the X-axis and the corresponding value on the Y-axis. Percentiles of these and other elements are given in table 5.4.

components Al_2O_3 , Fe_2O_3 , La, and Rb all have similar regional patterns while Na_2O and CaO have distinct patterns. The first four components are related to the clay content, while Na_2O and CaO respectively may be indicative for the silt fraction and carbonate sedimentation/leaching respectively.

The elements suspected of anthropogenic influence, all have more or less their own regional pattern, although some are partly comparable to patterns in figure 5.3. For example the regional pattern of As is quite similar to that of CaO while the same is true for Zn and Al_2O_3 . This is a first indication that these elements are largely related to natural processes. Elements with a more distinct pattern, like Cu, P_2O_5 , Cd, and Sb probably have an additional unique underlying process causing the regional variation, which is presumably of human origin. Also the regional scale of the variability differs. Components like P_2O_5 and Sb show small areas, i.e. a few locations together, with elevated concentrations, while Cu,

Table 5.4. Percentiles of the data of Zeeland topsoil. Optimal threshold percentiles, determined graphically, are noted in bold face. If no optimal threshold is given, then it is below the P90 and shown below the table

Element	P90	P95	P99	Element	P90	P95	P99
Al ₂ O ₃	10.94	11.56	12.05	Cs	5.46	6.23	6.90
CaO	5.99	6.91	7.93	Cu	21.44	24.26	29.14
Fe ₂ O ₃	5.32	5.60	6.15	Ga	12.53	13.36	14.54
K ₂ O	2.14	2.21	2.31	La	25.36	26.52	28.55
MgO	1.27	1.37	1.54	Nb	12.71	13.06	13.64
MnO	0.06	0.06	0.08	Ni	27.08	30.35	33.94
Na ₂ O	0.84	0.86	0.90	Pb	37	42.48	55
P ₂ O ₅	0.22	0.26	0.34	Rb	96.74	105.17	111.25
S	0.14	0.17	0.26	Sb	0.42	0.51	0.71
TiO ₂	0.60	0.62	0.67	Sn	2.38	2.59	3.95
As	20.67	21.64	24.81	Sr	146.02	158.66	182.99
Ba	100.87	108.51	116.12	V	95.62	102.01	114.67
Cd	0.37	0.40	0.59	Y	24.40	25.30	26.54
Ce	50.83	52.19	55.91	Zn	74.51	80.07	98.84
Cr	94.82	99.28	102.60	Zr	394.05	422.05	466.05

P80: Cu(18.93 ppm). P85: Cd(0.35 ppm), Sb(0.36 ppm)

As, and Zn show more regional features.

5.4.2 Univariate comparison with reference data and legal threshold levels

The upper continental crust

Figure 5.5 depicts the ratio of the median values of the concentrations as measured for the Zeeland soils and those given for the upper continental crust (UCC) (Wedepohl, 1995).

In this comparison, for Ba the XRF values are used instead of the much lower ICP-MS data, as the UCC is also based on total concentrations. Compared to the UCC and the overall pattern the elements As, Cd, Cr, Pb, and Zr are relatively enriched while especially Na, and to a lesser extent Ca, K, Mg, Mn, and Sr are depleted. Most other elements lie between a ratio of about 0.5 and 1.0. The Zeeland soils on average appear somewhat less aluminous and ferromagnesian than the UCC.

The elements Pb and Cd are known as contaminants in fertiliser (de López Camelo et al., 1997; Gimeno-García et al., 1996; Nash et al., 2003) and to be constituents of atmospheric




Figure 5.3. Regional distribution of a selection of components expected not to be influenced by human processes

deposition of traffic exhausts (Parkpian et al., 2003). It is expected that these processes, fertilization and atmospheric deposition, are at least partially responsible for the enrichment compared to the upper continental crust. Natural processes, like mineralisations of these elements, are not very likely for Zeeland. Ratios with the UCC from sedimentary studies




Figure 5.4. Regional distribution of a selection of elements expected to be influenced by human processes

in alluvial and fluvial sediments also indicate that concentrations of elements like As, Cd, Cr, and Pb are relatively high and can be related to anthropogenic processes (Ridgway et al., 2003; Singh et al., 2001). The high ratio of As might be caused by natural processes related to upward groundwater seepage in the lowlands of the delta of Zeeland (Nierop,




Figure 5.5. Ratio of median values for Zeeland topsoil and the average upper continental crust. The bars show the interval of the ratio between the 25% and 75% quantile of the data. For Ba, the XRF analyses were chosen.

2000). However, As occurs in certain pesticides as well (Folkes et al., 2001), but I expect its influence to be unlikely, as will be shown below. The low Na concentration might indicate a high degree of pre-depositional weathering of the sediment.

In figure 5.6 the ratio between the medians of the top and subsoil samples is shown. The ratio for Al (1.13) is depicted by a vertical line and shows that the topsoil has a higher Al content. This suggests that the topsoil consists of heavier clays, which can be attributed to a more quiet marine environment and the sedimentation of the finest fraction during the reclamation of the Zeeland polders. This confirms that a direct, local, comparison between topsoil and subsoil concentration for Zeeland would indeed be unsuitable, and that at least a normalisation according to equation 5.1 would be required.

The elements As, Cd, Cu, P, Pb, S, Sb, Sn, and Zn are clearly enriched compared to the subsoil and, according to Senesi et al. (1999), can be related to anthropogenic activities. CaO and Sr are depleted from the topsoil. Assuming that the CaO content is mostly determined by the calcite content of the soils it can be inferred that this depletion is caused by weathering. In general the element Sr is highly correlated with Ca in marine environments and often a replacement for Ca in calcite, hence Sr is depleted together with CaO.

Most of the other elements are close to the line round the 1.13 ratio. This suggest that these elements are closely related to the Al and clay content, and are not anthropogenically enriched in the Zeeland topsoil.

Comparing the local subsoil reference with the global UCC reference shows that the Cr and Zr enrichments found for Zeeland compared to the UCC are probably a natural phenomenon, related to a typical heavy mineral association. The high topsoil/subsoil ratio for Cu is not supported by the ratio of about 1.0 with the UCC, for this element the UCC is apparently an over-estimation of the Zeeland natural background. Given the fact that the As content of the subsoil is relatively close to that of the topsoil, in relation to the enrichment of the Zeeland soils in comparison to the UCC, it appears that at least part of the As enrichment is due to natural processes.

While the global UCC reference highlights the regional features and facilitates comparison with other areas, the local reference indeed provides much more insight into the (anthropogenic) nature of the enrichments found.

Legal thresholds

To compare the data with the legal threshold values, the ratio between the median topsoil values and the standard threshold value were calculated and depicted in figure 5.7. A comparison with soil specific threshold values is presented as well, based on the standard soil containing 25% clay and 10% organic matter using equation 5.5. Since no clay and organic matter values were available in the geochemical dataset, the clay content was estimated using the relation between Al_2O_3 and clay content obtained from another study in Zeeland (data not shown)* and the median clay content was used for the calculation. Organic matter was set at the median organic matter value from the same data set[†]. The smaller bars, as compared to figure 5.5 and 5.6, are caused by the higher values of the legal threshold values, resulting in much smaller ratios. Exceedance of the soil specific threshold values is

*The relation used is $4.76 \cdot [Al_2O_3] - 15.47$ and derived from the data set of chapter 7, $n=35$

[†]Median value organic matter is 3%




Figure 5.6. Ratio of median values for the Zeeland topsoil and the median of the subsoil. The bars show the interval of the ratio between the 25% and 75% quantile of the data while the vertical line shows the ratio of Al. For Ba, both the XRF and ICP-MS analysis were chosen.

sparse. Only Cu and Zn have the maximum number of 3 samples larger than the threshold. Most of the values show a ratio around 0.5, except Cr which has a ratio of about 0.8. This is probably caused by using the XRF for analysing Cr while the threshold values are based

on aqua regia leached digested samples (see chapter 4). This may also be the cause of the slightly higher ratio for Ni. Both Ni and Cr show a ratio between top and subsoil comparable to that of Al_2O_3 , suggesting that there is no anthropogenic enrichment. While Zeeland is less aluminous than the UCC, Ni and Cr are comparatively much higher, which for Cr was already ascribed in chapter 4 to its incorporation in heavy minerals. For Sb and Sn the reverse is found, a slightly higher ratio for the comparison between top and subsoil, while having a lower ratio compared to the legal threshold. Both Sb and Sn were analysed by ICP-MS using an aqua regia extraction, and were not found to be regionally low in Zeeland compared to the UCC. This suggests that, in comparison to the other priority elements, the legal thresholds for Sb and Sn are relatively high compared to their natural background in Zeeland.

For all elements it is clear that the legal threshold values are substantially higher than the median values found in the topsoils. This shows that the legal threshold value is not suitable to reveal the moderate, yet evident, enrichment in the topsoil of Zeeland.

5.4.3 Correlation and regression with Al

Explained variance versus natural variability

The Pearson correlation of the elements with Al_2O_3 is given in table 5.5. Most of the elements have a positive correlation with Al_2O_3 which more or less confirms the closely related ratios in figure 5.6; only Na_2O and Zr have a negative correlation. The correlations of CaO and Sr are not significantly different from zero. However, the correlation between CaO and Sr is 0.953 supporting that they both are the result of (biogenic) calcite precipitation and dissolution (pre- and post-depositional). The components P_2O_5 , Cd, Cu, Pb, S, Sb, and Sn are slightly less correlated with Al_2O_3 . These are also elements that are enriched in the topsoil according to figure 5.6. Both enrichment and lesser correlation with soil mineralogy proxies are suggesting that the soil concentrations of these elements are indeed altered by human processes.

Relation with Al content

To study the variability and concentrations due to natural mineralogical variability the measured values were plotted against the concentration of Al_2O_3 . Figure 5.8 shows scatter plots of selected elements versus Al_2O_3 , that illustrate the different types of relations found. Both topsoil and subsoil relations are depicted for clarity.

Cadmium, as best example, shows two distinct groups: a group of topsoil samples and a group of subsoil samples. The latter has lower Cd values and less residual variance than the first. If higher Cd concentration are caused by anthropogenic input, it can be assumed that this occurs more erratically than natural processes and thus causes higher variability in the influenced soil horizons. In chapter 3 it was already shown that presumed contaminants have a higher spatial variability than non contaminants. The other elements that also show




Figure 5.7. Ratio of median values and the legal threshold value. Circles are the comparison with the soil specific threshold value, based on the median clay content, and the bars show the range between the 25% and 75% quantile. Triangles are the comparison with the non standard values (see text for explanation). For Ba the ICP-MS analysis was chosen.

two different groups are As, Cu, Pb, Sn, Sb, and Zn. However, the separation between the top and subsoil is less evident than for Cd. Like in the case of Cd, the topsoil variability is always higher than that in the subsoil, presumably due to the same causes. However, for As and Zn the differences between top and subsoil variance seems smaller. For Ni and Cr all the points are close to one line and no grouping between top and subsoil can be seen. This also shows how Ni and Cr are highly correlated with Al_2O_3 . CaO and P_2O_5 are examples of elements that show a clear difference between top and subsoil, but are not very well correlated with Al_2O_3 . The topsoil is somewhat lower in CaO and higher in P_2O_5 than the subsoil and for CaO also within the top and subsoil different groups might be discerned. A poor relation with Al_2O_3 is also apparent for Zr. However, Zr shows no clear distinction between top- and subsoil.

Table 5.5. Pearson correlation with topsoil Al₂O₃. All correlations, except for CaO and Sr, differ significantly from zero.

Element	r	Element	r	Element	r
Al ₂ O ₃	1.000	Ba	0.882	Pb	0.571
CaO	-0.041	Cd	0.349	Rb	0.987
Fe ₂ O ₃	0.970	Ce	0.883	S	0.478
K ₂ O	0.970	Cr	0.954	Sb	0.396
MgO	0.946	Cs	0.928	Sn	0.401
MnO	0.675	Cu	0.461	Sr	-0.079
Na ₂ O	-0.280	Ga	0.987	V	0.973
P ₂ O ₅	0.351	La	0.880	Y	0.890
TiO ₂	0.975	Nb	0.960	Zn	0.843
As	0.735	Ni	0.970	Zr	-0.336

Subsoil baselines and topsoil enrichment

The natural, or background, concentration of Al related components can be estimated from the Al₂O₃ concentration. Although for some elements this relation is disturbed in the topsoil, for the subsoil it still exists, as is shown in figure 5.8, and can thus be used to calculate the enrichment of the topsoil. Based on the results above, baseline enrichments were calculated for those elements that are enriched in the topsoil and suspected to be anthropogenically influenced and that show good correlation with Al₂O₃ in the subsoil, i.e. As, Cd, Cu, Pb, Sb, Sn, and Zn.

Table 5.6 lists the slope and intercept of the regression model of each element with Al₂O₃ as the predictor, together with the correlation R^2 . From the subsoil regression the standard error (ϵ) of the prediction was calculated as well. Using the same method as for calculating e (equation 5.3), e_u and e_l were calculated for the median Al₂O₃ concentrations as

$$e_u = \frac{C_{est} + \epsilon}{C_{est}} \quad e_l = \frac{C_{est} - \epsilon}{C_{est}} \quad (5.6)$$

The range of e_l to e_u for a component is called the ‘ e -range’ and determines the range of enrichment values that might be expected solely by natural variation (5.6). Regression lines, confidence intervals, and standard error are depicted in figure 5.8 as well.

Table 5.7 shows the percentiles of the enrichment ratios e , while table 5.8 similarly shows the enrichments as the difference between the topsoil concentration (C_{top}) and the estimated concentration (C_{est}) (see equations 5.2 and 5.3).

To compare the elements among each other, the enrichment ratios (e) are graphically dis-




Figure 5.8. Scatter plots of Al_2O_3 versus selected elements. Black dots are subsoil samples, crosses are topsoil samples. Al_2O_3 is on the x-axis, elements are on the y-axis. Regression lines are drawn for the subsoil data with confidence intervals (inner dashed lines) and standard error (outer dashed lines). *continued at next page*

played in figure 5.9. These box plots show clearly that Cd and Cu are more enriched than the other elements. The range of e for the natural subsoil variation (table 5.6) is for these elements below the P_{25} showing that their enrichment is substantially higher than can be expected from natural (or random) variability. This again suggests that artificial fertilizer usage is a major process that influenced the Zeeland soils. The elements Pb, Sb, and Sn are less enriched, the upper limit of enrichment caused by natural variability lies between the P_{25} and the median topsoil enrichment. Zn and As have the smallest enrichment ratios and




Figure 5.8. *continued from previous page* Scatter plots of Al_2O_3 versus selected elements. *continued at next page*

their medians are within the range of the natural variability. Overall, the decreasing order of enrichment, based on the full range of table 5.7, is $Cd > Cu \gg Sn \approx Sb > Pb > Zn > As$. The differences between the topsoil concentration and estimated concentrations were compared with the enrichment found in a comparable agricultural and young marine clay area in the “Nord-Pas de Calais” (north-west France) (Sterckeman et al., 2004). The results are strikingly similar for the elements Cd, Pb, Sb, Sn, and Zn while Cu enrichment is almost twice as high in our research. This suggests that the enrichments in Zeeland are in general not specific for the area but are due to processes that are active within a large region or might be universal for similar areas. The reason for the higher concentration of Cu for




Figure 5.8. *continued from previous page* Scatter plots of Al_2O_3 versus selected elements.

Zeeland is unknown.

5.4.4 Spatial patterns of topsoil enrichments

Regional variability of enrichment ratios

The enrichment e for the seven selected components is geographically depicted in figure 5.10. Each component has more or less its own geographical pattern and only those of Sn and Cu show some similarities. The regional variability is large, and while it is possible

Table 5.6. Regression constants and R^2 of the regression of subsoil concentrations of selected elements as a function of Al_2O_3 . a is the slope and b is the intercept of the regression. The e range gives the range of enrichment based on the upper and lower limits of the regression standard error of prediction.

Element	a	b	R^2	e range
As	1.44	0.56	0.77	0.51-1.49
Cd	0.0074	0.072	0.29	0.57-1.43
Cu	0.80	0.48	0.64	0.62-1.38
Pb	1.58	4.76	0.65	0.63-1.37
Sb	0.011	0.12	0.29	0.57-1.43
Sn	0.086	0.40	0.42	0.49-1.51
Zn	5.73	-7.84	0.90	0.67-1.33

Table 5.7. Enrichment ratios in the topsoil, The minimum (min), 25-percentile (25 perc.), median, 75 percentile (75 perc.), and the maximum (max) ratio are given

element	min	25 perc.	median	75 perc.	max
As	0.63	1.07	1.22	1.46	2.23
Cd	1.10	1.93	2.14	2.39	6.66
Cu	1.01	1.53	1.86	2.46	5.9
Pb	1	1.27	1.41	1.71	4.19
Sb	0.85	1.33	1.50	1.72	5.21
Sn	0.56	1.25	1.56	1.88	5.16
Zn	1.07	1.25	1.33	1.45	4.62

to distinguish subregions they are less well expressed as the spatial patterns of the non influenced elements (compare figure 5.3). Considering the enrichment of these elements as anthropogenic this lesser regional correlation can be explained by the probably rather erratic human processes.

The spatial pattern of the overall contamination index d is shown in figure 5.11. Some locations show high index values but in general they lie around a ratio of 2. From a regional perspective the areas in the north-east of Zeeland (Schouwen-Duiveland, St. Philipsland, Tholen) seem to be more enriched, followed by Noord-Beveland and Zuid-Beveland. Besides regions with high values some local anomalies also can be seen, which comprise several enriched elements (compare figure 5.8).

Table 5.8. Enrichment differences for the topsoil [ppm]. The minimum (min), 25-percentile (25 perc.), median, 75 percentile (75 perc.), and the maximum (max) difference are given

element	min	25 perc.	median	75 perc.	max
As	-4.85	0.87	3.03	5.31	13.29
Cd	0.02	0.12	0.15	0.19	0.82
Cu	0.08	3.62	6.15	9.70	49.83
Pb	0.02	4.57	7.25	12.19	59.67
Sb	-0.03	0.07	0.10	0.15	0.76
Sn	-0.62	0.28	0.65	0.94	5.79
Zn	2.77	9.57	12.60	16.82	175.54




Figure 5.9. Box plots of the enrichment ratio e . Outliers are shown as circles.




Figure 5.10. Spatial pattern of the enrichment for selected elements. Gray dots have an enrichment within the natural variability of the subsoil. See table 5.7. *continued on next page*

Ratios with legal thresholds

It is also possible to calculate the ratios with the soil specific legal threshold level. Again calculation is based on equation 5.5 and table 5.3. For the value of clay C and organic matter O the same values are used as in §5.4.2. The results are shown for completeness in figure 5.12.

Since both the enrichment e and the ratio with the soil specific threshold s are based on the variation in Al_2O_3 (a constant value is taken for O) the resemblance between figures 5.10 and 5.12 is evident. However, the intercept for calculating s is larger (see table 5.3),




Figure 5.10. *continued* Spatial pattern of the enrichment for selected elements. Gray dots have an enrichment within the natural variability of the subsoil. See table 5.7.

and as a consequence the relative range of s -values is generally much smaller, resulting in enrichment maps that are still very similar to the element maps themselves. Exceptions are Cd, for which the relative effects of dividing by C_{est} or s are similar, and Sn, for which the relative range in s -values is much larger than the range in C_{est} , due to the steep slope associated with the clay-content.

A more important difference is that for the enrichment e the extent of enrichment values, that can be expected from natural variability is known through the standard error of the prediction. This variability is a distinct property of the linear regression model underlying the calculation of e while the legal threshold values is just a rigid, though normalised for




Figure 5.11. Spatial representation of the contamination index, i.e. the average enrichment of As, Cd, Cu, Pb, Sb, Sn, Zn.

parent material, value instead of a range. A second advantage of the enrichment model is that it uses a local baseline rather than a general model which tries to incorporate a wide range of parent materials.

5.5 General discussion and conclusions

A structured approach, using several statistical techniques that increase in complexity and the amount of auxiliary data needed, provides detailed insight in the soil geochemistry of




Figure 5.12. Spatial pattern of the ratio with the soil specific s -value for selected elements. *continued on next page*

Zeeland. This approach first of all shows which components are influenced by, most likely, anthropogenic processes. Then, this anthropogenic imprint can be quantified based on their (mutual) relation with non-influenced soil components.

Cumulative frequency plots are a good aid to show at a glance which elements have anomalous concentrations. From these plots percentiles can be derived dividing anomalous data from background. The use of the value corresponding to the P_{90} , an often used standard percentile in Dutch soil surveys, as a threshold for assumed to be influenced elements might be a responsible choice, but only if the corresponding cumulative distribution plots are studied and justify the choice. In this study most (trace) elements are chiefly associated with




Figure 5.12. *continued* Spatial pattern of the ratio with the soil specific c_s -value for selected elements

natural processes and the P_{90} is an obvious underestimation, the P_{99} is more appropriate. Also for Pb the P_{90} is an underestimation, while for some other probably anthropogenically influenced elements (Cd, Cu and Sb) the P_{90} is an overestimation of the threshold value. This suggests that the arbitrary choice for the P_{90} in many Dutch soil surveys should be reconsidered. Notwithstanding the advantage of its simplicity, the method only shows if a components distribution deviates from normality with a distinct tail towards higher values. It does not indicate the cause of the deviation, anthropogenic or natural, nor does it provide a clear overview of the extent of the enrichments, due to the subjectivity and sometimes ambiguity in the choice of the threshold percentile. However, at a reconnaissance stage the method can be very useful.

The spatial distribution of elements with low threshold percentiles (P_{80} to P_{90}) and suspected of human influence, shows regional variability that differs distinctly from the more or less similar patterns that emerge for elements related to natural processes like variability in clay/ Al_2O_3 . This indicates that these probably influenced elements indeed have a more unique process underlying their regional variability.

Calculation of median ratios, with the upper continental crust (UCC), median subsoil concentrations, and legal limits, is an easy method to determine which elements are enriched compared to the reference used. Ratios with the UCC indicate that As, Cd, Cr, Pb, and Zr are relatively enriched. For As it is assumed that this enrichment is partially caused by the low lying deltaic environment of Zeeland, Cr and Zr could be related to heavy mineral concentrations, while the other elements might be enriched due to, for example, fertiliser usage and/or traffic exhaust. Depleted elements are Ba, Ca, K, Mg, Mn, Na, and Sr, which are assumed to be a natural regional feature. On average, the Zeeland soils also appear somewhat less aluminous and ferromagnesian than the UCC. The ratios based on the subsoil show that the elements As, Cd, Cu, P, Pb, S, Sb, Sn, and Zn are enriched. These are all elements for which enrichments can be related to anthropogenic activities. The ratios with the official legal limits for Dutch soil, often used as reference values, show ratios of about 0.5-0.8, with the higher ratios (Cr, Ni) probably caused by the better analytical performance of the XRF method used as compared to the prescribed aqua-regia extraction. Even the P_{99} of the ratios, based on median clay values, are still below one for all elements.

The implications of the ratio results are basically twofold: 1) the area of Zeeland covered in this study is clean as regarding the governmental regulations and 2) the legal threshold limits are an overestimation of soil background concentrations, already including (minor) anthropogenic enrichments. They are thus less suitable as a reference value for a study into natural and anthropogenic geochemical patterns, and the ratios with the local reference (subsoil data) are preferred for identifying the influenced elements and determining the extent of their enrichment.

The correlation with Al_2O_3 is easily calculated and proved to be very useful to indicate elements which are possibly altered by human processes. The elements P_2O_5 , Cd, Cu, Pb, S, Sb, and Sn show relatively low correlations with Al_2O_3 in the topsoil compared to other elements and are also enriched in the topsoil. CaO, MnO, Na_2O , and Sr show relatively low, or negative, correlations with Al_2O_3 and are not enriched in the topsoil. The correlations confirm the similarities observed in the spatial distributions of the elements.

Overall higher correlations with Al_2O_3 in the reference subsoil confirm the hypothesis that the subsoil can be regarded as a pristine, non-influenced, reference. A graphical representation of the relation between the various elements and Al_2O_3 for both topsoil and subsoil clearly depicts the relative addition in the topsoil. Using a linear regression model based on the subsoil data the extent of the enrichment was quantified for each topsoil sample. When expressed as the ratio between observed concentration and calculated baseline concentration, the elements in order of decreasing enrichment are $\text{Cd} > \text{Cu} \gg \text{Sn} \approx \text{Sb} > \text{Pb} > \text{Zn} > \text{As}$. These enrichments are in line with data found for a similar area in the north of France, except for Cu, which is relatively higher in the Zeeland topsoils. Spatial distribution patterns of the enrichment ratios show that for most

influenced elements regional variability is high and that hardly any spatial or mutual regional relations appear. The overall regional perspective provided by the contamination index shows that some areas in the north-east and centre of Zeeland on average are slightly more enriched.

Using the subsoil regression as a baseline has many advantages above the current Dutch practice of calculating a fixed legal threshold value, though corrected for clay and organic matter content. The natural random variability in baseline concentrations, expressed as the standard error of the regression, is a property of the model. This contrasts with the legal threshold value, which is just a rigid value with no range given. Also the regression model provides a local baseline, specifically suited for the parent soil material of Zeeland, which obviously should be preferred over a general model that tries to include all possible soil types. Compared to all other techniques presented, this subsoil regression baseline method gives the most profound insight into 1) the geological/pedological pattern of geochemical variability, 2) random natural variation, and 3) the anthropogenic imprint on the Zeeland soil. These advantages definitely outweigh the greater complexity of the procedure.

6 Regional diffuse geochemical patterns and processes

6.1 Introduction

From the perspective of environmental management it can be useful to identify areas in which the soil geochemistry is more influenced by human processes than others. For example, given that soil in the Netherlands and elsewhere is often used as site preparation and building material, the further spreading of contaminants can be curtailed when soil with a certain degree of enrichment of anthropogenic components is not used elsewhere. The current policy in the Netherlands regarding regional mapping of human influence is to create risk maps of soil pollution (van der Gaast et al., 1998) in which areas can be compared based on whether or not permissible limits are exceeded. However, this method considers neither the interactions between polluting substances nor how their concentrations relate to natural levels in the soil.

It is well known that soil composition is the result of both natural and human influences, and is thus a summation of separate processes, such as sedimentary deposition, weathering of calcite and other minerals, input by manuring, or atmospheric deposition. This multidimensional space comprises interactions over the region and between soil components, both interactions generally being expressed in the form of observable patterns, like the intercorrelation between different elements and/or spatial variability over an area. These patterns thus represent an important key for assessing the condition of soils, especially if a distinction can be made between human induced and natural patterns.

The previous chapters show that, given a dedicated sampling design and analytical procedures, it is possible with univariate and bivariate statistical techniques to make a distinction between sites dominated by natural or anthropogenic processes. This distinction was based on the comparison with threshold and reference values or explained variance using no more than two variables. Although such a small dimensional space may give an indication of the underlying processes, a larger dimensional, multivariate, space must be considered to fully understand the soil composition. This multidimensional space can be divided into attribute space, in which processes among chemical components play a part, and geographical space in which the interaction of these processes between sample locations is assessed. Ultimately, this answers the question as to what combination of which

processes occur where, thereby providing a practically useful explanation of the processes that causes geochemical soil variability.

In this chapter I focus on the extent to which human contributions actually affect the regional soil composition compared to natural variability. Using the same data as in chapter 5, the processes or factors determining the regional variability, both in attribute and geographical space are assessed. The regional expression of the interaction of human and anthropogenic processes is further evaluated, to see if regions may be distinguished where specific human processes have had an effect on the soil composition.

The general covariance structure of the data was studied using principal component analysis (PCA) to reveal the processes within attribute space. Groups of covarying geochemical components, may indicate certain processes. To assess which processes explain features in geographical space, clustering techniques were used to identify groups of sample points with similar properties. Due to the expectation that each pattern is a combination of different processes and the general multivariate nature of geochemical data, fuzzy c-means clustering (FCMC) was chosen (Vriend et al., 1988).

6.2 Material and methods

6.2.1 Study area and geochemical topsoil data

The study area belongs largely to a Holocene estuary and consists of the mainland of Zeeuws Vlaanderen, bordering Belgium, and several islands/peninsulas that have a long history of land reclamation and inundation. The Holocene geology and pedology of Zeeland have been extensively described in chapter 2 of this thesis. In summary the major soil types of Zeeland are immature marine clay deposits ranging from heavy clays (clay fraction larger than 25%) to sandy tidal inlet and tidal flats (clay fraction <7%). The Zeeland landscape is divided into Heartlands, the oldest and first reclaimed areas with often very heavy pool-clays, and Newlands, younger areas with more sandy clay soils. The elevation of the field level, relative to the Dutch average sea level NAP, ranges from about 1.5 m for channel ridges to -1.3 m for the pool clay areas. Land use is mainly agricultural and in the Newlands is characterised by its larger scale and intensive artificial drainage.

Figure 6.1 depicts the 270 sample locations where geochemical data were obtained. Each sample is a composite of 15 to 20 subsamples from the upper soil layer (about 0-30 cm), usually the plough furrow. The samples were analysed by X-ray fluorescence (XRF) in pressed powder tablet or, after aqua-regia digestion, by inductively coupled plasma-mass spectrometry (ICP-MS). The analysis aimed at maximising the number of parameters and from these a selection was made based on accuracy, precision and minimum of missing values. For details about the sampling, analytical methods and the selection procedure, see chapter 4 and table 4.11 (p. 80).




Figure 6.1. Topography of Zeeland and sampling locations. Names of peninsula's and areas in italic, cities in bold

6.2.2 Principal component analysis

To discern the controlling processes in attribute space principal component analysis (PCA), also called principal factor analysis, was used which is based on the major axes in a data cloud. These axes are derived from the covariance matrix by extracting the eigenvectors and subsequently deriving a matrix with factor loadings with an *a priori* chosen number of factors. Subsequently, to maximise the variance of the loadings on the factors, varimax rotation was applied (Davis, 2002). A principal component, or factor, is assumed to represent a controlling process or feature, that captures a specific relevant source of variance.

PCA can reduce the number of variables of the multivariate dataset to a few factors that may be indicative for certain geochemical processes. Factor analysis is widely used in geochemistry and usually with success (Reimann et al., 2002; Bianchini et al., 2002; Chork & Salminen, 1993; van Helvoort, 2003; Mol, 2002). The method has its controversies though and three well known are: 1) the choice of the underlying method for the factor analysis: principal factor analysis or maximum likelihood; 2) the choice of which elements to include and the number of initial factors for the analysis; and 3) normality of the data is assumed while this most often is not the case (Reimann et al., 2002; van Helvoort, 2003).

It was shown by Reimann et al. (2002) that for geochemical data principal factor analysis, based on the eigenvector structure is a better solution than maximum likelihood. They considered that as the maximum likelihood depended even more on normally distributed data than PCA, its use increases the chance of misleading results.

For the desired number of variables or geochemical components compared to the number of cases, the so called dimensionality, there is no standard rule of thumb. If p is the number of variables and n is the number of cases than p should be substantially smaller than n . Given the many available suggestions for p and n (e.g. $n > p^2$, $n > 9p$, $n > 3p$, etc.) (Davis, 2002; Reimann et al., 2002) The dimensionality in the present dataset, where $p = 30$ and $n = 270$, here defined as p/n , can be considered as sufficient.

The data used for this study can be regarded as “near-normal” (compare figure 4.6, p. 85 and 5.2, p. 106), with some elements displaying a certain amount of positive skewness and outliers, that might have an adverse effect on a standard PCA analysis. As presented by Pison et al. (2003) using a robust technique for PCA overcomes these adverse effects. This technique is based on deriving a robust covariance matrix using the minimum covariance determinant (MCD) algorithm of Rousseeuw & van Driessen (1999). This MCD algorithm tries to find a subset of observations whose classical covariance matrix has the lowest determinant, thus the highest covariances for all variables. This reduces the influence of outliers which usually result in lower covariances for most variables. Following Pison et al. (2003), $0.75n$ of the datapoints were used to calculate the robust covariance matrix, allowing a maximum of $0.25n$ of outliers without effect on the estimation of the covariance matrix. Standardisation of data to compensate for the difference in units and magnitude for both XRF (wt-% and ppm) and ICP-MS (ppm) is also provided by using the robust correlation matrix. To prevent indeterminacy effects on the XRF data, SiO_2 was omitted from the dataset.

The number of factors in principle is determined by a combination of the Kaiser/Guttman criterion (only factors with eigenvalues larger than 1 are incorporated in the model) and geochemical interpretability. Also scree plots are used, in which the eigenvalue is plotted against component number, to evaluate if a factor is relevant or not. It is known that limiting the set of factors with the Kaiser/Guttman criterion may be too crude and factors with eigenvalues just smaller than 1 should be at least evaluated for their relevancy (e.g. van Helvoort (2003)). Besides, a subselection of elements related to one factor can be used for a second PCA to gain insight if this factor comprises more than one process.

Additionally, the covariance structure of the components not included in the selected

dataset was estimated by calculating the correlation of these components with the PCA factor scores.

To provide insight in the regional variability of the processes inferred from the patterns in attribute space, the factor scores are spatially depicted. This may reveal which areas are particularly influenced by certain processes (Tao, 1998; Reimann et al., 2002). For spatial plotting of map attributes, the same plotting method as in chapter 5, based on the methods of Gustavson et al. (1997), is used.

6.2.3 Fuzzy c-means clustering

Although spatial presentations of PCA factor scores can reveal their regional variation, they only show the effects of the processes related to each of the factors, not the interaction between the processes. Yet, at each location the overall geochemistry is determined by a summation of the different processes described by the PCA. This interaction in geographical space is not revealed since the PCA is used for separating the processes regardless of their spatial location. The PCA procedure in principle results in independent (i.e. uncorrelated in the overall attribute space) processes, yet at each location they have some specific relation. To see if these interacting processes have a regional pattern, the sample locations with similar properties can be grouped together with a clustering technique.

Since the degree of interaction of the different processes varies gradually from one location to the other it is not expected that the grouping will result in clear, well separated clusters in attribute space but rather in dense areas in a continuous data cloud. Due to this expected fuzziness it was chosen to use fuzzy c-means clustering (FCMC). This technique has proved to be very useful in the field of soil science and geochemistry. (Vriend et al., 1988; Hanesch et al., 2001; McBratney & Odeh, 1997; Burrough et al., 1997; Kramar, 1995). With FCMC the interest is not to assign samples, or cases, unambiguously to a group but rather to give a similarity to each group in the cluster model. This similarity is expressed as a membership value ranging from 0, no similarity, to 1, identical to the cluster centre. The memberships of a sample for all clusters sum up to one. The advantage of this technique is that intermediate cases, falling in between clusters, can be recognized and are not forced into just one cluster. Outliers contribute equally to all clusters and since the calculation of the cluster centres uses a weighted membership, outliers have hardly any adverse influence on the final result.

The choice of elements for FCMC is more delicate than for PCA. Simply selecting all elements may result in additional noise for the clustering algorithm, but also that just one major process, e.g. parent material, determines the outcome of the clustering. Considering that the PCA differentiates the major processes determining the geochemical soil composition, then elements with high loadings on the corresponding factors should be representative for these processes. To prevent that the variability of one process alone will determine the outcome of the FCMC, a moderately equal number of elements of each factor is therefore selected.

The results of the FCMC are geographically plotted by using the cluster membership as the attribute for the maps, again using the approach of Gustavson et al. (1997). A sample was

regarded to belong to one cluster when exceeding a threshold value according to

$$\text{Threshold} = 0.7 \cdot \text{highest membership} > \text{next highest membership} \quad (6.1)$$

If the location did not meet this condition the location was regarded as belonging to more than one cluster.

6.2.4 Statistical software packages

Both multivariate analyses (PCA and FCMC) were applied using S-Plus and the R statistical software (Venables & Ripley, 2002; R Development Core Team, 2004). Using R, for the FCMC the `cmeans` function from the *e1071* library was used and for the robust covariance matrix the MCD version of the `cov.rob` routine from the *MASS* library was used. For the PCA an R implementation, based on the S-Plus `princomp` function was applied for extracting the eigenvalues and calculating the factor loadings, rotation was provided by the `varimax` function.

6.3 Results and interpretation

6.3.1 Principal components analysis

The first 8 eigenvalues, as extracted from the robust correlation matrix of the selected components included in the optimal dataset (see chapter 4), are plotted in figure 6.2. The first 4 components were considered to be geochemically relevant. They explain 76.6% of the total variability in all 30 components selected. The factor loadings are listed in table 6.1 and plotted in figure 6.3.

The first rotated factor explains the major part, 50.4%, of the variance and has the highest loadings for geogenic elements like Al_2O_3 , TiO_2 , Ce, La, Nb, and Y. Most additional elements, like the REE, are also related to this factor (table 6.2). It is interpreted as representing natural variability in the pristine soil composition. The component Al_2O_3 is related to Al-bearing phyllosilicates and known to have a close relationship with clay (Tack et al., 1997; Wilcke et al., 1998; Sharma et al., 2000). Many other elements are primarily influenced by the variability in clay content of the parent material and load high on this factor as well (van Gaans et al., 1995; Mol, 2002; Tebbens et al., 2000; Bianchini et al., 2002; Huisman et al., 1997). This relation has already been demonstrated in chapter 5. Relatively high loadings on this component were also found for the elements Cr, Ni, Zn, which within the Dutch situation are usually regarded as potential contaminants, but here the PCA indicates otherwise.

The second rotated factor, explaining 10.3% of the variance, shows high loadings of around 0.9 for CaO and Sr, with minor loadings for Mg and Mn. This factor was interpreted as representing the variability in carbonate content




Figure 6.2. Scree plot of the eigenvalues from the first 8, unrotated, components. The horizontal line indicates an eigenvalue of 1.

The third rotated factor explains about 8.9% of the variance, and contains mostly those elements that had already been identified as anthropogenically enriched in chapter 5. It is thus interpreted as representing variability of an anthropogenic input. The elements Cd, Cu, Sn, and Sb have the highest loading on this factor, while As, Pb, and Zn have minor loadings. As an additional element, Bi is to some extent associated with this factor as well (table 6.2). The loadings of K_2O , MgO , and P_2O_5 on this factor are suggestive of the role of fertiliser application in anthropogenic enrichment.

The elements with the highest loadings are the most enriched elements as well (table 5.7 p. 119). It should be noted that the elements that load high on this factor in general show low communalities indicating that they in part show unique behaviour. Zinc exhibits only a small loading on this 3rd factor while having a higher loading on the first factor. Also its communality is substantially higher, indicating that only a small part of the variability of this element is related to anthropogenic input. No distinct loading is shown for P_2O_5 on either factor 1, 2, or 3 but rather small loadings for each factor. The third factor suggests that anthropogenic processes form a distinct feature of the Zeeland topsoils, in univariate, bivariate and multivariate attribute space.

The last factor, explaining 7.0% of the variance, is related to Na_2O and Zr with SiO_2 as a correlated component (table 6.2). These elements are primarily associated with the silt

Table 6.1. Varimax rotated factorloadings and communalities. Method indicates analytical procedure, I=aqua regia digestion/ ICP-MS analysis, X=XRF analysis. Loadings between -0.25 and 0.25 are not shown for clarity.

	method	PC1	PC2	PC3	PC4	communality
Al ₂ O ₃	X	0.93				0.98
CaO	X		0.93			0.94
Fe ₂ O ₃	X	0.89			-0.31	0.98
K ₂ O	X	0.89		0.27		0.90
MgO	X	0.77	0.47	0.27	-0.26	0.95
MnO	X	0.59	0.51		-0.27	0.69
Na ₂ O	X		-0.38		0.65	0.63
P ₂ O ₅	X	0.31		0.26		0.22
S	X	0.49				0.31
TiO ₂	X	0.97				0.98
As	X	0.35	0.50	0.42		0.58
Ba	I	0.87				0.83
Cd	I	0.30		0.58		0.43
Ce	I	0.93				0.90
Cr	X	0.92				0.91
Cs	I	0.89			-0.29	0.91
Cu	X			0.59		0.41
Ga	X	0.92				0.97
La	I	0.95				0.93
Nb	X	0.96				0.94
Ni	X	0.85	0.32		-0.27	0.95
Pb	X	0.37	-0.31	0.46		0.46
Rb	X	0.91			-0.30	0.97
Sb	I			0.60		0.41
Sn	I			0.62		0.40
Sr	X		0.91			0.89
V	X	0.93			-0.28	0.97
Y	X	0.95				0.95
Zn	X	0.86		0.36	-0.29	0.95
Zr	X		-0.36		0.67	0.64
variance explained		50.4%	10.3%	8.9%	7.0%	

fraction and/or heavy minerals.




Figure 6.3. PCA loading plots of the selected elements. Dashed lines denote zero loadings.

The PCA shows that the major part, about 68%, of the variability in soil geochemistry is determined by natural processes related to a marine depositional environment and the distinction made by the PCA between respectively “clay”, “calcite”, “anthropogenic” and “silt” factor is in line with previous studies (van Gaans et al., 1995) and chapter 3 (table 3.2, p. 50).

The regional variation of the factor scores is depicted in figure 6.4. The plots of the clay, calcite and silt factor show the distribution of the geogenic components. The resemblance with the spatial distribution of Al_2O_3 , CaO , and Na_2O (compare figure 5.4, p. 109) is evident. The anthropogenic factor shows high scores in the North East of the province, at

Table 6.2. Robust correlation of selected elements with PCA factor scores. Correlations between -0.25 and 0.25 are left out for clarity. For information about precision, see chapter 4

	PC1	PC2	PC3	PC4	method		PC1	PC2	PC3	PC4	method
SiO ₂	-0.72	-0.63	0.22	0.39	X	Lu	0.88				I
Ba	0.59	-0.61		0.20	X	Mg	0.77	0.42			I
La	0.88	0.37			X	Mn	0.60	0.47	0.27		I
Nb	0.96				X	Nb		-0.29			I
Nd	0.87	0.34			X	Nd	0.93				I
Sc	0.88	0.44	0.49		X	Ni	0.87	0.26	0.37	-0.29	I
As	0.36	0.52		-0.27	I	Pb	0.70			-0.45	I
B	0.82			-0.31	I	Pr	0.93				I
Be	0.91		0.37	-0.32	I	Rb	0.92			-0.42	I
Bi	0.86			-0.27	I	Sc	0.90			-0.27	I
Ca		0.74			I	Sm	0.93				I
Cr	0.93		0.56	-0.28	I	Sr		0.88			I
Cu					I	Ta					I
Dy	0.92				I	Tb	0.92				I
Er	0.92				I	Th	0.91				I
Eu	0.92				I	Ti	0.89				I
Fe	0.92			-0.33	I	Tl	0.87				I
Ga	0.93			-0.31	I	Tm	0.91		0.26		I
Gd	0.93		0.42		I	U	0.78				I
Hf	0.26	-0.33			I	Y	0.93				I
Ho	0.90				I	Yb	0.90		0.29		I
La	0.94				I	Zn	0.80		0.42	-0.29	I
Li	0.90				I	Zr	0.65	-0.32			I

Walcheren and Noord-Beveland. This image is in line with the results obtained from the contamination index, as derived in chapter 5 (compare figure 5.11, p. 123), except that for Walcheren the extent of anthropogenic influence appears as in other areas while the contamination index indicates a smaller enrichment.

6.3.2 Anthropogenic subprocesses

Regarding the interest in the different subprocesses (manuring, atmospheric deposition, etc.) of the anthropogenic factor, the elements closely associated with this factor were studied in more detail. Both standard Pearson and MCD correlation matrix for the elements associated with the anthropogenic factor are shown in table 6.3. Robust and non-robust cor-




Figure 6.4. Regional distribution of rotated factor scores. Factor PC1 is clay factor, PC2 is calcite factor, PC3 is anthropogenic factor, and PC4 is silt factor. For explanation, see text.

relations are similar, except for Bi which was not analysed in survey 1. Robust moderate correlations group Cu with As, and Zn with Pb and Bi, and Cd with Sb. However, when a PCA is performed on only the elements of table 6.3, using the same method as for the complete dataset, no distinctive factor model could be derived. Apparently there are no clear distinguishable subfeatures within the covariance structure for the enriched elements, and apart from the one common factor, each anthropogenic element shows unique behaviour.

6.3.3 Fuzzy clustering

Based on the results of the PCA the elements (or their oxides) Al_2O_3 , TiO_2 , CaO , Sr , Na_2O , Zr , Sb , and Cd were selected for the FCMC, as they have the highest loadings on the PCA, have high communalities, and may be considered representative for the associated

Table 6.3. Correlation of anthropogenic elements. Upper part are MCD robust correlations, lower part are standard Pearsons correlations. All standard Pearsons correlations differ significantly from 0.

	As	Bi	Cd	Cu	P	Pb	Sb	Sn	Zn
As		0.08	0.32	0.58	0.38	0.21	0.42	0.48	0.50
Bi	0.54		0.26	-0.11	0.24	0.54	-0.02	-0.03	0.55
Cd	0.38	0.61		0.26	0.40	0.43	0.58	0.37	0.49
Cu	0.53	0.31	0.22		0.29	0.27	0.16	0.44	0.31
P	0.42	0.37	0.40	0.29		0.26	0.28	0.23	0.47
Pb	0.23	0.55	0.56	0.33	0.26		0.27	0.32	0.65
Sb	0.39	0.31	0.54	0.03	0.28	0.34		0.41	0.40
Sn	0.34	0.42	0.38	0.29	0.18	0.34	0.33		0.37
Zn	0.51	0.88	0.66	0.34	0.52	0.62	0.35	0.35	

process. For convenience they are called respectively “clay elements” (Al, Ti), “calcite elements” (Ca, Sr), “silt elements” (Na, Zr), and “anthropogenic elements” (Sb, Cd). The data were scaled to equal range after removal of extreme values larger than 3 times the interquartile range plus the third quartile (see equation 4.4, p. 70, $f = 3$). For determining the number of FCMC clusters the partition coefficient (F) and partition entropy (H) criteria (Vriend et al., 1988; Bezdek et al., 1984) were used together with *a priori* knowledge of the soils and earlier results presented in this thesis. A systematic approach was applied to vary the number of clusters between 2 and 6, and varying the fuzziness (q -value) from 1.2 to 1.7 trying to maximise F and minimise H . This resulted in an optimal model with 4 clusters and a q of 1.3.

The cluster centroids of the scaled data are shown in figure 6.5. The variability in centroids is high for the elements for the clay, calcite, and silt elements, while it is much smaller, especially between clusters 1,3, and 4 for the anthropogenic elements.

Figure 6.6 shows box-and-whisker plots of the concentrations of the components grouped by cluster. In this crisp presentation, the samples are assigned to the cluster for which they have the highest membership. This shows a gradual decrease in concentrations for the clay elements from cluster 1 to 4, with the largest gap between clusters 2 and 3. The calcite factor elements have significantly higher concentrations in cluster 1 and 3, and lower concentrations in cluster 2 and 4. The silt factor elements show a more or less moderate increase in concentrations from cluster 1 to 4, which confirms the inverse relation between clay and silt. Here two major gaps appear between cluster 1 and 2, and between cluster 3 and 4. The FCMC apparently divides the samples into 1) calcareous heavy clays, 2) non-calcareous heavy clays, 3) moderately calcareous silty clays, and 4) non-calcareous silts.

The anthropogenic elements (Cd and Sb) barely show differences between clusters, except




Figure 6.5. FCMC centroids of the scaled data.

for cluster 4 that has moderately lower concentrations. The latter probably reflects the naturally lower metal concentrations in this clay and carbonate poor, silty cluster. Exchanging Cd and/or Sb for other anthropogenic elements in the clustering procedure gives similar results with interquartile concentrations for the first three clusters within the same range. Adding more anthropogenic elements to the cluster model made no difference as well. Using Cd and Sb as representatives for human influence may thus be considered a sound choice. The distribution of concentrations for other anthropogenic elements is indeed similar to Cd and Sb (see figure 6.7), although the degree of departure from the natural pattern varies. Zinc, for example, still more or less follows the pattern of Al₂O₃, while the pattern for Sn is much flatter, without clear differences between the clusters. Copper, and also As,




Figure 6.6. Box and Whisker plots of the distribution of the data for 8 elements for each cluster. The y-axis denotes the concentration while the x-axis gives the cluster number.

appear to be relatively enhanced in the silty clay cluster 3, while Pb (like Cd) is highest in the non-calcareous heavy clays (cluster 2). I conclude that the extra variability caused by the anthropogenic enrichment in part counteracts the natural differentiation by soil types. The interaction of natural variability and human impact thereby does not lead to any cluster of samples that is discriminated solely on the basis of elevated anthropogenic enrichment. The median concentrations for all elements for each cluster are given in table 6.4.

Figure 6.8 shows the cluster memberships in geographical space. Their distribution can be compared with the regional distribution of Al_2O_3 , CaO and Na_2O as depicted in figure 5.3 (p. 108) and with the Zeeland soil map (compare figure 2.8, p. 33), since both clustering




Figure 6.7. Box and Whisker plots of the distribution of the data selected elements for each cluster. The y-axis denotes the concentration while the x-axis gives the cluster number.

and soil map have clay content as principal classifier (van der Sluis et al., 1965; Pleijter & Wallenburg, 1994). The calcareous heavy clays of cluster 1 are mainly found at south Noord-Beveland and in west Zeeuws-Vlaanderen. The non-calcareous heavy clay cluster 2 mainly shows high memberships in pool-clay areas at Walcheren and Schouwen-Duiveland. The Eastern part of Noord-Beveland, a silty soil area rich in CaO, has high memberships in cluster 3 (calcareous silty clays), while the non-calcareous silts of the 4th cluster show high memberships in areas at Schouwen-Duiveland and Tholen.




Figure 6.8. Regional distribution of the FCMC memberships of the sample locations. Samples are assigned to the cluster for which they have the highest membership. Samples not exclusively belonging to one cluster, see equation 6.1, are plotted in gray.

6.4 General discussion and conclusions

The consecutive use of principal component analysis (PCA) and fuzzy c-means clustering (FCMC) provided a profound insight into the process affecting the regional geochemical variability in the Zeeland soils and into the extent of the variability caused by human processes.

The PCA reveals 4 factors that explain 76.6% of the overall variance, these factors are

named for convenience “clay”, “calcite”, “anthropogenic” and “silt” respectively. The largest part of the variability (68%) is akin to geogenic clay, calcite, and silt related processes while 8.9% can be explained by anthropogenic processes. The elements related to the latter were the so called “heavy metals” Cd, Cu, Sb, Sn, and to a lesser extent Pb, Zn, and As. The regional variability of the clay, calcite, and silt factor resembles the regional variability of elements associated with this factor. The regional variability of the anthropogenic factor can be largely compared with the contamination index as obtained in chapter 5. When the elements of the anthropogenic factor were examined more closely, no separate subprocesses other than unique behaviour for each element could be discerned.

Considering that the soils of Zeeland are clearly enriched for some elements it is expected that this enrichment will also be revealed when looking at the processes within attribute space. The PCA confirmed this, however, the natural clay and calcite related processes still comprise the largest part of the soil variability. The anthropogenic influence on the soil only explains a small part of the variability emphasising that these anthropogenic processes only affect a small number of inorganic elements and are really an imprint on the overall natural variability.

The grouping of locations with FCMC, using the elements with the highest loadings of the PCA, resulted in an optimal 4 cluster model mainly dividing the area in calcareous heavy clays, non-calcareous heavy clays, moderately calcareous silty clays, and non calcareous silts. No clusters exist where samples are discriminated solely on the basis of anthropogenic enrichment. This suggests that the extent of these human processes, clearly imprinted on the Zeeland soils, are not related to certain regions but rather are diffusely distributed over the province.

Table 6.4. Median values for each FCMC cluster. See chapter 4 for information about precision and accuracy of the values.

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	method/unit
Al ₂ O ₃	10.46	9.79	7.68	6.98	XRF [wt-%]
CaO	6.37	1.55	3.58	1.53	XRF [wt-%]
Fe ₂ O ₃	5.06	4.44	3.33	2.77	XRF [wt-%]
K ₂ O	2.05	2.02	1.78	1.74	XRF [wt-%]
MgO	1.27	0.96	0.80	0.54	XRF [wt-%]
MnO	0.05	0.04	0.04	0.03	XRF [wt-%]
Na ₂ O	0.58	0.73	0.73	0.80	XRF [wt-%]
P ₂ O ₅	0.20	0.19	0.18	0.16	XRF [wt-%]
S	0.10	0.12	0.09	0.09	XRF [wt-%]
SiO ₂	63.58	71.25	74.03	79.23	XRF [wt-%]
TiO ₂	0.57	0.55	0.44	0.42	XRF [wt-%]
As	19.75	17.19	16.79	12.71	XRF [ppm]
Ba	274.87	311.45	282.56	296.70	XRF [ppm]

Continued on next page

Table 6.4 – continued from previous page

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	method/unit
Cr	91.08	89.40	74.43	71.25	XRF [ppm]
Cu	16.54	13.68	14.77	11.59	XRF [ppm]
Ga	11.93	11.25	8.68	8.03	XRF [ppm]
La	30.47	31.14	25.03	23.75	XRF [ppm]
Nb	11.66	11.87	9.76	9.51	XRF [ppm]
Nd	29.15	29.30	25.30	25.45	XRF [ppm]
Ni	26.78	21.93	17.45	14.38	XRF [ppm]
Pb	27.44	29.93	23.44	22.78	XRF [ppm]
Rb	96.12	88.19	72.03	65.78	XRF [ppm]
Sc	11.48	10.50	8.40	7.40	XRF [ppm]
Sn	4.29	3.62	3.97	3.44	XRF [ppm]
Sr	146.27	78.13	104.86	76.06	XRF [ppm]
Th	8.13	8.20	8.01	6.77	XRF [ppm]
U	2.17	2.48	2.03	2.14	XRF [ppm]
V	85.65	84.28	58.67	51.57	XRF [ppm]
Y	21.74	23.46	19.18	18.86	XRF [ppm]
Zn	66.40	65.57	49.11	42.92	XRF [ppm]
Zr	264.75	319.62	315.43	371.21	XRF [ppm]
Ag	0.08	0.08	0.07	0.06	ICP [ppm]
As	17.28	15.17	14.66	10.81	ICP [ppm]
Au	0.01	0.03	0.01	0.03	ICP [ppm]
B	50.24	47.69	39.32	34.98	ICP [ppm]
Ba	92.05	89.24	68.19	65.56	ICP [ppm]
Be	1.36	1.19	0.93	0.81	ICP [ppm]
Bi	0.21	0.23	0.17	0.15	ICP [ppm]
Ca	41173	10329	20860	9801	ICP [ppm]
Cd	0.29	0.32	0.29	0.25	ICP [ppm]
Ce	45.90	48.10	38.53	39.09	ICP [ppm]
Cr	48.53	46.54	35.35	32.01	ICP [ppm]
Cs	5.31	4.59	3.27	2.99	ICP [ppm]
Cu	13.17	11.04	12.03	9.90	ICP [ppm]
Dy	2.64	2.74	2.29	2.30	ICP [ppm]
Er	1.23	1.30	1.07	1.06	ICP [ppm]
Eu	0.81	0.85	0.67	0.66	ICP [ppm]
Fe	26750	24162	18555.85	15357	ICP [ppm]
Ga	7.31	6.47	4.77	4.39	ICP [ppm]
Gd	3.72	3.87	3.11	3.16	ICP [ppm]

Continued on next page

Table 6.4 – continued from previous page

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	method/unit
Hf	0.24	0.35	0.23	0.22	ICP [ppm]
Ho	0.48	0.50	0.40	0.41	ICP [ppm]
La	23.58	23.99	19.88	19.48	ICP [ppm]
Li	32.75	26.89	21.57	18.30	ICP [ppm]
Lu	0.17	0.17	0.14	0.15	ICP [ppm]
Mg	6549.81	5266.79	4710.06	3928.54	ICP [ppm]
Mn	311.60	229.09	227.63	187.15	ICP [ppm]
Mo	0.47	0.64	0.33	0.29	ICP [ppm]
Nb	0.57	0.66	0.60	0.57	ICP [ppm]
Nd	22.27	22.82	19.05	18.64	ICP [ppm]
Ni	22.35	19.82	15.35	13.45	ICP [ppm]
Pb	24.43	24.67	18.83	18.60	ICP [ppm]
Pr	5.58	5.68	4.81	4.70	ICP [ppm]
Pt	0.01	0.01	0.01	0.01	ICP [ppm]
Rb	62.61	56.20	40.15	38.29	ICP [ppm]
Sb	0.33	0.31	0.33	0.25	ICP [ppm]
Sc	7.89	7.66	5.97	5.60	ICP [ppm]
Se	0.41	0.98	0.34	0.64	ICP [ppm]
Sm	4.07	4.24	3.55	3.52	ICP [ppm]
Sn	2.08	1.96	1.92	1.60	ICP [ppm]
Sr	130.99	55.38	82.91	51.59	ICP [ppm]
Ta	0.03	0.02	0.03	0.02	ICP [ppm]
Tb	0.53	0.54	0.44	0.44	ICP [ppm]
Th	5.91	6.26	5.13	5.15	ICP [ppm]
Ti	702.83	727.14	630.68	620.85	ICP [ppm]
Tl	0.36	0.35	0.26	0.28	ICP [ppm]
Tm	0.17	0.18	0.14	0.14	ICP [ppm]
U	1.06	1.09	0.94	0.93	ICP [ppm]
Y	12.12	12.90	10.37	10.22	ICP [ppm]
Yb	1.09	1.11	0.92	0.94	ICP [ppm]
Zn	70.08	64.91	55.15	51.26	ICP [ppm]
Zr	11.97	14.26	10.12	9.44	ICP [ppm]

7 Assessment of regional DDT concentrations in the soils of Zeeland

7.1 Introduction

Since the 1940s the organochlorine pesticide DDT has been used in western Europe, and despite its ban in the 1970s its legacy is still present as residues in the soil compartment. For several decades its risks for the environment have been acknowledged (Organisation, 1989; Edwards, 1970) and this makes it important to have a good overview of the occurrences of DDT in our environment. Especially for soil, as a major environmental compartment, and being both a source and a sink, knowledge about DDT residue concentrations is vital to evaluate the ecotoxicological risk of our notorious legacy. One of the questions in such an evaluation is whether the DDT residue level is part of the ubiquitous global background or is the result of more local features.

Studies in the Arctic and Antarctic, glaciers, tundras, i.e. pristine areas from where it can be assumed that DDT was not actively applied, show significant concentrations in soils, sediment and water (Rovinsky et al., 1995; Harner et al., 1999; Villa et al., 2003; Politov et al., 2000). Most of these studies regard marine and atmospheric transport as the main transportation mechanisms of DDT. Considering this ubiquitous occurrence of DDT one can assume the existence of a global DDT background. However, this background is neither geogenic nor biogenic but solely anthropogenic and there are no reasons to assume that a natural source of DDT exists.

Knowledge about DDT residue concentrations can be provided by large scale regional soil surveys, for example, as are common in geochemistry (Darnley, 1995; Bölviken et al., 1996) However, examples of large scale surveys aimed at organochlorine pesticides (OCP), including DDT, are scarce (Gong et al., 2004). More often soil surveys aimed at DDT concentrations are regional and have a limited number of samples, usually about 10 to 50 samples (e.g. Covaci et al. (2001); Kim & Smith (2001)). When speculating about the reasons why OCP surveys are as yet less common than geochemical surveys, both would have similar environmental goals (Darnley, 1995), the time, complexity, and effort that are needed for the analytical determination of OCP, appear likely culprits.

Also within the Netherlands interest in the general level of organochlorine pesticides occurrence in soil exists, but only small parts of the country have been sampled and the number

of samples is usually limited (Groot et al., 2001). A country wide, or even regional scale, survey of DDT concentrations does not yet exist, despite the fact that at the local scale DDT occurrence in soils frequently results in legal issues (e.g. building permits, sale of land).

An alternative approach to obtain regional information about DDT or other contaminants is to use existing data. A larger regional assessment can be obtained by combining data from several subregions, which are gathered over the years (Licht & Tarvainen, 1996). Ideally, these data should have been sampled and analysed using standardised methods to minimise survey and analytical bias. Even then the combined dataset may still suffer from bias caused by different choices during survey and analytical procedures. However, by acknowledging the presence of bias, the data may still provide relevant information, and using limited resources and effort an indication can be obtained if larger or more detailed studies are necessary.

In the Netherlands, municipal and provincial authorities are currently putting much effort in drafting so called soil pollution risk maps (van der Gaast et al., 1998; van Lienen et al., 2000). These maps are used for environmental legislation, building permits, and permits for transport of (non-polluted) soil and are to some extent standardised regarding sampling and determination of chemical composition (VROM, 1999; Eikelboom et al., 2001). The data for these maps are held in soil information systems (SIS) and include some information about OCP, including DDT and residues.

This study aims at providing a regional assessment of current DDT residue concentrations, variability, and degradation for the province of Zeeland. Data from several separate SIS data are merged. By comparing the concentrations with legal limits and data from other areas the extent of contamination will be assessed, providing insight into whether the concentrations are just part of the ubiquitous background or exceeding those. The geographical distribution of the DDT residue levels might indicate patterns, if any, that possibly relate to distinct processes. In addition, the data integration exercise will assess the value of the combined SIS for regional environmental policy and may lead to recommendations to the (local) authorities for further research and monitoring.

7.2 DDT, properties and history

DDT^{*}, and its metabolites DDE[†] and DDD[‡] are well known as contaminants in our present day environment (Braune et al., 1999). The physiochemical properties of these substances make them a real environmental threat as they are resistant to breakdown, easily taken in by organisms, and are readily absorbed on to the soil (organic) matrix, and they have hazardous effects on aquatic and terrestrial organisms (Organisation, 1989; Zitko, 2003). Soil can both be a sink and a long-term source for exposure to DDT. This not only affects soil biota (Megharaj et al., 1999) but can also affect larger organisms (Zitko, 2003). Al-

^{*} 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane

[†] 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene

[‡] 1,1-dichloro-2,2-bis(p-chlorophenyl)ethane

though absorption is often assumed, DDT is also reported as soil-bound and unextractable by solvents (Singh & Agarwal, 1992).

The metabolites DDE and DDD are formed by the breakdown of DDT. It is assumed that the degradation of DDT in aerobic soils proceeds relatively slowly, with reported half-lives ranging between 20 to 30 years (Howard, 1991; Dimond & Owen, 1996)

DDT has been in use since the beginning of the 20th century, as part of a larger group of OCP that were seen as new state of the art methods to engineer nature. It was not until the emotional call in the book “*Silent Spring*” (Carson, 1962) that awareness about the deleterious effects of pesticide usage in society was raised. Since the beginning of the 1970s DDT usage has been banned in most industrialised countries, in the Netherlands DDT has not been allowed since 1973.

DDT, DDE and DDD each have two different isomers depending on the position of the phenyl-bound chlorine. When data for the individual isomers are presented they will be reported as *p,p*-DDT and *o,p*-DDT (or *p,p*-DDE/*o,p*-DDE, and *p,p*-DDD/*o,p*-DDD). When purity is unknown the simple abbreviations DDT, DDE, and DDD will be used in a general sense. All isomers and metabolites together will be referred to as DDx in a general sense or $\sum DDx$ when quantitatively indicating the sum of all concentrations.

7.3 Study area

The location of the study area is shown in figure 7.1. The province of Zeeland is located in the south-west of the Netherlands along the coast of the North Sea. Besides the mainland of Zeeuws-Vlaanderen, it consists of several islands and peninsulas. The area is the result of a continuous shaping by the marine environment and human activity. It has been more extensively described in chapter 2.

Zeeland has an agricultural land use history where crop farming has long been the most important factor. Nowadays the total area of 2930 km² consists of 1140 km² water and 1440 km² agricultural area. Common crops are corn, root- and tuberous plants while livestock are mostly sheep and some cattle. In the 20th century the economy shifted towards more industrial and service activities and today only 4% of the economic revenues are gained from agriculture even though it still encompasses the largest geographical area (see also chapter 2). Based on communications with the provincial authorities, it is known that DDT has been used for pest control for several years in Zeeland, especially in orchards. This usage ended in the mid 1960s. Still the concentrations of DDT and derivatives existing in the soil is of concern for local authorities. The legal limits, set up by the Dutch government, are often exceeded, resulting in litigations.




Figure 7.1. Topography of Zeeland. Names of peninsula's and areas in italic, cities in bold

7.4 Method

7.4.1 Collection of data

Data were obtained from several databases of soil information systems (SIS) maintained by the municipal and local governments in Zeeland. These SIS are basically intended for management of soil pollution and soil remediation and therefore primarily information about environmental priority compounds is recorded. Each SIS holds data for a distinct area of the province usually bound by municipal boundaries.

The available databases were merged and only topsoil (0-50 cm depth) data from rural areas, mainly agricultural land, were selected. These georeferenced data comprise several inorganic parameters like priority metals, and organic parameters such as polycyclic aromatic hydrocarbons, polychlorinated biphenyls and OCP including DDx. Some databases contained replicate samplings which will be used to assess the variability at different sampling scales.

Sampling and analytical methods are generally based on national standards for which the NEN 5740 standard for sampling (NNI, 1999b) and the NEN 5734 standard for OCP analysis (NNI, 1999a) are the most relevant in this context. The sampling in general comprises taking composite samples containing about 10 to 20 subsamples, taken from a distinct range from the soil profile. Depending on the purpose, scale, and if contamination is expected or not, the number of subsamples and sampling density varies. The depth range of the samples usually varies from 1-30 cm to 1-50 cm depth for the topsoil and a second sample from a depth of 50-100 cm.

The OCP analysis, according to NEN 5734, is based on a wet extraction with acetone and petroleum ether. The extracts are dried with anhydrous Na₂SO₄ and eluted over an aluminum oxide column. Subsequent gas-chromatographic analysis, mostly with a mass-spectrometric detector, is used for detection of the residues. The chemical analyses are mostly carried out in commercial laboratories.

Although standard methods are used, small deviations are to be expected due to different interpretations of the protocol or adaptations to local circumstances, during both surveying and laboratory analysis. These departures are usually not recorded in the datasets. No method to compensate for the resulting bias is used since neither control samples were available nor were analytical standards reported in the SIS. Other known methods for removing bias between datasets are not suitable due to the low number of samples per survey (Daneshfar & Cameron, 1998; Darnley, 1995).

7.4.2 Basic statistical analysis

The statistical analysis should provide insight in the ranges of DDx concentrations, while being robust against the expected high variability within the data, many outliers, and censored data below the detection limit occur. Considering the possible bias between the different datasets, the analysis should also provide some basic insight into the differences between the datasets.

Univariate data exploration techniques, such as cumulative distribution frequency (CDF) plots, were used to provide the necessary insight in the statistical DDx distributions. With CDF plots, in which the data are plotted against the data percentiles on a probability scale, the type of data distribution is determined and possible outliers can be identified. This technique has already been demonstrated in chapter 5 and has been also used by other authors (Sinclair, 1976; Matschullat et al., 2000; Bounessah & Atkin, 2003). Standard histograms were viewed as well.

Since outliers might denote important features, such as areas with more extensive DDT

usage, they are as important as the bulk of the data. Therefore outliers are treated separately when necessary. The CDF plots were used to find a threshold value above which a value should be regarded as an outlier. This threshold value is the percentile where a clear infection in the CDF curve occurs.

When replacement of censored data, i.e. values below the detection limit, was necessary they were replaced by a value of 0.7 times the detection limit.

7.4.3 Analysis of variance

The observed variability of DDT and its derivatives in soil comes from several sources. For example variability in application, degree of breakdown, transport through the soil column, together with the variability caused by sampling and analytical procedures may all contribute to the total variability. In chapter 3 it is already shown how the contribution of local scale variability can influence the regional variability. If the local variability is relatively high then it is to be expected that observed regional patterns are mainly caused by local features instead of true regional processes.

In some datasets replicate samples exist, i.e. some areas or fields were purposely sampled more than once within a survey, or samples were analysed by two different laboratories. The occasional replicates in the dataset were used to calculate local variance (i.e. fieldscale) and this can be compared with the overall regional variance. Using an analysis of variance (ANOVA) the variance components can be calculated, which explain the contribution of each level of variance to the total variance (Youden & Mehlich, 1937; Nortcliff, 1978; Oliver & Webster, 1986; Webster & Oliver, 1990; Searle et al., 1992). For the ANOVA and calculation of the variance components the raov function of *S-PLUS* was used (Venables & Ripley, 2002).

Since not all datasets have a sampling design suitable for ANOVA, and considering the interest in the overall spatial variance as well, an experimental semivariogram was calculated by using all available data (Journel & Huijbregts, 1981). By using the range of the variogram the extent of the local semivariance can be compared to the regional one and it can be assessed how far local noise determines the observed regional variability. The semivariogram was calculated using the *Gstat* program (Pebesma & Wesseling, 1998). Both ANOVA and semivariance can provide insight into how the observed regional variability should be interpreted.

7.4.4 Regional variability

To reveal spatial DDx patterns over the region of Zeeland, the concentrations were plotted using symbol maps. From these, groups or areas with similar concentrations can be discerned. Spatial anomalies, single samples with anomalous concentrations compared to the surrounding samples, can be recognised as well. Examples of symbol maps, for which the size of the symbol varies with the associated attribute can be found in Bounessah & Atkin (2003) and Reimann & Filzmoser (1999). In chapter 5 (§5.3.2) the plotting method of

the Geological Survey of Finland is described and demonstrated (Gustavson et al., 1997), which will also be used here.

7.4.5 Comparison with external data and normative values

Considering that there is no natural, geogenic, background for DDx it is more difficult to evaluate the soil concentrations of Zeeland than in the case of potentially toxic metals. To put the data in perspective, values found in other areas, mostly outside the Netherlands, are compared with those found in Zeeland. Although it is expected that the sampling and analytical methods of the other studies will often differ from those for the current study, the comparison will still point out the relative magnitude of the DDT contamination of Zeeland.

The values are obtained from literature for areas which were presumed to be to some extent comparable to Zeeland or from pristine areas where it can be assumed that no DDT application has occurred. It is recognised that the probable differences in soil lithology and soil ecology also will lead to differences in ecotoxicological risk between areas.

In the Netherlands it is standard practice to compare DDx values with soil specific legal limits to judge the extent of the contamination, using the limits as a standard reference. These limits are provided by the Dutch normative framework (VROM, 1994; Swartjes, 1999) and include a threshold value (*S*, or “Streefwaarde”), which is the maximum permissible level for sustainable soil quality, and an intervention value (*I* or “Interventiewaarde”) which indicates the level of severe pollution. These values apply to a so called standard soil, as defined in the framework, containing 25 % of clay and 10 % organic matter (VROM, 1994). The *S* and *I* values for standard soil for $\sum DDx$ are 0.01 mg/kg and 4 mg/kg respectively. For pesticides the normative framework requires correction for the actual organic matter content in the soil according to

$$s = S \cdot \frac{O}{10} \quad (7.1)$$

Here *S* is the value for standard soils and *s* is the soil specific legal limit normalised on percentage organic matter, *O* (VROM, 1994). The comparison of DDx concentrations against these limits does not take into account sample and analytical variability and this can make deciding between a “clean” soil (concentration < *S*) and “polluted” soil (concentration > *S*) a chance event.

7.4.6 DDT breakdown

If the extent of DDT breakdown calculated from the actually observed values in Zeeland for metabolic and remaining DDT is less than expected under local circumstances, compared with a half-life of 20-30 year (Howard, 1991; Dimond & Owen, 1996), then the DDT breakdown has evolved more slowly than expected, or fresh application has occurred. Also,

it indicates that the Zeeland soils will be a source of DDx components for a longer period than probably anticipated. Presuming that DDT was banned at the beginning of the 1970s it can be expected that DDT breakdown should now be at a level of at least 1 half-life time. As no single validated value for DDT half-life exists, the exact time elapsed since its application cannot be assessed. However, from the molar ratio between DDT and its breakdown products, DDE and DDD, it can be calculated how many half-lives ($\frac{t}{T_{\frac{1}{2}}}$) have passed. Assuming exponential decay the formula

$$\frac{t}{T_{\frac{1}{2}}} = \frac{\ln\left(\frac{[DDE+DDD]}{DDT} + 1\right)}{\ln(2)} \quad (7.2)$$

can be used, as was suggested by van Gaans et al. (1995).

As DDE is the major breakdown product under aerobic conditions, which is the predominating condition in soils, DDE is likely to be the predominant metabolite to be found in Zeeland.

7.5 Results

7.5.1 Data merging and selection

It became apparent that neither a common standard nor a well described method was used when recording the data in the different databases. The recording of the data itself was regularly quite minimal. For instance, in many datasets only a sum DDT value was recorded instead of separate values for DDT, DDE, and DDD. Information about which isomer was determined (*o,p* or *p,p*) was even more scarce. In general, neither information about sampling strategy nor analytical procedure or laboratory was available. Therefore it was decided to use the data “as is” and not to make assumptions about the differences between the methods.

From the initial merged dataset, locations within the rural area of the province were selected, and from each location the sample, if any, for which DDT, DDD, DDE or $\sum DDx$ was determined, was selected. Locations known to be orchards were left out, as these were suspected of showing divergently higher values. This resulted in a final selection from 7 databases and 30-60 samples per database (table 7.1). The sample locations are reasonably well distributed over the province although not every area is covered (figure 7.2).

All surveys, except surveys 1 and 3, were derived from datasets related to local pollution risk maps. Surveys 2, 5, and 6 were distinct surveys specifically aimed at creating a pollution risk map while surveys 4 and 7 are in themselves already a combination of several subsurveys with different aims.

Survey 1 had a somewhat different sampling design than proposed by the standardised design according to the NEN 5740. A composite topsoil sample, consisting of 5 randomly

Table 7.1. Databases and survey area used for merged dataset

Number	Area	samples
1	Koegorspolder (Terneuzen)	45
2	Schouwen-Duiveland	30
3	Tholen, Zuid-Beveland	35
4	Walcheren	38
5	Borsele (Zuid-Beveland)	40
6	Zeeuws-Vlaanderen	30
7	Noord- and Zuid Beveland	59
Total		277

located subsamples, was taken in triplicate from a field ($\pm 1-2$ ha) for 15 adjacent fields (total area approximate 400x500 m). The aim of the sampling design was to obtain information about both soil concentrations and field variability, and to construct a pollution risk map. In this study the data can be used to gain information about the variability at the moderately local scale.

Survey 3 was part of the geochemical mapping of Zeeland as presented in this thesis. The sampling design is comparable to the NEN 5740 and is extensively described in chapter 4. Sample pre-treatment and analysis of DDT for these samples were carried out in a commercial laboratory similar to those engaged in the surveys related to pollution risk maps. For a separate study (Van Gaans et al, in prep.), 12 samples with $\sum DDx$ values above the detection limit were re-analysed in our own laboratory, including replicates to assess within laboratory error. In this study these samples are used to assess the between laboratory variability.

In survey 5, for 10 locations a NEN 5740 based sampling design was carried out in duplicate. A composite sample was taken from 20 subsamples on a rigid 40x50 m grid. The duplicate sample was taken from an identical grid, displaced by 1 m from the original. With these data it is possible to obtain insight into the short scale variability compared to the larger scale of the total survey.

7.5.2 Exploratory statistics

Table 7.2 gives summary statistics of the data used in this study. This table shows that the number of data on DDE and DDD is limited. Information on different isomers (e.g. *o,p*-DDT and *p,p*-DDT) is only available for survey 1 and 3; the sum of the isomers are reported in table 7.2. The $\sum DDx$ concentrations are the only data that are available for all the datasets. Therefore the regional assessment concentrates on $\sum DDx$.

After studying the distribution of the data by means of boxplots, CDF plots and histograms, a logarithmic transformation (to base e) was used to obtain a more normal distribution.




Figure 7.2. Selected sample locations from the soil information systems of Zeeland. The location of survey 1 is depicted as a single icon, but consists of 45 sample locations in a small area (about two times the size of the area covered by the icon).

Figure 7.3 shows boxplots for each survey and for the overall data on a logarithmic scale. The large differences in distributions and medians, as well as the presence of many outliers, are evident. Surveys 2 and 4 have relatively low concentrations of $\sum DDx$ while survey 5 has relatively high concentrations. Since the surveys are restricted to administrative boundaries, except that survey 3 and 7 have some overlap, the differences in concentrations might indicate regional differences. However, since sampling and, especially, analytical techniques also might differ per survey, it is very likely that the differences are at least partly caused by bias.

Figure 7.4 depicts the CDF-plot of the $\sum DDx$ concentration on a logarithmic scale. It shows that about 12% is censored data due to detection limits, and a clear inflection point is visible around the 90-percentile, or P_{90} . Based on this inflection point the values larger than the P_{90} , corresponding to a concentration of 0.0945 mg/kg, will be considered as outliers. The other data, non-censored and non-outlier, show an almost straight line in the CDF plot

Table 7.2. Summary values for DDDx[mg/kg] and total organic carbon [wt-%] (TOC). Format is (minimum:maximum/median). NA is no available data. Italic numbers are detection limits.

survey	DDD	DDE	DDT
1	(<i>0.001</i> :0.025)/0.002	(0.001:0.025)/0.016	(<i>0.001</i> :0.057)/0.016
2	NA	NA	NA
3	(<i>0.001</i> :0.005)/ <i>0.001</i>	(<i>0.001</i> :0.026)/0.003	(<i>0.01</i> :0.050)/0.007
4	NA	NA	NA
5	NA	NA	NA
6	NA	NA	NA
7	NA	NA	NA

survey	Σ DDDx	TOC
1	(<i>0.001</i> :0.083)/0.028	(0.6:2.6)/1.5
2	(0.001:0.850)/0.002	(2.5:6)/3.6
3	(<i>0.001</i> :0.090)/0.009	(3.3:9)/4.7
4	(<i>0.001</i> :0.070)/ <i>0.001</i>	(3:96)/4.5
5	(<i>0.001</i> :1.100)/0.028	(2.1:7)/4.2
6	(<i>0.001</i> :0.060)/0.010	(2.3:6.2)/4.05
7	(<i>0.001</i> :0.850)/0.015	(1.1:15.6)/3.6

confirming that a log-transformation is a sound choice.

7.5.3 Local and regional variability

The results of the ANOVA are shown in table 7.3 and the variance components, ordered by approximate scale, are graphically depicted in figure 7.5. All data used were below the P_{90} value of the total dataset (comp. figure 7.4), detection limits were multiplied by 0.7.

To compare estimated variances between the surveys, recalculation of the variance components was necessary. Survey 1 had only 5 subsamples for each composite while other surveys had 15-20 subsamples per composite. Therefore in figure 7.5 the recalculated within field estimated variance is based on the \sqrt{n} ratio of the original estimated variance. The between field variance of survey 3 was recalculated to compensate for the decreased variability by leaving out the samples with values below the detection limit.

From figure 7.5 it is clear that the between laboratory variability is much smaller than the variability on even the smallest field scale, implying that analytical variance is sufficiently low. The within field variability is always a large component compared to between field variability, while between field variability clearly increases with increased scale of the sub-region covered in the survey. Since the largest scale indicated (15x15 km) is still much




Figure 7.3. Boxplots of log scaled concentration $\sum DDx$ by survey. The upper dashed line denotes the overall P_{90} (see figure 7.4) and the lower dashed line the legal, non soil specific, S limit. The total of all surveys together is shown in the last (T) boxplot.

smaller than the province as a whole, overall variance is expected to be even larger.

Comparing the variability of the concentrations of $\sum DDx$ to the variability of inorganic compounds, e.g. As and metals like Pb, Cd, Zn as obtained in chapter 3, it is clear that the variability of $\sum DDx$ is substantially larger. Given the fact that the samples from the study described in chapter 3 are single samples, while the samples from the datasets of this study are composite samples, it can be expected that the actual small scale variance is even higher.

For the calculation of the semivariogram of the log transformed concentrations $\sum DDx$ outliers were removed based on the P_{90} . It is expected that these outliers result in an overestimation of the regional semivariance. The influence of censored values below the detection limits was assessed by both calculating the semivariance with and without these values. The censored values might result in an underestimation due to lack of variability between these samples. However, such samples indicate very low or absent $\sum DDx$ concentrations which are an important feature of the regional variability. Therefore it was decided to calculate the semivariogram both with censored data, replaced by 0.7 times the detection limit, and without the censored data.

The calculated variogram is given in figure 7.6 and it shows that the nugget effect for the




Figure 7.4. Cumulative frequency plot of the concentration $\sum DDX$ [mg/kg]. Percentiles are plotted along the X axis and concentrations along the logarithmic Y axis. The line depicts the P_{90} for the X-axis and the corresponding value ($94.5 \mu\text{g}/\text{kg}$) on the Y-axis.

data without the censored values is nearly as high as the sill. This means that at small scales already the regional semivariance is reached and that the regional variability is mainly determined by small scale variance. For the data where the censored values were replaced, the overall semivariance as expected is larger. Also the difference between nugget and sill seems a bit more pronounced, probably due to the larger number of sample pairs at small scales and the larger overall semivariance.

7.5.4 Regional overview

Figure 7.7 shows the regional overview of the $\sum DDX$ concentrations in Zeeland. This overview shows large regional and subregional variance in concentrations while the outliers appear to be relatively concentrated in the south-east of Zuid-Beveland. The high values in this area might be related to the presence of orchards, which might still be represented

Table 7.3. Results ANOVA on the replicated samples. DF=degrees of freedom, sum sq=sum of squares, mean sq=mean square, Est. var=variance component.

Survey 5	DF	sum sq	mean sq	est. var.	approx. scale
Between fields	11	49.73	4.52	1.92	15x15 km
Within field	12	8.25	0.69	0.69	40x50m

Survey 1	DF	sum sq	mean sq	est. var.	approx. scale
Between fields	14	36.26	2.59	0.46	1.5x2 km
Within Field	30	36.36	1.21	1.21	100x200m

Survey 3	DF	sum sq	mean sq	est. var.	approx. scale
Between fields	11	9.40	0.85	0.40	10x10 km
Between labs	1	0.51	0.51	0.04	–
Error	11	0.33	0.30	0.03	–

in the dataset due to incomplete information in the SIS, or which could have influenced the general level in the area. Relative low concentrations are found in Walcheren and at Schouwen-Duiveland. Similar low concentrations, together with high concentrations, are visible in the south-east of Zuid-Beveland. Variability appears less in areas like Noord-Beveland and Zeeuws-Vlaanderen.

However, considering the results of the ANOVA and the semivariogram which indicated that local features are a large component in the regional variability, the distribution of $\sum DDx$ concentrations over the area should be judged with some precautions. Together with the differences in $\sum DDx$ concentrations between surveys (see figures 7.2 and 7.3) the intrinsic value of the regional overview is limited.

7.5.5 Comparison with external data and normative values

Table 7.5 gives values of DDx levels from other natural and agricultural areas around the world for comparison with the $\sum DDx$ values observed in Zeeland. This shows that the DDx residue levels of a former study in Zeeland has substantially higher values than this study (van Gaans et al., 1995). This might indicate that the DDx residue levels are lowered due to breakdown but it is more likely, based on the DDT breakdown values as presented below, that this illustrates variability resulting from different research approaches. The variability among all the studies in table 7.5 is also evident. The one study of orchards show relatively high values compared to the other areas. The study of Cavanagh et al. (1999) show relative low values and as they report probably related to the just occasional DDT usage.

Considering the overall median values for DDT and $\sum DDx$, both 0.01 mg/kg, the extent of




Figure 7.5. Variance versus increased approximate scale. Upper graph shows the cumulative variance component, lower graph shows the variance components for each scale. Surveys are denoted between brackets. *) recalculated for a composite sample of 15-20 subsamples instead of 5 as used in the Koegorspolder, **) recalculated for the total dataset of 35 samples instead of the subset of 12 selected samples (original values as unfilled bars).

the contamination in Zeeland seems limited. Concentrations in natural areas, i.e. a tundra and the Tibetan high plateau (Rovinsky et al., 1995; Fu et al., 2001), are roughly a factor 2 to 10 lower than the concentrations in this study.

To compare the $\sum DDx$ concentration with the Dutch legal limit, it should be adjusted for organic matter content according to the regulations of the normative framework (VROM, 1994). Figure 7.8 shows the relation of the $\sum DDx$ with the total organic carbon content




Figure 7.6. Semivariogram of the regional ΣDDx concentrations. Numbers indicate the number of pairs.

(TOC). Outliers and samples with values below the detection limit were removed from the plot. From the graph it is clear that they are unrelated, which is confirmed by the correlation coefficient of 0.001. Survey 1 is clearly different from the other surveys, probably the TOC variability is smaller due to the smaller size of the sampled area.

Without a relation between ΣDDx and TOC the correction of the S values to soil specific values seems less obvious since this correction is based on the assumption of such a relationship. The normalisation could still be applied based on the arguments of the ecotoxicological risk associated (Swartjes, 1999). However, it was decided not to apply the correction for organic matter in this study, as it is concluded that for a regional inventory of soil concentrations the usage of the normalisation procedure will in part obscure the results. Table 7.4 lists the percentage of sample points exceeding the S value. Roughly half of the samples have ΣDDx concentrations larger than S . For DDT almost half the samples exceeds S , while the number for DDE is somewhat lower and few samples exceed for DDD. Note




Figure 7.7. Regional overview of $\sum DDX$ concentrations. Normal values are gray, outliers (concentrations $> P_{90}$) are black (see also figure 7.2).

that $\sum DDX$ is a summed value; if one of the components (DDD, DDE, DDT) exceeds S , $\sum DDX$ will exceed also. As is evident from figure 7.8, the number of samples exceeding the soil specific s limit will be even higher, in the order of 75 %.

Table 7.4. Percentage of sample points exceeding the S -value.

	DDD	DDE	DDT	$\sum DDX$
percentage $> S$	3 %	28 %	49 %	50 %




Figure 7.8. Relation of organic matter (TOC) and $\sum\text{DDX}$ with outliers and not detected values not shown. The solid line depicts the S value while the dashed line shows the soil specific $c s$. The correlation coefficient r is 0.001

7.5.6 DDT breakdown

For survey 1 and 5 not only is the $\sum\text{DDX}$ available but also the separate parameters for DDT, DDE and DDD (both o,p and p,p), their relation is plotted in figure 7.9. This shows a strong correlation ($r=0.935$) between DDT and DDE and, using the medians, a DDE:DDT ratio of 0.5. The relation between DDT and DDD is less evident. For estimating the number of half-lives passed censored data were recalculated using equation 7.2.

By calculating a $\frac{t}{T_{1/2}}$ for each available sample, a range of values is obtained. The P_{25} , median, and P_{75} are 0.4, 0.6, and 0.8 respectively. It appears that the higher DDT values tend to a $\frac{t}{T_{1/2}}$ of 0.5, while lower values have higher $\frac{t}{T_{1/2}}$ (figure 7.2). This indicates that the residue level is only about one half to two-thirds $\frac{t}{T_{1/2}}$ and implies that the breakdown is not at the expected rate or that DDT applications occurred after the ban in 1973. The latter,

however, is very unlikely at such a large scale.

7.6 Discussion and conclusions

The combination of the different SIS derived datasets resulted in a database that can be used for a regional assessment. However, the differences between the surveys, although in general based on the same sampling and analytical standards, are evident in the data. It is hard to determine if these differences denote differences in true $\sum DDx$ soil values or that they are largely caused by the slight variations in field and laboratory methods.

The $\sum DDx$ concentrations as found in Zeeland are comparable to those in similar agricultural areas, and based on these values there is no reason to believe that the current level of DDT contamination is more severe than can be expected for an agricultural area. Compared to pristine areas the level in Zeeland is about a factor 2 to 10 higher, which indicates Zeeland can be regarded as enriched compared to a general ubiquitous $\sum DDx$ background. Despite that the values are within a normal range for agricultural areas, the standard soil normative threshold value S is exceeded for roughly half the cases and the soil specific value s in even more cases, which may result in litigations.

The relatively large contribution of local scale variability to the overall variability of $\sum DDx$ concentrations implies that a regional overview has limited value since most of the observed variance can be regarded as noise caused by small scale features. This is both indicated by the analysis of variance on replicate samples and by the nugget effect of the calculated semivariance.

Given both small scale variability and between survey bias, a final regional overview will contain large uncertainties. Despite that large scale features appear visible, like the relatively higher concentrations in the south of Zuid-Beveland which are presumably caused by the presence of orchards, these should be judged with caution. Extreme caution is needed when the values are compared with (soil specific) legal limits. Due to the uncertainty caused by the various sources of variability, such comparison might have very limited value.

The degradation level of DDT residues indicates that the Zeeland soils will remain a source for $\sum DDx$ longer than expected. Since it is not assumed that new applications of DDT will occur, the monitoring of DDT residue levels could be cautionary. The most important issue which should be addressed is the reduction of variability in the monitoring data. Probably the largest reduction can be obtained in more standardisation of field survey procedures, reducing the between survey bias. The small scale variance may be assessed by increasing the number of subsamples and sampling on smaller scales but this might be less cost-effective than reducing the between survey bias. However, policy and regulation based on these data should always incorporate the uncertainties as a result of the remaining variability.




Figure 7.9. Relation between DDT, its isomers DDE and DDD, and the number of half-times. Circles denote the values of survey 3, diamonds denote survey 5.

Table 7.5. Selected \sum DDx values from different areas and studies. Format is (min:max)/statistic where statistic, noted in column 'stat' is median (m), average (a), or geometric mean (g). Detection limits are presented with italic font while non available data is denoted as NA. Type denotes type of area, agricultural (a), orchard (o), uninfluenced (u) i.e. no active DDT application, unknown (-). The column 'n' gives the number of samples. Column 'ref' refers to the references below the table.

type	DDT [mg/kg]	DDE [mg/kg]	DDD [mg/kg]	\sum DDx [mg/kg]	stat	n	ref
a				(0.001:1.1)/0.010	m	1	1
a	(0.010:0.160)/0.040	(0.010:0.094)/0.042	(0.010:0.080)/0.024		m	16	2
o	(0.01:17.07)/0.40	(0.01:3.11)/0.75	(0.01:2.55)/0.16	(0.03:28.7)/1.30	m	6	3
-	(0:0.005)/0				m	13	4
-		(0.005:0.0077)/0.0065			g	3	5
a	(0.002:0.0173)/0.002 (o, p-DDT)				g	19	5
a	(0.002:0.367)/0.0124 (p, p-DDT)				g	19	5
u	(NA:NA)/0.060	(NA:NA)/0.060					6
a	(0.005·10 ⁻³ :0.411·10 ⁻³)/0.047·10 ⁻³	(0.005·10 ⁻³ :6.217·10 ⁻³)/0.092·10 ⁻³	(0.005·10 ⁻³ :0.031·10 ⁻³)/0.005·10 ⁻³		m	13	7
a	(0.005·10 ⁻³ :0.972·10 ⁻³)/0.029·10 ⁻³	(0.005·10 ⁻³ :19.9·10 ⁻³)/0.005·10 ⁻³	(0.005·10 ⁻³ :0.313·10 ⁻³)/0.005·10 ⁻³		m	16	7
u	(0.0005:0.012)/0.0055				a	9	8
a				(0.0035:0.1195)/0.0355	a	11	9
a				(0.007:0.9722)/0.056	a	180	10
-				(0.006:0.0224)/0.0068	a	16	11
u	(0.00006-0.0006)/0.00021	(0.00008-0.00283)/0.0006		(0.0006:0.003)/0.001	m	8	12
a				(0.0002:0.0897)/0.012	g	15	13

References: 1) This study, 2) van Gaans et al. (1995), 3) Gaw et al. (2003), 4) Feng et al. (2003), 5) Harner et al. (1999), 6) Miglioranza et al. (1999), 7) Cavanagh et al. (1999), 8) Rovinsky et al. (1995), 9) Covaci et al. (2001), 10) Gong et al. (2004), 11) Covaci et al. (2002), 12) Fu et al. (2001), 13) McGrath (1995)

8 Synthesis

The aim of this thesis is to assess how patterns in the geochemical soil composition can be used to distinguish natural variability from anthropogenic impact both in attribute and geographical space. The patterns found in this study are related to processes determining variability in soil composition, including variability caused by human influence. The many soil quality indicators presented can be used as a first initiative for sustainable management of soil quality.

In the following I will evaluate the methods used, the achievements of this study, and give an overview of the implications of the results, together with possible future applications.

8.1 Methods used

The methods used in this study are in general based on state of the art analytical methods combined with well established statistical techniques from the field of geochemical surveying and exploration. With the use of X-ray fluorescence spectrometry (XRF) and inductively coupled plasma mass spectrometry (ICP-MS) a broad range of elements could be analysed on many samples. For future comparison with data obtained from regular environmental Dutch soil surveys, the ICP-MS samples were leached using a hot aqua-regia solution. Both multi-element techniques were specifically chosen to broaden the analytical focus with major components, trace elements, and the rare earth elements (REE). This wide range of components can be obtained with minimal extra effort compared to a smaller selection such as environmental priority metals. This resulted in a better understanding of not only the human effect but also of the natural composition and relations in the soil.

By using both techniques some overlap between measured components occurred, which was useful to obtain extra information about analytical quality and bias. Due to the different approaches of the techniques, partial aqua-regia leach for ICP-MS and total analysis for XRF, some basic mineralogical information was also obtained.

Using well established statistical techniques for data analysis has the advantage that possibilities and limitations are well known and that many examples are available. This makes the techniques more transparent to the researcher and easily explainable to users. However, more complex methods have not been avoided, such as robust principal components analysis, so long as they provide a true advantage over standard techniques.

The methodology of this study incorporated several techniques to obtain information about the accuracy, variability and bias on analytical and sampling scales and, if applicable, methods to reduce noise and bias. It is essential to be able to distinguish the contribution of geochemical soil variability from that due to chemical analysis and sampling.

8.2 Achievements of this study

It was expected from the history of Zeeland that the soil contained slightly elevated concentrations of so called “heavy metals” (Cd, Cu, Pb, and Zn) compared to pristine background concentration levels. Also persistent organochlorine pesticide residues were expected to be present. This is confirmed in this study.

It was shown that the spatial variability of the heavy metals, Fe, and calcite related components, has a relatively large contribution of local scale variance. This means that for these components small scale variability is an important source of the total variation and compositing for these elements is considered useful. For most other elements the regional variance predominates, reducing the need for compositing.

The main part of this study comprises the geochemical characterisation of the province. The main hypotheses were that 1) based on insight into the natural variability the anthropogenic imprint could be recognised and 2) the subsoil is suitable for obtaining a geochemical baseline as a reference to calculate the anthropogenic impact. An additional question studied was if the imprint results in spatially distinct areas where specific human processes have had a relevant contribution to the soil composition.

To test these hypotheses a dataset of 31 chemical components was selected from the overall dataset of 83 components, based on precision, accuracy and completeness. For this dataset a measurement was available for each variable at each location and for each depth, topsoil and subsoil. The dataset still comprised major, minor, and trace elements, providing a reference for a wide range of components. It formed the core for the geochemical assessment of the soils of Zeeland.

To determine the anthropogenic imprint it is necessary to have a reference, or baseline, comprising the natural soil composition. For the Netherlands it was believed that no good baseline existed (van der Meent et al., 1990), and reference values like the legal threshold values are not considered representative for natural soil composition. When a true baseline is available, deviations from this baseline can be attributed to separate processes. Human processes, for example the local application of fertilisers or atmospheric deposition of contaminants from remote areas, will in general lead to increased concentration levels, or enrichment, of certain components. Using a baseline, such enrichment can be determined and the extent of the influence of the associated process can be assessed.

A good method for many elements to obtain a baseline is the use of the relation with Al_2O_3 . The correlation of an element with Al_2O_3 can indicate if the element is uninfluenced by human processes. Overall higher correlations with Al_2O_3 in the subsoil in Zeeland confirm the hypothesis that the subsoil can be used as a pristine, non-influenced, reference. Using

a regression model based on the subsoil relation of an element with Al_2O_3 , the extent of the enrichment of the topsoil can be quantified both in absolute terms and as ratio with the subsoil baseline. The enrichment can be assessed using individual enrichments of soil components or by the average enrichment, called the contamination index, of all components with elevated concentration. Despite the presence of enrichments for many elements the legal threshold level is not exceeded, thus the soils may be regarded as legally clean with regard to inorganic components. This implies also that the legal threshold values are not a suitable reference for natural background soil concentrations. Compared to all other techniques illustrated in chapter 5, the subsoil regression baseline method gives the best insight into 1) the geological/pedological pattern of geochemical variability, 2) random natural variation, and 3) the anthropogenic imprint on Zeeland soils.

The use of the above described baseline provides enrichments for individual elements based on the covariability with only one other variable. To fully understand the soil factors that determine the variability in soil composition a more multidimensional space must be considered. Multi-element covariability (principal component analysis) revealed four main factors that can be related to clay, calcite, silt and anthropogenic processes. The natural clay and calcite related processes comprise the largest part of the overall soil variability. The anthropogenic factor explains only a small part of the overall variability emphasising that these anthropogenic processes are really an imprint on the overall natural variability. Yet, for individual elements such as Sn, the anthropogenic factor may explain up to about 40% of the variance.

The regional patterns of both the contamination index and the scores on the anthropogenic factor show many similarities. However, since the anthropogenic factor explains the variability caused by human processes based on the total soil composition while the contamination index is an average of ratios from a few bivariate relations, I prefer the former. Nevertheless, enrichments or the contamination index can be quantified as concentrations of individual components or average of more components, making it easier to compare them with normative values and values from other areas.

When the samples in attribute space are grouped, i.e. based on their similarity with respect to interactions between the processes distinguished, no spatial regions emerge where samples are discriminated solely on the basis of anthropogenic enrichment. This suggests that the extent of the human imprint on the Zeeland soil, is not related to certain regions but rather is distributed diffusely over the province.

Using the techniques and experience from the former chapters, chapter 7 gave an assessment of DDT residue levels based on separate datasets provided by the local authorities. Compared with other areas, the DDT residue concentrations in Zeeland are such that there is no reason to believe that the DDT contamination is more severe than can be expected of an agricultural area. The extent of the values compared to uninfluenced areas is about a factor 2 to 10 higher, indicating that values for Zeeland must be regarded as enriched compared to the non-natural ubiquitous DDT residue background.

Unfortunately, a regional assessment of DDT residue values is hampered by the large contribution of small scale (sampling) variance. This means that for a regional overview the

observed variance is largely influenced by local scale noise. Together with the clear differences between the sub datasets, this implies that the regional variability should be judged with caution, despite the fact that large scale features are still visible.

8.3 Implications

The collection of soil quality data over time is a prerequisite of sustainable management. With a dynamic assessment the direction and magnitude of change can be evaluated (Wienhold et al., 2004). This study showed that with the use of state of the art analytical techniques and well established methods from geochemistry, a profound insight can be obtained in a single survey into the natural soil composition and how it is influenced by human processes. This both provides methods to obtain a reference for natural soil composition and to quantify the anthropogenic imprint. It is shown in this study that the framework of legal limits does neither provides such a reference nor is able to detect subtle anthropogenic enrichment. This, in a way, is remarkable considering that this framework is also used for the building material decree and related to soil pollution risk maps that should prevent spreading of contaminants. I therefore suggest that references for soil composition should be derived from (local) baselines rather than legal limits. When obtaining such a baseline, quantification of the variability due to sampling and analytical procedures, and methods to reduce or compensate bias between sub-surveys are as important as the baseline itself. If multi-element techniques are used, as most laboratories do, focus should not only be on environmental priority components but also on components which might provide information about natural geochemical soil composition and variability.

Soil pollution risks appear to be, as defined in the Dutch normative framework, mainly determined by DDT residues. Unfortunately, the on average low concentrations in soil, complex analytical determination, and high small-scale variability result in large uncertainties concerning the regional variability of this component. Consequently, to obtain a more trustworthy georeferenced overview of DDT residue levels, more effort aimed at reduction of variance and bias should be put in the assessment of this component.

The question remains what “soil quality” should mean for the Dutch situation and which indicators to use. The definition of soil quality as given by Karlen et al. (1997): “the capacity of a specific kind of soil to function, within natural or managed boundaries, to sustain plant and animal productivity, maintaining or enhance water and air quality, and support human health and habitation”, clearly implies not only a characterisation, but an evaluation against certain criteria as well. The so called “soil quality maps”, as used in the Netherlands to facilitate spatial development, clearly do not provide information about soil quality as envisaged by Karlen et al. (1997), despite their name. Therefore, one of the goals to reach sustainable soil management should be a reconsideration of what is understood by “soil quality”. Only then it is possible to define which soil quality indicators should be assessed. The reference data and methods provided in this study could be a starting point for obtaining such indicators, finally leading towards sustainable soil management.

Bibliography

- ABRAHAM, P. W. (2002), Soils: their implications to human health. *Science of the Total Environment* 291, pp. 1–32.
- ALLEGRE, C. J. & E. LEWIN (1995), Scaling Laws and geochemical distributions. *Earth and Planetary Science Letters* 132, pp. 1–12.
- ANONYMOUS (1999), Vrijstellingsregeling grondverzet. *Staatscourant* 180.
- BAUER, I. & J. BOR (1995), Lithogene, geogene und anthropogene Schwermetallgehalte von Lössboden an den Beispielen von Kupfer, Zink, Nickel, Blei, Quecksilber und Cadmium. *Mainzer Geowiss. Mitt.* 24, pp. 47–70.
- BAUER, I., V. M. SPRENGER & J. BOR (1992), Die Berechnung lithogener und geogener Schwermetallgehalte von Lössboden am Beispiel von Kupfer, Zink und Blei. *Mainzer Geowiss. Mitt.* 21, pp. 7–34.
- BAZEN, M. (1987), Bodemkaart van Nederland 1:50.000. Toelichting bij de kaartbladen 48 Oost Middelburg en 49 West Bergen op Zoom. Stichting bodemkartering, Wageningen.
- BEZDEK, J. C., R. EHRLICH & W. E. FULL (1984), FCM; The fuzzy c-means clustering algorithm. *Computer & Geosciences* 10, pp. 191–203.
- BIANCHINI, G., R. LAVIANO, S. LOVO & C. VACCARO (2002), Chemical-mineralogical characterisation of clay sediments around Ferrara (Italy): a tool for an environmental analysis. *Applied Clay Science* 21, pp. 165–176.
- BLASER, P., S. ZIMMERMANN, J. LUSTER & W. SHOTYK (2000), Critical examination of trace element enrichments and depletions in soils: As, Cr, Cu, Ni, Pb, and Zn in Swiss forest soils. *Science of the Total Environment* 249, pp. 257–280.
- BÖLVIKEN, B., J. BOGEN, A. DEMETRIADES, W. DE VOS, J. EBBING, R. HINDEL, M. LANGEDAL, J. LOCUTERA, P. O'CONNOR, R. T. OTTESEN, E. PULKKINEN, R. SALMINEN, O. SCHERMANN, R. SWENNEN, J. VAN DER SLUYS & T. VOLDEN (1996), Regional geochemical mapping of Western Europe towards the year 2000. *Journal of Geochemical Exploration* 56, pp. 141–166.
- BOUNESSAH, M. & B. P. ATKIN (2003), An application of exploratory analysis (EDA) as a robust non-parametric technique for geochemical mapping in a semi-arid climate. *Applied Geochemistry* 18, pp. 1185–1195.
- BRAUNE, B., D. MUIR, B. DEMARCH, M. GAMBERG, K. POOLE, R. CURRIE, M. DODD, W. DUSCHENKO, J. EAMER, B. ELKIN, M. EVANS, S. GRUNDY, C. HEBERT, R. JOHNSTONE, K. KIDD, B. KOENIG, L. LOCKHART, H. MARSHALL, K. REIMER, J. ANDERSON & L. SHUTT (1999), Spatial and temporal trends of contaminants in the Canadian Arctic freshwater and terrestrial ecosystems: a review. *Science of the Total Environment*, pp. 145–207.
- BRIDGES, E. M. & J. H. V. VAN BAREN (1997), Soil: an overlooked, undervalued and vital part of the human environment. *The Environmentalist* 17, pp. 15–20.

- BURENKOV, E. K., A. A. GOLOVIN, I. MOROZOVA & E. I. FILATOV (1999), Multi-purpose geochemical mapping (1:1,000,000) as a basis for the integrated assessment of natural resources and ecological problems. *Journal of Geochemical Exploration* 66, pp. 159–172.
- BURROUGH, P. A., P. F. M. VAN GAANS & R. HOOTSMANS (1997), Continuous classifications in soil survey: spatial correlation, confusion and boundaries. *Geoderma* 77, pp. 115–135.
- BUSINK, E. R. V. & S. POSTMA (2000), Provincial soil-quality monitoring networks in the Netherlands as an instrument for environmental protection. *Netherlands Journal of Geosciences* 4, pp. 429–440.
- CARSON, R. (1962), *Silent Spring*. Houghton Mifflin Company, Boston.
- CAVANAGH, J. E., K. A. BURNS, G. J. BRUNSKILL & R. J. COVENTRY (1999), Organochlorine pesticide residues in soils and sediments of the Herbert and Burdekin River regions, North Queensland: implications for contamination of the Great Barrier Reef. *Marine pollution Bulletin* 39, pp. 367–375.
- CHORK, C. Y. & R. SALMINEN (1993), Interpreting exploration geochemical data from Outokumpu, Finland: A MVE-robust factor analysis. *Journal of Geochemical Exploration* 48, pp. 1–20.
- COCKER, M. D. (1999), Geochemical mapping in Georgia, USA: a tool for environmental studies, geologic mapping and mineral exploration. *Journal of Geochemical Exploration* 67, pp. 345–360.
- COVACI, A., C. HURA & P. SCHEPENS (2001), Selected persistent organochlorine pollutants in Romania. *Science of the Total Environment* 280, pp. 143–152.
- COVACI, A., P. MANIRAKIZA & P. SCHEPENS (2002), Persistent organochlorine pollutants in soils from Belgium, Italy, Greece, and Romania. *Environmental Contamination and Toxicology* 68, pp. 97–103.
- DANESHFAR, B. & E. CAMERON (1998), Leveling geochemical data between map sheets. *Journal of Geochemical Exploration* 63, pp. 189–201.
- DARNLEY, A., A. BJÖRKLAND, B. BØLVIKEN, N. GUSTAVSSON, P. V. KOVAL, J. A. PLANT, A. STEENFELT, M. TAUCHID & X. XUEJING (1995), A global geochemical database for environmental and resource management. Final Report of IGCP Project 259, Unesco Publishing.
- DARNLEY, A. G. (1995), International geochemical mapping, a review. *Journal of Geochemical Exploration* 55, pp. 5–10.
- DAVIS, J. C. (2002), *Statistics and Data Analysis in Geology*. John Wiley and Sons, New York.
- DE GRUIJTER, J. J. & C. J. F. TER BRAAK (1990), Model-Free Estimation from Spatial Samples: A Reappraisal of Classical Sampling Theory. *Mathematical Geology* 22, pp. 407–415.
- DE LÓPEZ CAMELO, L. G., S. R. DE MIGUEZ & L. MARBÁN (1997), Heavy metals input with phosphate fertilizers used in Argentina. *Science of the Total Environment* 204, pp. 245–250.
- DE ROO, G. (2003), *Environmental planning in the Netherlands: Too good to be true*. Ashgate, Hampshire.
- DE VOS, W., J. EBBING, R. HINDEL, J. SCHALICH, R. SWENNEN & I. V. KEER (1996), Geochemical mapping based on overbank sediments in the heavily industrialised border area of Belgium, Germany and the Netherlands. *Journal of Geochemical Exploration* 56, pp. 91–104.
- DIMOND, J. B. & R. B. OWEN (1996), Long-term residue of DDT compounds in forest soils in Maine. *Environmental Pollution* 92, pp. 227–230.
- DORAN, J. W. (2002), Soil health and global sustainability: translating science into practice. *Agriculture, Ecosystems and Environment* 88, pp. 119–127.
- EDELMAN, T. (1984), *Achtergrondgehalten van stoffen in de bodem*. Staatsuitgeverij, Den Haag.
- EDWARDS, C. A. (1970), *Persistent pesticides in the environment*. CRC monoscience series, Butterworths, London.

- EIKELBOOM, R. T., E. RUWIEL & J. J. J. M. GOUMANS (2001), The Building Material Decree: an example of a Dutch regulation based on the potential impact of materials on the environment. *Waste Management* 21, pp. 295–302.
- FACCHINELLI, A., E. SACCHI & L. MALLEN (2001), Multivariate statistical and GIS-based approach to identify heavy metal sources in soils. *Environmental Pollution* 114, pp. 313–324.
- FAO (1990), FAO-UNESCO soil map of the world, revised legend. *Soils Bulletin* 60, FAO, Rome.
- FENG, K., B. Y. YU, D. M. GE, M. H. WONG, X. C. WANG & Z. H. CAO (2003), Organochlorine pesticide (DDT and HCH) residues in the Taihu Lake Region and its movement in soil-water system; I. Field survey of DDT and HCH residues in ecosystems of the region. *Chemosphere* 50, pp. 683–687.
- FERREIRA, A., M. M. INÁCIO, P. MORGADO, M. J. BATISTA, L. FERREIRA, V. PEREIRA & M. S. PINTO (2001), Low-density geochemical mapping in Portugal. *Applied Geochemistry* 16, pp. 1323–1331.
- FOLKES, D. J., T. E. KEUHSTER & R. A. LITTLE (2001), Contributions of pesticide use to urban background concentrations of arsenic in Denver, Colorado, U.S.A. *Environmental Forensics* 2, pp. 127–139.
- FU, S., S. CHU & S. XU (2001), Organochlorine pesticide residue in soils from Tibet. *Bulletin of Environmental Contamination and Toxicology* 2001, pp. 171–177.
- GARRETT, R. G. (1983), Sampling Methodology. In: R. J. Howarth, ed., *Handbook of exploration geochemistry*. Volume 2: statistics and data analysis in geochemical prospecting, Elsevier Science.
- GARRETT, R. G. & T. I. GOSS (1979), The Evaluation of Sampling and Analytical variation in Regional Geochemical Surveys. In: J. R. Watterson & P. K. Theobald, eds., *Geochemical Exploration 1978*, Association of Exploration Geochemists, Rexdale.
- GARRETT, R. G. & T. I. GOSS (1980), Uanova: A FORTRAN IV program for unbalanced nested analysis of variance. *Computer & Geosciences* 6, pp. 35–60.
- GAW, S. K., G. PALMER, N. D. KIM & A. L. WILKINS (2003), Preliminary evidence that copper inhibits the degradation of DDT to DDE in pip and stonefruit orchard soils in the Auckland region, New Zealand. *Environmental pollution* 122, pp. 1–5.
- GIMENO-GARCÍA, E., V. ANDRUE & R. BOLUDA (1996), Heavy metals incidence in the application of inorganic fertilizers and pesticides to rice farming soils. *Environmental Pollution* 92, pp. 19–25.
- GOEMANS, T. & T. VISSER (1987), The Delta Project; The Netherlands experience with a megaproject for flood protection. *Technology in Society* 9, pp. 97–111.
- GONG, Z. M., S. TAO, F. L. XU, R. DAWSON, W. X. LIU, Y. H. CUI, J. CAO, X. J. WANG, W. R. SHEN, W. J. ZHANG, B. P. QING & R. SUN (2004), Level and distribution of DDT in surface soils from Tianjin, China. *Chemosphere* 54, pp. 1247–1253.
- GREGORAUSKIENE, V. & V. KADUNAS (1997), Experience and goals of geochemical mapping for environmental protection in Lithuania. *Journal of Geochemical Exploration* 60, pp. 67–76.
- GROOT, M. S. M., J. J. B. BRONSWIJK & T. V. VAN LEEUWEN (2001), Landelijk meetnet bodemkwaliteit, resultaten 1996. rapport nr. 714801026, Rijksinstituut voor Volksgezondheid en Milieu.
- GUSTAVSON, N., E. LAMPPIO & T. TARVAINEN (1997), Visualisation of geochemical data on maps at the Geological Survey of Finland. *Journal of Geochemical Exploration* 59, pp. 197–207.
- HALE, M. & R. KÜHNEL (1997), Exploration of contaminated soils and sediments. *Applied Clay Science* 11, pp. 373–380.
- HALFWERK, W. (1996), *Natuur in Zeeland, tussen Zee en Zoom* (Nature in Zeeland, between sea

- and zoom). De Koperen Tuin, Goes.
- HANESCH, M., R. SCHOLGER & M. J. DEKKERS (2001), The application of fuzzy c-means cluster analysis and non-linear mapping to a soil data set for the detection of polluted sites. *Physical Chemical Earth* 26, pp. 885–891.
- HARNER, T., J. L. WIDEMAN, L. M. M. JANTUNEN, T. F. BIDLEMAN & W. J. PARKHURST (1999), Residues of organochlorine pesticides in Alabama soils. *Environmental Pollution* 106, pp. 323,332.
- HEIJBOER, D. & J. NELLESTIJN (2002), *Klimaatatlas van Nederland: de normaalperiode 1971-2000*. Elmar, Rijswijk.
- HOWARD, P. H. (1991), *Environmental degradation rates*. Lewis Publishers, Michigan.
- HUISMAN, D. J., F. J. H. VERMEULEN, J. BAKER, A. VELDKAMP, S. B. KROONENBERG & G. T. KLAVER (1997), A geological interpretation of heavy metal concentrations in soils and sediments in the southern Netherlands. *Journal of Geochemical Exploration* 59, pp. 163–174.
- JOURNAL, A. G. & C. J. HUIJBREGTS (1981), *Mining Geostatistics*. Academic Press, London.
- KARLEN, D. L., M. J. MAUSBACH, J. W. DORAN, R. G. CLINE, R. F. HARRIS & G. E. SCHUMAN (1997), Soil quality: a concept, definition, and framework for evaluation. *Journal Soil Science Society of America* 61, pp. 4–10.
- KHATREE, R., D. N. NAIK & R. L. MASON (1997), Estimation of variance components in staggered nested designs. *Journal of Applied Statistics* 24, pp. 395–408.
- KIM, J.-H. & A. SMITH (2001), Distribution of organochlorine pesticides in soils from South Korea. *Chemosphere* 43, pp. 137–140.
- KOELEMAN, M., W. J. VAN DER LAAK & H. IETSWAART (1999), Dispersion of PAH and heavy metals along motorways in the Netherlands: an overview. *Science of the Total Environment* 235, pp. 347–349.
- KRAMAR, U. (1995), Application of limited fuzzy clusters to anomaly recognition in complex geological environments. *Journal of Geochemical Exploration* 55, pp. 81–92.
- LEENAERS, H., J. DE BOER, D. BOELS., M. HOOGERWERF & A. WEENK (1999), *Bodemkwaliteitsbeeld 2005: de lokale en regionale informatiebehoefte*. Tech. Rep.
- LICHT, O. A. B. & T. TARVAINEN (1996), Multipurpose geochemical maps produced by integration of geochemical exploration data sets in The Paraná Shield, Brazil. *Journal of Geochemical Exploration* 56, pp. 167–182.
- MATSCHULLAT, J., R. OTTENSTEIN & C. REIMANN (2000), Geochemical background - can we calculate it? *Environmental Geology* 39, pp. 990–1000.
- MCBRATNEY, A. B. & I. O. A. ODEH (1997), Application of fuzzy sets in soils science: fuzzy logic, fuzzy measurements and fuzzy decisions. *Geoderma* 77, pp. 85–113.
- MCGRATH, D. (1995), Organic micropollutant and trace element pollution of Irish soils. *Science of the Total Environment* 164, pp. 125–133.
- MCGRATH, D. & C. ZHANG (2003), Spatial distribution of soil organic carbon concentrations in grassland of Ireland. *Applied Geochemistry* 18, pp. 1629–1639.
- MEGHARAJ, M., H. L. BOUL & J. H. THIELE (1999), Effects of DDT and its metabolites on soil algae and enzymatic activity. *Biol Fertil Soils* 29, pp. 130–134.
- MIESCH, A. T. (1975), Variograms and variance components in geochemistry and ore evaluation. In: E. H. T. Whitten, ed., *Quantative studies in the geological sciences - A memoir in honor of William C. Krumbein*, vol. 142 of *Geological Society of America*.
- MIESCH, A. T. (1976), *Geochemical survey of Missouri: methods of sampling, laboratory analysis and statistical reduction of data*. U.S. Geol. survey prof. paper .
- MIGLIORANZA, K. S. B., J. E. A. DE MORENO, V. J. MORENO, M. L. OSTERRIETH & A. H.

- ESCALANTE (1999), Fate of organochlorine pesticides in soils and terrestrial biota of Los Padres pond watershed, Argentina. *Environmental Pollution* 105, pp. 91–99.
- MOL, G. (2002), Soil acidification monitoring in the Netherlands. Ph.D. thesis, Faculty of Earth Sciences, Utrecht University.
- MOL, G., S. P. VRIEND & P. F. M. VAN GAANS (2001), Environmental monitoring in the Netherlands: past developments and future challenges. *Environmental Monitoring and Assessment* 68, pp. 313–335.
- MOSIER, A. R. (1998), Soil processes and global change. *Biol. Fertil. Soils* 27, pp. 221–229.
- NASH, D., D. HALLIWELL, M. HANNAH, L. CLEMOW & B. WEBB (2003), Phosphorus and selected metals mobilized from two commercial fertilizers into overland flow during border irrigation. *Nutrient Cycling in Agroecosystems* 67, pp. 255–264.
- NIEROP, T. M. (2000), Samen slim met water; Waterhuishoudplan 2001-2007. Tech. Rep., Provincie Zeeland.
- NIKKARINEN, M., E. KAUNISKANGAS & K. NENONEN (1996), Use of geological and geochemical data for the derivation of maps for environmental planning in Iisalmi, Finland. *Applied Geochemistry* 11, pp. 261–270.
- NNI (1999a), NEN 5734: Bepaling van het gehalte PCB's en OCB's in grond. Tech. Rep., Netherlands Normalisatie Instituut (NNI).
- NNI (1999b), NEN 5740: Bodem-onderzoeksstrategie bij verkennend onderzoek. Tech. Rep., Netherlands Normalisatie Instituut (NNI).
- NORTCLIFF, S. (1978), Soil Variability and Reconnaissance Soil Mapping: A Statistical Study in Norfolk. *Journal of Soil science* 29, pp. 403–418.
- NORTCLIFF, S. (2002), Standardisation of soil quality attributes. *Agriculture, Ecosystems and Environment* 88, pp. 161–168.
- OLIVER, M. A. & A. L. KHAYRAT (2001), A geostatistical investigation of the spatial variation of radon in soil. *Computer & Geosciences* 27, pp. 939–957.
- OLIVER, M. A. & R. WEBSTER (1986), Combining nested and linear sampling for determining the scale and form of spatial variation of regionalized variables. *Geographical Analysis* 18, pp. 227–242.
- ORGANISATION, W. H. (1989), Environmental Health Criteria 83. DDT and its derivatives- Environmental aspects. Tech. Rep.
- PARKPIAN, P., S. T. LEONG, P. LAORTANAKUL & N. THUNTHAISONG (2003), Regional monitoring of lead and cadmium contamination in a tropical grazing land site, Thailand. *Environmental Monitoring and Assessment* 85, pp. 157–173.
- PEBESMA, E. J. & C. G. WESSELING (1998), Gstat: A program for geostatistical modelling, prediction and simulation. *Computer & Geosciences* 24, pp. 17–31.
- PETTITT, A. N. & A. B. MCBRATNEY (1993), Sampling Designs for Estimating Spatial Variance Components. *Applied Statistics* 42, pp. 185–209.
- PISON, G., P. J. ROUSSEUW, P. FILZMOSEER & C. CROUX (2003), Robust factor analyses. *Journal of Multivariate Analysis* 84, pp. 145–172.
- PLEIJTER, G. & C. WALLENBURG (1994), Bodemkaart van Nederland 1:50.000. Toelichting bij de kaartbladen 42 West en 42 Oost Zierikzee, 47/48 West Cadzand/Middelburg. DLO-Staringcentrum, Wageningen.
- POLITOV, S. V., Y. A. ANOKHIN & N. N. SURNINA (2000), Organochlorine pesticides and polychlorinated biphenyls in the environment of the Baikal region. *Aquatic Ecosystems Health and Management* 3, pp. 235–238.
- R DEVELOPMENT CORE TEAM (2004), R: A language and environment for statistical com-

- puting. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-00-3, URL <http://www.R-project.org>.
- RAPANT, S., M. RAPOSOVÁ, D. BODIS, K. MARSINA & I. SLANINKA (1999), Environmental-geochemical mapping program in the Slovak Republic. *Journal of Geochemical Exploration* 66, pp. 151–158.
- REIMANN, C. & P. FILZMOSER (1999), Normal and lognormal data distribution in geochemistry: death of a myth. Consequences for the statistical treatment of geochemical and environmental data. *Environmental Geology* 39, pp. 1001–1014.
- REIMANN, C., P. FILZMOSER & R. G. GARRETT (2002), Factor analysis applied to regional geochemical data: problems and possibilities. *Applied Geochemistry* 17, pp. 185–206.
- RIDGWAY, J., N. BREWARD, W. J. LANGSTON, R. LISTER, J. G. REES & S. M. ROWLATT (2003), Distinguishing between natural and anthropogenic sources of metals entering the Irish Sea. *Applied Geochemistry* 18, pp. 283–309.
- RIJKSWATERSTAAT (1971), Beschrijving van de provincie Zeeland behorende bij de waterstaatskaart. Tech. Rep., Directie Algemene Dienst van de Rijkswaterstaat.
- RIVM (2004), Milieu- en Natuurcompendium. Tech. Rep., RIVM-MNP, Bilthoven; CBS, Den Haag; Stichting DLO, Wageningen. URL <http://www.milieuennatuurcompendium.nl>.
- ROSE, A. W., H. E. HAWKES & J. S. WEBB (1979), *Geochemistry in mineral exploration*. Academic Press, London.
- ROUSSEEUW, P. J. & K. VAN DRIESSEN (1999), A fast algorithm for the minimum covariance determinant estimator. *Technometrics* 41, pp. 212–223.
- ROVINSKY, F., B. PASTUKHOV, Y. BOUYVOLOV & L. BURTSEVA (1995), Present day state of background pollution of the natural environment in the Russian Arctic in the region of the UST-Lena Reserve. *Science of the Total Environment* 160/161, pp. 193–199.
- RUTTEN, K. (2004), een vergelijking van de geochemie en bodemkunde van Zeeland. Master's thesis, Utrecht University, department of geosciences.
- SALMINEN, R. & V. GREGORAUSKIENĖ (2000), Consideration regarding the definition of a geochemical baseline of elements in the surficial materials in areas differing in basic geology. *Applied Geochemistry* 15, pp. 647–653.
- SALMINEN, R. & T. TARVAINEN (1995), Geochemical mapping and databases in Finland. *Journal of Geochemical Exploration* 55, pp. 321–27.
- SALMINEN, R. & T. TARVAINEN (1997), The problem of defining geochemical baselines. A case study of selected elements and geological materials in Finland. *Journal of geochemical exploration* 60, pp. 91–98.
- SCANCAR, J., R. MILACIC & M. HORVAT (2000), Comparison of various digestion and extraction procedures in analysis of heavy metals in sediments. *Water, Air and Soil Pollution* 118, pp. 87–99.
- SEARLE, S. R., G. CASELLA & C. E. MCCULLOCH (1992), *Variance Components*. John Wiley & Sons.
- SELINUS, O. S. & K. ESBENSEN (1995), Separating anthropogenic from natural anomalies in environmental geochemistry. *Journal of Geochemical Exploration* 55, pp. 55–66.
- SENESI, G. S., G. BALDASSARRE, N. SENESI & B. RADINA (1999), Trace elements inputs into soils by anthropogenic activities and implications for human health. *Chemosphere* 39, pp. 343–377.
- SHARMA, V. K., K. B. RHUDY, J. C. CARGILL, M. E. TACKER & F. G. VAZQUEZ (2000), Metals and grain size distributions in soil of the middle Rio Grande basin, Texas USA. *Environmental Geology* 39, pp. 698–704.

- SHOTYK, W., D. WEISS, J. D. KRAMERS, R. FREI, A. K. CHEBURKIN, M. GLOOR & S. REESE (2001), Geochemistry of the peat bog at Etang de la Gruère, Jura Mountains, Switzerland, and its record of atmospheric Pb and lithogenic trace metals (Sc, Ti, Y, Zr, and REE) since 12,370 14C yr BP. *Geochimica et Cosmochimica Acta* 65, pp. 2337–2360.
- SINCLAIR, A. J. (1976), Applications of probability graphs in mineral exploration. Association of Exploration Geochemists.
- SINGH, D. K. & H. C. AGARWAL (1992), Chemical release and nature of soil-bound DDT-residues. *Journal of Agricultural Food Chemistry* 40, pp. 1713–1716.
- SINGH, M., G. MULLER & I. B. SING (2001), Geogenic distribution and baseline concentration of heavy metals in sediments of the Ganges River, India. *Journal of Geochemical Exploration* 80, pp. 1–17.
- SPOSITO, G. (1989), *The Chemistry of Soils*. Oxford University Press.
- STERCKEMAN, T., F. DOUAY, D. BAIZE, H. FOURRIER, N. PROIX & C. SCHVARTZ (2004), Factors affecting trace element concentrations in soils developed on recent maine deposits from Northern France. *Applied Geochemistry* 19, pp. 89–103.
- STIBOKA (1964), Bodemkaart van Nederland 1:50.000. Toelichting bij kaartblad 43 West Willemstad. Stichting voor Bodemkartering, Wageningen.
- STIBOKA (1967), Bodemkaart van Nederland 1:50.000. Toelichting bij kaartblad 53 Sluis en 54 West Terneuzen. Stichting voor Bodemkartering, Wageningen.
- STIBOKA (1980), Bodemkaart van Nederland 1:50.000. Toelichting bij kaartblad 54 Oost Terneuzen, 55 Hulst en het Zeeuws-Vlaamse deel van de kaartbladen 48 Oost Middelburg, 49 West Bergen op Zoom. Sluis en 54 West Terneuzen. Stichting voor Bodemkartering, Wageningen.
- SWARTJES, F. A. (1999), Risk-based assessment of soil and groundwater quality in the Netherlands: standards and remediation urgency. *Risk Analysis* 19, pp. 1235–1249.
- SWENNEN, R. & J. VAN DE SLUYS (2002), Anthropogenic impact on sediment composition and geochemistry in vertical overbank profiles of river alluvium from Belgium and Luxembourg. *Journal of Geochemical Exploration* 75, pp. 93–105.
- SZAVA-KOVATS, R. C. (2002), Outlier-resistant errors-in-variables regression: anomaly recognition and grain-size correction in stream sediments. *Applied Geochemistry* 17, pp. 1149–1157.
- TACK, F. M. G., M. G. VERLOO, L. VANMECHELEN & E. V. RANST (1997), Baseline concentration levels of trace elements as a function of clay and organic carbon contents in soils in Flanders (Belgium). *Science of the Total Environment* 201, pp. 113–123.
- TAO, S. (1998), Factor score mapping of soil trace element contents for the Shenzhen Area. *Water, Air and Soil Pollution* 102, pp. 415–425.
- TEBBENS, L. A., A. VELDKAMP & S. B. KROONENBERG (2000), Natural compositional variation of the river Meuse (Maas) suspended load: a 13 ka bulk geochemical record from the upper Kreftenheye and Betuwe Formations in Northern Limburg. *Geologie en Mijnbouw/Netherlands Journal of Geosciences* 79, pp. 391–409.
- TSOLAKIDOU, A., J. B. I GARRIGÓS & V. KILIKOGLU (2002), Assessment of dissolution techniques for the analysis of ceramic samples by plasma spectrometry. *Analytica Chimica Acta* 474, pp. 177–188.
- TUKEY, J. W. (1977), *Exploratory data analysis*. Addison-Wesley, Philipines.
- URBAT, M., M. J. DEKKERS & S. P. VRIEND (1997), Fuzzy c-means clustering as an aid in the interpretation of the natural remanent magnetization in various types of sediments. In: G. V. Pawlowsky, ed., *Proceedings of IAMG '97: Third annual conference of the International Association for Mathematical Geology*.

- VAN DEN BERG, G. A. & J. P. G. LOCH (2000), Decalcification of soils subject to periodic waterlogging. *Journal of Soil Science* 51, pp. 27–33.
- VAN DER GAAST, N., H. LEENAERS & J. ZEGWAARD (1998), The grey areas in soil pollution risk mapping. The distinction between cases of soil pollution and increased background levels. *Journal of Hazardous Materials* 61, pp. 249–255.
- VAN DER MEENT, D., T. ALDENBERG, J. H. CANTON, C. A. M. VAN GESTEL & W. SLOOFF (1990), *Desire for Levels*. rapport nr 670101002, Rijksinstituut voor Volksgezondheid en Milieu.
- VAN DER SLUIS, P., G. G. STEUR & I. OVAA (1965), *De bodem van Zeeland, Toelichting bij blad 7 bodemkaart van Nederland, schaal 1:200.000*. Stiboka, Wageningen.
- VAN DER WEIJDEN, C. H. (2002), Pitfalls of normalization of marine geochemical data using a common divisor. *Marine Geology* 184, pp. 167–187.
- VAN DIJK, D. & V. J. G. HOUBA (2000), Homogeneity and Stability of Materials Distributed Within the Wageningen Evaluating Programmes for Analytical Laboratories. *Communications in Soil Science and Plant Analysis* 31, pp. 1745–1756.
- VAN DRECHT, G., L. J. M. BOUMANS, D. FRATERS, H. F. R. REIJNDERS & W. VAN DUIJVENBODEN (1996), Landelijke beelden van diffuse metaalbelasting van de bodem en de metaalgehalten in de bovengrond, alsmede de relatie tussen gehalten en belasting. rapport nr. 714801006, Rijksinstituut voor volksgezondheid en milieu.
- VAN GAANS, P. F. M., S. P. VRIEND, S. BLEYERVELD, G. SCHRAGE & A. J. VOS (1995), Assessing Environmental Soil Quality in Rural Areas, a base line study in the Province of Zeeland, the Netherlands and reflections on soil monitoring network design. *Environmental Monitoring and Assessment* 34, pp. 73–102.
- VAN GAANS, P. M. & S. P. VRIEND (1996), PAK en de verspreiding van onderhoudsbaggerspecie over landbouwpercelen. Tech. Rep. ISBN 90-71577-96-1.
- VAN HELVOORT, P. J. (2003), Complex confining layers. A physical and geochemical characterization of heterogeneous unconsolidated fluvial deposits using a facies-based approach. Ph.D. thesis, Dutch geographical Studies.
- VAN LIENEN, F., G. FRAPPORTI & A. STEIN (2000), Construction of maps for soil recycling in regional infrastructural works integrating soil-quality laws. *Netherlands Journal of Geosciences* 4, pp. 449–457.
- VENABLES, W. N. & B. D. RIPLEY (2002), *Modern Applied Statistics with S*. Fourth Edition. Springer.
- VILLA, S., M. VIGHI, V. MAGGI, A. FINIZIO & E. BOLZACCHINI (2003), Historical trends of organochlorine pesticides in an alpine glacier. *Journal of Atmospheric Chemistry* 46, pp. 295–311.
- VOS, P. C. & R. M. VAN HEERINGEN (1997), Holocene geology and occupation history of the Province of Zeeland (SW Netherlands). In: M. M. Fischer, ed., *Holocene evolution of Zeeland (SW Netherlands)*, vol. 59 of *Medelingen Nederlands Instituut voor Toegepaste geowetenschappen*, Netherlands institute of applied geosciences TNO.
- VRIEND, S. P., P. F. M. VAN GAANS, J. MIDDELBURG & A. DE NIJS (1988), The application of fuzzy c-means cluster analysis and non-linear mapping to geochemical datasets: examples from Portugal. *Applied Geochemistry* 3, pp. 213–224.
- VROM (1994), circulaire streef en interventiewaarden. rapport nr. DBO/0749013, Ministerie van VROM.
- VROM (1997), Nationaal milieubeleidsplan 3. Tech. Rep., Ministry of Housing, Spatial Planning and the Environment, The Netherlands.
- VROM (1999), Interim Richtlijn. Opstellen en toepassen bodemkwaliteitskaarten in het kader van

- de Vrijstellingsregeling grondverzet. Beleidsstuk, Ministerie van VROM.
- WALSH, J. N., R. GILL & M. F. THIRLWALL (1997), Dissolution procedures for geological and environmental samples. In: *Modern analytical geochemistry, an introduction to quantitative chemical analysis for earth, environmental and materials scientists.*, Addison Wesley Longman Ltd, Harlow, UK, pp. 29–40.
- WEBSTER, R. (1985), Quantitative spatial analysis of soil in the field. In: Stewart, ed., *Advances in Soil Sciences*, vol. 3, Springer Verlag.
- WEBSTER, R. & M. A. OLIVER (1990), *Statistical methods in soils and land resource survey*. Oxford University Press.
- WEDEPOHL, K. H. (1995), The composition of the continental crust. *Geochemica Cosmochemica Acta* 59, pp. 1217–1232.
- WIENHOLD, B. J., S. ANDREWS & D. L. KARLEN (2004), Soil quality: a review of the science and experiences in the USA. *Environmental Geochemistry and Health* 26, pp. 89–95.
- WILCKE, W., S. MÜLLER, N. KANCHANAKOOL & W. ZECH (1998), Urban soil contamination in Bangkok: heavy metal and aluminium partitioning in topsoils. *Geoderma* 86, pp. 211–228.
- YOU DEN, W. J. & A. MEHLICH (1937), Selection of efficient methods for soil sampling. *Contr. Boyce Thompson Inst. Plant. Res.* 9, pp. 59–70.
- ZAGWIJN, W. H. (1991), *Nederland in het Holoceen*. Staatsuitgeverij, Den Haag.
- ZITKO, V. (2003), Chlorinated pesticides: Aldrin, DDT, Endrin, Dieldrin, Mirex. In: H. Fiedler, ed., *Persistent Organic Pollutants*, Springer-Verlag, Berlin Heidelberg.

Abstract - Geochemical patterns in the soils of Zeeland, natural variability versus anthropogenic impact

Chapter 1. Introduction

The close relation between soil, health, and global sustainability confirm that soil plays a critical role as a major interface in our environment and that soil quality can be an important indicator for sustainable environmental management.

This thesis is concerned with obtaining an overview of general (geo)chemical soil quality, within the framework of diffuse anthropogenic pollution and sustainable soil management in the Netherlands. It aims at assessing patterns in the geochemical soil composition and distinguishing the natural variability from the anthropogenic impact. The variability is assessed in geographical space, in which the spatial interaction of soil components takes place, and in attribute space, the space where interaction between soil constituents is exposed. Patterns from both spaces can then be related to processes influencing the soil composition.

Based on these patterns, and related processes, anthropogenic influence can be distinguished from natural variability. This can provide tools and information which can be part of the country wide overview needed as required by the Dutch government in their national environmental policy plan (NEPP 3).

The Province of Zeeland in the south west of the Netherlands was the area chosen for this research. The large rural area of young Holocene deposits and rich human history make it a suitable area for testing the main hypothesis that human influence leaves a distinct and quantifiable pattern of soil pollutants, based on variability within the geographical and attribute space of the soil composition.

Chapter 2. Geology and pedology of Zeeland

Zeeland is located in the marine delta of the rivers Rijn, Maas en Schelde. It was the sea level rise following the end of the Weichselian ice age that brought Zeeland within the

marine realm. The soils of Zeeland can be regarded as relatively homogeneous and young. They were moulded in a continuous interaction of natural processes and human endeavour. Considering the fact that soils of Zeeland are in general developed in marine clay deposits, it can be expected that a main source of geochemical variability is the varying clay content. The natural pattern of creek ridges and pool grounds creates relatively abrupt transitions. As local homogeneity or gradual variation may be further disturbed by extensive human activity, variability in soil composition is expected to be still higher than is directly evident from the soil map. Each (embanked) area with its own history, both sedimentary and human, may have its own pattern of soil variability. Given the fact that 80% of terrestrial Zeeland is used for agriculture, nearly 70% of which is arable land, human processes related to fertilisation and pesticide use are further expected to have influenced soil composition. This will result in elevated concentrations of so called “heavy metals” (Cd, Cu, Pb, and Zn) and persistent organochlorine pesticide residues. Finally, atmospheric inputs should also be considered as contributing to soil geochemistry.

Chapter 3. Natural and anthropogenic patterns of covariance and spatial variability of minor and trace elements in agricultural topsoil

The aim of this chapter was to analyze the sources of variation in soil geochemical composition related to spatial scale and sampling procedure. Apart from an intrinsic interest, research was motivated by practical questions regarding sampling strategies, using single or composite samples, for regional soil surveys. An unbalanced nested sampling design, based on single samples, was implemented at Walcheren/Zuid-Beveland. The samples were digested with aqua regia and analyzed by ICP-MS. The resulting data for 40 elements were examined through principal component analysis in combination with varimax rotation (PCA-V), unbalanced analysis of variance (UANOVA) and a fuzzy c-means clustering of the UANOVA results. The increase in noise when using single samples was further assessed through comparison with results of a previous study in the area based on composite samples. The three factors of the PCA-V are interpreted as being the results of natural, human, and geohydrological processes. The UANOVA analysis distinguishes three spatial variance patterns. Two groups show a gradual increase of variance with distance and contain respectively clay/feldspar related elements and heavy rare earth elements. The third group encompasses all anthropogenically and geohydrologically influenced elements with a larger part of the variance at small scales.

In general regional variance predominates over local variance and the extra effort of compositing is of limited value for a regional survey. However, environmental priority metals like Cd, Cu, Pb, Sb, Sn and Zn, were found to have larger local variance components. For these elements compositing is relatively more beneficial.

Chapter 4. Sampling and analyses for a regional environmental soil geochemical survey

When assessing soil composition for individual concentrations of elements it is often difficult to distinguish anthropogenic effects from natural variation. Establishing a bivariate or multivariate geochemical baseline, that incorporates a major part of natural variability, is an accepted method to derive background concentrations. Divergence from this background might then be caused by human disturbance. To quantitatively assess which components are affected by human activities, a consistent regional dataset is needed where special attention is paid to quantifying data quality. This chapter documents the construction of such a dataset for the province of Zeeland.

The sampling design was aimed at sampling present-day soil and pristine soil concentrations by sampling an influenced topsoil layer and an assumed pre-anthropogenic deeper subsoil layer at a density of 1 sample per 1-10 km². The analytical methods used (ICP-MS with aqua regia digestion and XRF on pressed powder tablets) and the broad group of elements measured (major elements like Al₂O₃ and Fe₂O₃, anthropogenically affected elements like Cd, Pb, Sb, trace elements like the REE) resulted in a data set of 87, partly overlapping, parameters. Each group of elements is thought to provide specific information on natural variation, human influence, or their interaction, as will the comparison between top and subsoil.

Precision was estimated by analysing about 15% of the ICP-MS samples and about 4% of the XRF samples in duplicate. The reproducibility is in general better for the XRF (0.5-10% relative to the mean) than for the ICP-MS (5-13% relative). The analytical variability is generally smaller than the local sampling variance as found in chapter 3, and therefore adequate for the desired regional evaluation of soil concentrations. The number of outliers found in the data also appears quite acceptable.

In this study the levelling of the data from each survey, based on Between Survey Duplicates (BSD), proved to be a good method to eliminate subsurvey biases. The fact that BSD levelling uses a set of samples that has the full range of element composition and concentration present, and is thus representative of the total survey, makes this method preferable to corrections based solely on analytical standards.

Based on the precision and accuracy a selection was made of 31 parameters that were measured in all surveys, and hence were sampled for virtually all locations and soil depths. This selection is thus optimised for multivariate statistical analysis by maximizing data quality and minimizing the number of missing values. This optimal dataset is limited compared to all the data analysed, but still comprises a sufficient subset of major, minor, and trace elements for environmental assessment of the soils of Zeeland. It is intended to form the core for all multivariate statistical and geochemical analysis, from which cross reference to the complete data set can still be made.

Summary statistics show that for most of the parameters neither a normal nor a lognormal distribution should be assumed. Although most major elements tend to a more or less

normal distribution, while most minor and trace elements are positively skewed, deviations from the standard type distributions are apparent. Suspected contaminants like Cd, Sb, As, Pb and Zn are more skewed than other trace elements.

Chapter 5. Enrichment and natural variability versus anthropogenic impact

Chapter 5 adopts a structured approach, using several statistical techniques that increase in complexity and the amount of auxiliary data needed. This approach provides detailed insight in the soil geochemistry of Zeeland. It first shows which components are most likely influenced by anthropogenic processes. Then, this anthropogenic imprint can be quantified based on their relation with non-influenced soil components.

Cumulative frequency plots are a good aid to show at a glance which elements have anomalous concentrations. From these plots percentiles can be derived thereby dividing anomalous data from background. The use of the value corresponding to the P_{90} , an often used standard percentile in Dutch soil surveys, as a baseline or threshold for assumed to be influenced elements might be a responsible choice, but only if the corresponding cumulative distribution plots are studied. In this study most (trace) elements are chiefly associated with natural processes and the P_{90} is an obvious underestimation, the P_{99} is more appropriate. Also for Pb the P_{90} is an underestimation, while for some other probably anthropogenically influenced elements (Cd, Cu and Sb) the P_{90} is an overestimation of the threshold value. This suggests that the arbitrary choice for the P_{90} in many Dutch soil surveys should be reconsidered. Notwithstanding the advantage of its simplicity, the method only shows if a components distribution deviates from normality with a distinct tail towards higher values. It does not show the nature of the deviation, anthropogenic or natural, nor does it provide a clear overview of the extent of the enrichments, due to the subjectivity and sometimes ambiguity in the choice of the threshold percentile. However, at a reconnaissance stage the method can be very useful.

The spatial distribution of elements with low threshold percentiles (P_{80} to P_{90}) and suspected of human influence, shows regional variability that differs distinctly from the more or less similar patterns that emerge for elements related to natural processes like variability in clay/ Al_2O_3 . This indicates that these probably influenced elements indeed have a single process underlying their regional variability.

Calculation of median ratios, with the upper continental crust (UCC), median subsoil concentrations, and legal limits, is an easy method to determine which elements are enriched compared to the reference used. Ratios with the UCC indicate that As, Cd, Cr, Pb, and Zr are relatively enriched. For As it is assumed that this enrichment is partially caused by the low lying deltaic environment of Zeeland, Cr and Zr could be related to heavy mineral concentrations, while the other elements might be enriched due to fertiliser usage and/or traffic exhaust. Depleted elements are Ba, Ca, K, Mg, Mn, Na, and Sr, which is assumed to be a natural regional feature. On average, the Zeeland soils also appear somewhat less

aluminous and ferromagnesian than the UCC. The ratios based on the subsoil show that the elements As, Cd, Cu, P, Pb, S, Sb, Sn, and Zn are enriched. These are all elements for which enrichments can usually be related to anthropogenic activities. The ratios with the official legal limits for Dutch soil, often used as reference values, show ratios of about 0.5-0.8, with the higher ratios (Cr, Ni) probably caused by the better analytical performance of the XRF method used as compared to the prescribed aqua-regia extraction. Even the P_{99} of the ratios, based on median clay values, are still below one for all elements.

The implications of the ratio results are basically twofold: 1) the area of Zeeland covered by this study is clean as regarding the governmental regulations and 2) the legal threshold limits are an overestimation of soil background concentrations, already including (minor) anthropogenic enrichments. They are thus less suitable as a reference value for a study into natural and anthropogenic geochemical patterns, and the ratios with the local reference (subsoil data) are preferred for identifying the influenced elements and determining the extent of their enrichment.

The correlation with Al_2O_3 is easily calculated and proved to be very useful to indicate elements which are possibly altered by human processes. The elements P_2O_5 , Cd, Cu, Pb, S, Sb, and Sn show relatively low correlations with Al_2O_3 in the topsoil compared to other elements and are also enriched in the topsoil. CaO, MnO, Na_2O , and Sr show relatively low, or negative, correlations with Al_2O_3 but are not enriched in the topsoil. The correlations confirm the similarities observed in the spatial distribution of the elements.

Overall higher correlations with Al_2O_3 in the reference subsoil confirm the hypothesis that the subsoil can be regarded as a pristine, non-influenced, reference. A graphical representation of the relation between the various elements and Al_2O_3 for both topsoil and subsoil clearly depicts the relative addition in the topsoil. Using a linear regression model based on the subsoil data the extent of the enrichment was quantified for each topsoil sample. When expressed as the ratio between observed concentration and calculated baseline concentration, the elements in order of decreasing enrichment are $Cd > Cu \gg Sn \approx Sb > Pb > Zn > As$. These enrichments are in line with data found for a similar area in the north of France, except for Cu, which is relatively higher in the Zeeland topsoils. Spatial distribution patterns of the enrichment ratios show that for most influenced elements regional variability is high and that hardly any spatial or mutual regional relations appear. The overall regional perspective provided by the contamination index shows that some areas in the north-east and centre of Zeeland are slightly more enriched on average.

Using the subsoil regression as a baseline has many advantages above the current Dutch practice of calculating a fixed legal threshold value, though corrected for clay and organic matter content. The natural random variability in baseline concentrations, expressed as the standard error of the regression, is a property of the model. This contrasts with the legal threshold value, which is just a rigid value with no range given. Also the regression model provides a local baseline, specifically suited for the parent soil material of Zeeland, which obviously should be preferred over a general model that tries to include all possible soil types. Compared to all other techniques presented, this subsoil regression baseline method gives the most profound insight into 1) the geological/pedological pattern of geochemical

variability, 2) random natural variation, and 3) the anthropogenic imprint on the Zeeland soil. These advantages definitely outweigh the greater complexity of the procedure.

Chapter 6. Regional diffuse geochemical patterns and processes

In this chapter I focus on the extent to which the human contributions actually affect the regional soil composition compared to the natural variability. Using the same data as in chapter 5, the processes or factors determining the regional variability, both in attribute and geographical space are assessed. The regional expression of the interaction of human and anthropogenic processes is further evaluated, to see if regions may be distinguished where specific human processes have had an effect on the soil composition.

The general covariance structure of the data was studied using robust principal component analysis (PCA) with varimax rotation to reveal the processes within attribute space. Groups of covarying geochemical components, may indicate certain processes. To assess which processes explain features in geographical space, clustering techniques were used to identify groups of sample points with similar properties. Due to the expectation that each pattern is a combination of different processes and the general multivariate nature of geochemical data, fuzzy c-means clustering (FCMC) was chosen.

The consecutive use of principal component analysis and fuzzy c-means clustering provided a profound insight into the processes affecting the regional geochemical variability in the Zeeland soils and into the extent of the variability caused by human processes.

The PCA reveals 4 factors that explain 76.6% of the overall variance, these factors are named for convenience “clay”, “calcite”, “anthropogenic” and “silt” respectively. The largest part of the variability (68%) is akin to geogenic clay, calcite, and silt related processes while 8.9% can be explained by anthropogenic processes. The elements related to the latter were the so called “heavy metals” Cd, Cu, Sb, Sn, and to a lesser extent Pb, Zn, and As. The regional variability of the clay, calcite, and silt factor resembles the regional variability of elements associated with this factor. The regional variability of the anthropogenic factor can be largely compared with the contamination index as obtained in chapter 5. When the elements of the anthropogenic factor were examined more closely, no separate subprocesses could be discerned other than unique behaviour for each element.

Considering that the soils of Zeeland are clearly enriched for some elements it is to be expected that this enrichment will also be revealed when looking at the processes within attribute space. The PCA confirmed this, but, the natural clay and calcite related processes still comprise the largest part of the soil variability. The anthropogenic influence on the soil only explains a small part of the variability emphasising that these anthropogenic processes only affect a small number of inorganic elements and are really an imprint on the overall natural variability.

The grouping of locations with FCMC, using the elements with the highest loadings of the PCA, resulted in an optimal 4 cluster model mainly dividing the area in calcareous heavy

clays, non-calcareous heavy clays, moderately calcareous silty clays, and non calcareous silts. No clusters exist where samples are discriminated solely on the basis of anthropogenic enrichment. This suggests that the extent of these human processes, clearly imprinted on the Zeeland soils, are not related to certain regions but rather are diffusely distributed over the province.

Chapter 7. Assessment of regional DDT concentrations in the soils of Zeeland

This chapter aims at providing a regional assessment of current DDT residue concentrations, variability, and degradation for the province of Zeeland. Based on a merging of several separate soil information systems (SIS) data, and comparing the concentrations with legal limits and data from other areas allowed the extent of contamination to be assessed, providing insight into whether the concentrations are just part of the ubiquitous background or exceed them. The geographical distribution of the DDT residue levels might indicate patterns, if any, that possibly relate to distinct processes. In addition, the data integration exercise may help assess the value of the combined SIS for regional environmental policy and may lead to recommendations for the (local) authorities for further research and monitoring.

The combination of the different SIS derived datasets resulted in a database that can be used for a regional assessment. However, the differences between the surveys, although in general based on the same sampling and analytical standards, are evident. It is hard to determine if these differences denote differences in true DDT soil values or that they are largely caused by the slight variations in surveying and laboratory methods.

The $\sum\text{DDx}$ concentrations (sum of the concentrations of all DDT isomers and metabolites) as found in Zeeland are comparable to those in similar agricultural areas, and based on these values there is no reason to believe that the current level of DDT contamination is more severe than can be expected for an agricultural area. Compared to pristine areas the level in Zeeland is about a factor 2 to 10 higher, which indicates that Zeeland can be regarded as enriched compared to a general ubiquitous $\sum\text{DDx}$ background. Despite that the values are within a normal range for agricultural areas, the standard soil normative threshold value S is exceeded for roughly half the cases and the soil specific value s in even more cases, which may result in litigations.

The relatively large contribution of local scale variability to the overall variability of $\sum\text{DDx}$ concentrations implies that a regional overview has limited value since most of the observed variance can be regarded as noise caused by small scale features. This is both indicated by the analysis of variance on replicate samples and by the nugget effect of the calculated semivariance.

Given both small scale variability and between survey bias, a final regional overview will contain large uncertainties. Despite that large scale features appear visible, like the relatively higher concentrations in the south of Zuid-Beveland which are presumably caused by

the presence of orchards, these should be judged with caution. Extreme caution is needed when the values are compared with (soil specific) legal limits. Due to the uncertainty caused by the various sources of variability, such comparison might have very limited value.

The degradation level of DDT residues indicates that the Zeeland soils will remain a source for $\sum\text{DDT}$ longer than expected. Since it is not assumed that new applications of DDT will occur, the monitoring of DDT residue levels could be cautionary. The most important issue which should be addressed is the reduction of variability in the monitoring data. Probably the largest reduction can be obtained in more standardisation of surveying, reducing the between survey bias. The small scale variance may be assessed by increasing the number of subsamples and sampling on smaller scales but this might be less cost-effective than reducing the between survey bias. However, policy based on these data should always incorporate the uncertainties as a result of the remaining variability.

Chapter 8. Synthesis

This study showed that with the use of state of the art analytical techniques and well established methods from geochemistry, much insight can be obtained from a single survey into the natural soil composition and how it is influenced by human processes. This both provides methods to obtain a reference for natural soil composition and to quantify the anthropogenic imprint. It is shown in this study that the framework of legal limits neither provides such reference nor is it able to detect subtle anthropogenic enrichment. This, in a way, is remarkable considering that this framework is also used for the building material decree and related to soil pollution risk maps that should prevent spreading of contaminants. I therefore suggest that references for soil composition should be derived from (local) baselines rather than legal limits. When obtaining such a baseline, quantification of the variability due to sampling and analytical procedures, and methods to reduce or compensate bias between sub-surveys are as important as the baseline itself. If multi-element techniques are used, like most laboratories do, focus should not only be on environmental priority components but also on components which might provide information about natural geochemical soil composition and variability.

Soil pollution risks in Zeeland appear to be, as defined in the Dutch normative framework, mainly determined by DDT residues. Unfortunately, the low average concentrations in soil, complex analytical determination, and large small-scale variability result in large uncertainties towards the regional variability of this component. Consequently, to obtain a more trustworthy georeferenced overview of DDT residue levels, more effort aimed at reduction of variance and bias should be put in the assessment of this component.

The question remains what “soil quality” should mean for the Dutch situation and which indicators to use. In this chapter it is argued that one of the goals to reach sustainable soil management should be a reconsideration of what is understood by “soil quality”. Only then it is possible to define which soil quality indicators should be assessed. The reference data and methods provided in this study could be a starting point for obtaining such indicators, finally leading towards sustainable soil management.

Samenvatting - Geochemische patronen in de bodems van Zeeland, natuurlijke variabiliteit versus antropogene impact

Hoofdstuk 1. Introductie

De sterke band tussen bodem, gezondheid en algemene duurzaamheid bevestigt dat de bodem een belangrijke rol speelt als grensvlak tussen milieu-compartimenten en dat bodemkwaliteit een belangrijke potentiële indicator is voor het duurzaam beheer van ons milieu.

Dit proefschrift wil een overzicht geven van de algemene (geo)chemische bodemkwaliteit in de provincie Zeeland, binnen het kader van diffuse antropogene vervuiling en duurzaam bodembeheer in Nederland. Het voornaamste doel daarbij is om de natuurlijke variabiliteit in de geochemische bodemsamenstelling te onderscheiden van antropogene beïnvloeding. De totale variabiliteit is geëvalueerd in zowel de geografische ruimte, waar de ruimtelijke interactie tussen de bodem-componenten plaatsvindt, alsook in de eigenschappen- of attribuut-ruimte, waarin die interactie eveneens tot uitdrukking komt. Patronen in beide ruimten kunnen dan gerelateerd worden aan processen die van invloed zijn op de bodemsamenstelling. Gebaseerd op deze patronen en de daaraan gerelateerde processen kan de antropogene invloed worden onderscheiden van de natuurlijke variabiliteit. Dit voorziet in informatie en gereedschappen ten nutte van het in het Nationaal Milieubeleids Plan 3 van de Nederlandse regering beoogde landelijk beeld van de algemene bodemkwaliteit.

De provincie Zeeland heeft een omvangrijk landelijk gebied, bestaand uit jonge Holocene afzettingen, en kent een rijke historie. Hierdoor is dit een bij uitstek geschikt gebied om de hoofdhypothese bij dit onderzoek te testen: dat menselijke invloed leidt tot onderscheidbare en kwantificeerbare patronen in bodemsamenstelling, in termen van ruimtelijke variabiliteit en onderlinge samenhang tussen bodem-componenten.

Hoofdstuk 2. Geologie en pedologie van Zeeland

Zeeland is gesitueerd in de mariene delta van de rivieren Rijn, Maas en Schelde. Door de zeespiegelstijging aan het einde van de laatste grote ijstijd, het Weichsel, kwam Zeeland

in de greep van de zee. De bodems van Zeeland kunnen worden beschouwd als relatief homogeen en jong. Zij zijn gevormd in een continue interactie van natuurlijke processen en menselijke ambities.

De bodems van Zeeland zijn in het algemeen ontwikkeld in mariene klei-afzettingen. Het ligt daarom in de verwachting dat variatie in het kleigehalte de belangrijkste omstandigheid is die de geochemische variatie bepaalt. Het natuurlijke patroon van kreekruggen en poelgronden creëert hierin al relatief abrupte overgangen. Homogeniteit of graduele variatie binnen dergelijke eenheden kan verder fysiek zijn verstoord door menselijke activiteiten. Waarschijnlijk is de variabiliteit in de bodemsamenstelling daarom hoger dan direct blijkt uit de bodemkaart. Elk bedijkt gebied heeft zijn eigen geschiedenis, zowel menselijk als sedimentair, en daarmee een eigen, uniek patroon van bodemvariatie. Daarnaast kan verwacht worden, gezien het feit dat 80% van het Zeeuwse landoppervlak in gebruik is voor de landbouw, waarvan 70% voor akkerbouw, dat gebruik van bemestings- en bestrijdingsmiddelen de variabiliteit in de bodem verder hebben beïnvloed. Hierbij zullen met name verhoogde concentraties van zogenaamde “zware metalen” (Cd, Cu, Pb en Zn) en restanten van persistente organochloor bestrijdingsmiddelen optreden. Als laatste moet ook atmosferische toevoer worden beschouwd als bijdrage aan de bodemgeochemie.

Hoofdstuk 3. Natuurlijke en antropogene covariantie patronen en ruimtelijke variabiliteit van sporenelementen in de bovenlaag van landbouwgronden

Het doel van dit hoofdstuk is om de mate van geochemische variabiliteit in de bodemsamenstelling te analyseren als functie van de ruimtelijke schaal en de bemonsteringsprocedure (wel of geen mengmonsters). Het onderzoek was ingegeven door vragen uit de praktijk aangaande bemonsteringsstrategieën voor regionale bodemonderzoeken. Een zogenaamde ongebalanceerde geneste bemonsteringsstrategie, met enkelvoudige monsters, is uitgevoerd op Walcheren en Zuid-Beveland. Hierbij worden series van monster-paren op (exponentieel) oplopende afstand (schaal) gecreëerd. De toename in ruis-variantie door gebruik van enkelvoudige monsters is onderzocht door vergelijking met de resultaten van een eerdere studie in het gebied op basis van mengmonsters.

De monsters zijn opgelost in koningswater en geanalyseerd met ICP-MS. De resulterende data (40 elementen) zijn onderzocht met behulp van principaal component analyse in combinatie met een varimax rotatie (PCA-V), ongebalanceerde variantie-analyse (UANOVA) en fuzzy c-means clusteranalyse op de UANOVA resultaten. De drie factoren van de PCA-V representeren variabiliteit ten gevolge van respectievelijk natuurlijke, menselijke en geohydrologische processen. De UANOVA analyse onderscheidt drie patronen van ruimtelijke variantie. Twee groepen laten een graduele toename zien van de variantie met de afstand. De ene groep bevat klei/veldspaat gerelateerde elementen inclusief de lichtere lantaniden of zeldzame aarden, de andere groep bevat de zwaardere zeldzame aarden. De derde groep bevat alle antropogene en geohydrologisch beïnvloede elementen. Voor deze

laatste groep treedt een groter deel van de variatie al op lokale schaal op.

In het algemeen domineert de regionale variantie over de lokale variantie en de extra inspanning die het nemen van mengmonsters vraagt blijkt van beperkte waarde voor een regionaal onderzoek. Echter, prioritaire metalen zoals Cd, Cu, Pb, Sb, Sn en Zn, hebben een relatief grotere lokale variantie-component. Voor deze elementen is het nemen van mengmonsters relatief gezien gunstiger.

Hoofdstuk 4. Bemonstering en analyse voor een regionale milieukundig geochemisch bodemonderzoek

Het onderkennen van antropogene effecten (aanrijking) ten opzichte van mogelijke natuurlijke variabiliteit op basis van waargenomen concentraties van individuele elementen is uiterst moeilijk. Het vaststellen van een bi-variate of multi-variate geochemische referentie (baseline) is daarom een algemeen geaccepteerde methode om achtergrondwaarden vast te stellen. Hierbij wordt een eventuele aanrijking relatief beschouwd ten opzichte van andere bodemchemische kenmerken die een groot deel van de natuurlijke variabiliteit beschrijven. Een afwijking ten opzichte van de aldus bepaalde achtergrond is dan te interpreteren als mogelijk veroorzaakt door menselijke verstoring. Om op die manier kwantitatief te onderzoeken welke componenten zijn beïnvloed door menselijke activiteiten is dan wel een consistente regionale multivariate dataset nodig, die ook aan hoge eisen betreffende betrouwbaarheid en juistheid moet voldoen. Dit hoofdstuk documenteert de constructie van zo'n dataset voor de provincie Zeeland.

De bemonsteringsstrategie was gericht op het bemonsteren van zowel hedendaagse als oorspronkelijke, onbeïnvloede bodemconcentraties. De bemonsteringsdichtheid bedroeg 1 locatie per 1-10 km² en zowel de beïnvloede bovenlaag van de bodem als een dieper gelegen, verondersteld pre-antropogene, laag werden bemonsterd. De gebruikte analyse-technieken (ICP-MS met koningswater-ontsluiting en XRF met tabletten) en de brede groep gemeten elementen (hoofdelementen zoals Al₂O₃ en Fe₂O₃, antropogene beïnvloede elementen zoals Cd, Pb, Sb, sporenelementen zoals de lantaniden) resulteerde in een dataset van 87, gedeeltelijk overlappende, parameters. Van elke groep elementen, alsook van de vergelijking tussen de bovenlaag en diepere bodem, wordt verwacht dat zij specifieke informatie kunnen leveren over natuurlijke variatie, menselijke invloed of hun interactie.

De betrouwbaarheid (precisie, reproduceerbaarheid) van de bepalingen is geschat door 15% van de ICP-MS monsters en circa 4% van de XRF monsters in duplo te analyseren. De reproduceerbaarheid is in het algemeen beter voor de XRF (0.5-10% relatief) dan voor de ICP-MS (5-13% relatief). Deze analytische variabiliteit is in het algemeen kleiner dan de lokale veld-variantie zoals gevonden in hoofdstuk 3 en daarmee adequaat voor de gewenste regionale evaluatie van de bodemconcentraties. De dataset bleek slechts een beperkt aantal onverklaarde uitbijters te bevatten, wat eveneens als zeer acceptabel is beoordeeld.

In deze studie is het corrigeren van de data, gebaseerd op "Between Survey Duplicates" (BSD) een goede methode gebleken om systematische afwijkingen tussen de verschil-

lende meetrondes in het totale bodemonderzoek te elimineren. De BSD methode maakt gebruik van een deelverzameling van de monsters die de volledige bandbreedte in elementconcentraties beslaat, en dus representatief is voor het onderzoek. De BSD methode verdient daarmee de sterke voorkeur boven correctie-methoden gebaseerd op alleen analytische standaarden.

Op basis van de geraamde precisie en juistheid van analyse is uit het totaal aan parameters een selectie gemaakt van 31 parameters, die bovendien zijn bepaald in elke meetronde en dus voor alle locaties en bodemlagen. Door het op deze manier maximaliseren van de kwaliteit van de data en het minimaliseren van het aantal ontbrekende waarden is deze selectie optimaal voor multivariate statistische analyse. Deze optimale dataset bevat nog steeds voldoende diversiteit in hoofd- en sporenelementen voor de gewenste milieukundige beoordeling van de bodems van Zeeland. Hij vormt derhalve de gewaarborgde kern voor alle data-analyses, waarbij echter nog steeds gerefereerd kan worden aan de totale dataset.

Statistische kengetallen laten zien dat de meeste parameters niet de standaard normale of lognormale verdeling volgen. Hoewel de meeste hoofdelementen neigen naar een min of meer normale distributie, en de meeste sporenelementen scheef verdeeld zijn naar de hogere concentraties, zijn afwijkingen van de standaardverdeling overduidelijk. Mogelijke contaminanten zoals Cd, Sb, As, Pb en Zn zijn meer scheef verdeeld dan andere sporenelementen.

Hoofdstuk 5. Aanrijking en natuurlijke variabiliteit versus antropogene impact

Voor het bepalen en kwantificeren van antropogene beïnvloeding wordt een gestructureerde aanpak gevolgd, gebruik makend van diverse statistische technieken die toenemen in complexiteit en het gebruik van extra data. Deze aanpak verschaft gedetailleerd inzicht in de bodemgeochemie van Zeeland. Hij laat ten eerste zien welke componenten zeer waarschijnlijk zijn beïnvloed door menselijke activiteit. Vervolgens blijkt deze antropogene invloed gekwantificeerd te kunnen worden, gebaseerd op een relatie met niet beïnvloede bodem-componenten.

Cumulatieve frequentie grafieken zijn een goede hulp om in één oogopslag te zien welke elementen voor een deel van de monsters een afwijkende concentratie hebben. Uit deze grafieken kunnen in principe percentielen in de frequentieverdeling worden afgeleid die de grens vormen tussen achtergrond en afwijkende data. Een veel gebruikte standaard bij Nederlands bodemonderzoek is om de waarde corresponderend met de P_{90} te kiezen als referentie- of drempelwaarde voor potentieel beïnvloede elementen. Dit is echter alleen verantwoord als ook de bijbehorende cumulatieve frequentie grafieken worden bestudeerd. In deze studie zijn de meeste (sporen) elementen voornamelijk geassocieerd met natuurlijke processen en de P_{90} is een overduidelijke onderschatting, de P_{99} is meer geschikt. Ook voor Pb is de P_{90} een onderschatting, terwijl voor sommige andere waarschijnlijk antropogeen beïnvloede elementen (Cd, Cu en Sb) de P_{90} een overschatting van de drempelwaarde

is. Dit suggereert dat de arbitraire keuze voor de P_{90} in veel Nederlandse bodemonderzoeken in heroverweging genomen moet worden. De cumulatieve frequentie methode heeft het voordeel van haar eenvoud, maar laat alleen zien of de statistische frequentie verdeling van een bodemcomponent al of niet afwijkt van normaliteit met een duidelijk neiging naar hogere waarden. De methode geeft geen aanwijzing voor de aard van de afwijking, natuurlijk of antropogeen. Ook wordt geen duidelijk beeld voorzien van de mate van aanrijking door de subjectiviteit en soms ambivalentie in de keuze van de drempelwaarde. Echter in een verkennende fase kan deze methode zeer bruikbaar zijn.

De ruimtelijke verdeling van de elementen met lage drempelwaarden (P_{80} tot P_{90}), en waarschijnlijk beïnvloed door de mens, laat een variabiliteit zien die duidelijk afwijkt van de min of meer overeenkomende ruimtelijke patronen die de elementen laten zien die gerelateerd zijn aan natuurlijke processen, zoals de variabiliteit in klei-gehalte. Dit duidt erop dat voor deze mogelijk beïnvloede elementen een uniek proces ten grondslag ligt aan de regionale variabiliteit.

Berekening van mediane ratio's met de standaardwaarden voor de buitenste continentale korst (upper continental crust, UCC), met de mediane waarden zoals bepaald voor de diepere Zeeuwse bodem, en met de wettelijke Streefwaarden, is een eenvoudige methode om te bepalen welke elementen aangerijkt zijn in vergelijking tot de gebruikte referentie. Ratio's met de UCC duiden erop dat As, Cd, Cr, Pb en Zr relatief aangerijkt zijn. Voor As wordt aangenomen dat deze aanrijking deels wordt veroorzaakt door de laag gelegen deltaïsche positie van Zeeland. Cr en Zr kunnen gerelateerd worden aan concentraties van zware mineralen, terwijl de ander elementen mogelijk zijn aangerijkt door kunstmestgebruik en/of uitstoot van verkeer. Relatief verarmde elementen zijn Ba, Ca, K, Mg, Mn, Na en Sr, waarvan wordt aangenomen dat dit een natuurlijke regionale eigenschap is voor Zeeland. Gemiddeld lijkt het dat de Zeeuwse bodems iets armer in Al en Fe is dan de UCC. De ratio's gebaseerd op de diepere bodem laten zien dat de elementen As, Cd, Cu, P, Pb, S, Sb, Sn en Zn aangerijkt zijn. Dit zijn allemaal elementen waarvoor aanrijking in het algemeen gerelateerd kan worden aan antropogene activiteiten. De ratio's met de Streefwaarden, vaak gebruikt als referentiewaarde voor de mate van verontreiniging, laten waarden zien tussen de 0.5-0.8. Hierbij worden de hogere ratio's (Cr, Ni) waarschijnlijk veroorzaakt door de hogere analytische opbrengst van de XRF totaal-methode in vergelijking met de koningswater extractie. Zelfs de P_{99} van de ratio's gebaseerd op Streefwaarden voor het mediane klei-gehalte in Zeeland, zijn lager dan 1 voor alle elementen.

De implicaties van de ratio-resultaten zijn tweeledig: 1) het landelijk gebied in Zeeland zoals beschouwd in deze studie is schoon volgens wettelijke maatstaven en 2) de Streefwaarden, die duidelijk een zekere algemene(lichte) mate van antropogene aanrijkingen al incorporeren, zijn een overschatting van de natuurlijke achtergrondwaarden in de bodem. Zij zijn dus minder geschikt als referentiewaarden in een studie naar natuurlijke en antropogene geochemische patronen. De ratio's met de lokale referentie (data van de diepere bodem) hebben de voorkeur bij de identificatie van de beïnvloede elementen en het vaststellen van de mate van aanrijking.

De correlatie met Al_2O_3 is eenvoudig te berekenen en is zeer bruikbaar gebleken om elementen te duiden die mogelijk zijn beïnvloed door menselijk processen. De componenten

P₂O₅, Cd, Cu, Pb, S, Sben Sn laten, vergeleken met andere elementen, een relatief lagere correlatie zien met Al₂O₃ in de bovenlaag van de bodem en zijn ook aangerijkt ten opzichte van de diepere bodem. CaO, MnO, Na₂O en Sr laten een relatief lage, of negatieve correlatie zien met Al₂O₃ maar zijn niet aangerijkt in de bovenlaag. De correlaties bevestigen de overeenkomsten zoals waargenomen in de ruimtelijke patronen van de elementen.

De in het algemeen hogere correlaties met Al₂O₃ in de diepere bodem bevestigen de hypothese dat de laatste kan worden beschouwd als een oorspronkelijke, niet beïnvloede, referentie. Een grafische weergave van de relatie tussen de verschillende elementen en Al₂O₃ voor zowel de bovenlaag als de diepere bodem laat duidelijk de relatieve aanrijking van de bovenlaag zien. Door gebruik te maken van een lineair regressiemodel gebaseerd op alle data van de diepere bodem, kan de mate van aanrijking in de bovenlaag voor iedere afzonderlijke locatie gekwantificeerd worden. De elementen in volgorde van afnemende aanrijking, uitgedrukt als de gemiddelde of mediane ratio tussen de werkelijk gemeten waarde en de uit de regressievergelijking berekende referentiewaarde, is Cd > Cu ≫ Sn ≈ Sb > Pb > Zn > As. Deze mate van aanrijking komt overeen met data uit een vergelijkbaar gebied in het noorden van Frankrijk, behalve voor Cu dat in Zeeland relatief meer is aangerijkt. Ruimtelijke verdelingspatronen van de aanrijgings-ratio's laten zien dat voor de meeste beïnvloede elementen de regionale variabiliteit groot is en dat nauwelijks enige ruimtelijke of onderlinge relaties voorkomen. Het algemene regionale beeld zoals blijkend uit de gesommeerde contaminatie-index laat zien dat sommige gebieden in het noord-oosten en midden van Zeeland relatief meer zijn aangerijkt.

De regressie met de diepere bodem als referentie biedt veel voordelen ten opzichte van de huidige Nederlandse praktijk van het vergelijken met de vaste Streefwaarden, ofschoon gecorrigeerd voor klei en organisch stof gehalte. De natuurlijke willekeurige variabiliteit in de referentie-concentratie, uitgedrukt als de standaardfout van de regressie, is een via het regressiemodel bepaald kenmerk. Dit contrasteert met de wettelijke Streefwaarde, die slechts n getal zonder bandbreedte geeft. Tevens voorziet het regressiemodel in een lokale referentie, specifiek voor het natuurlijke bodemmateriaal van Zeeland, hetgeen zeker de voorkeur verdient boven een algemeen model dat probeert alle mogelijke bodemtypen te omvatten. Vergeleken met de andere technieken, geeft de regressie met de diepere bodem als referentie het meest diepgaand inzicht in 1) de geologische/pedologische patronen in geochemische variabiliteit, 2) de willekeurige natuurlijke variatie en 3) de antropogene aanrijking van de Zeeuwse bodem. De nadelen van de grotere complexiteit wegen niet op tegen bovengenoemde voordelen van deze methode.

Hoofdstuk 6. Regionale diffuse geochemische patronen en processen

In dit hoofdstuk concentreer ik me op de mate waarin de menselijke bijdrage werkelijk de regionale bodemsamenstelling beïnvloedt in vergelijking met de natuurlijk variabiliteit. Gebruik makend van dezelfde data als in hoofdstuk 5 worden de processen en factoren, die

de regionale variabiliteit in de geografische en attribuut ruimte bepalen, onderzocht. De regionale expressie van de interactie van menselijke en natuurlijke processen wordt nader bekeken om te zien of er regio's te onderscheiden zijn waar specifieke menselijke processen effect hebben gehad op de bodemsamenstelling.

De algemene covariantiestructuur van de data wordt bestudeerd met behulp van principaal component analyse (PCA) om zo processen in de attribuut-ruimte te laten zien. De groepering van onderling covariërende geochemische componenten vormt aanwijzingen voor de aard van deze processen. Een clustering-techniek is gebruikt om groepen van monsterpunten met gelijke eigenschappen te identificeren, om zo de relatie tussen de geïdentificeerde processen en locatie in de geografische ruimte te leggen. Een fuzzy clustering (FCMC) werd gekozen omdat verwacht werd dat het uiteindelijke totale patroon een combinatie of overlap is van verschillende processen.

Het achtereenvolgende gebruik van PCA en FCMC leverde een duidelijk inzicht in de processen die van invloed zijn op de regionale geochemische bodemsamenstelling in Zeeland en in de mate van de variabiliteit veroorzaakt door de mens.

De PCA onthult 4 factoren die 76.6% van de variantie verklaren. Deze factoren zijn voor het gemak "klei", "calciet", "antropogeen" en "silt" genoemd. Het grootste gedeelte van de variabiliteit (68%) is gerelateerd aan de geogene klei, calciet en silt gerelateerde processen terwijl 8.9% verklaard kan worden met antropogene processen. De elementen die gerelateerd zijn aan het laatste zijn de zogenaamde "zware metalen" Cd, Cu, Sb, Sn en in mindere mate Pb, Zn en As. De regionale variabiliteit van de klei, calciet en silt factoren komt overeen met de regionale variabiliteit van de specifieke elementen die aan deze respectieve factoren geassocieerd zijn. De regionale variabiliteit van de antropogene factor kan grotendeels worden vergeleken met die van de contaminatie index zoals verkregen in hoofdstuk 5. Wanneer de elementen op de antropogene factor nader worden bekeken kunnen geen aparte subprocessen onderscheiden worden, anders dan uniek gedrag van elk element.

Gegeven de al eerder aangetoonde aanrijking van de bovenlaag van de Zeeuwse bodems met sommige elementen, is het te verwachten dat dit ook naar voren komt bij het bekijken van proces-gerelateerde patronen in de attribuut-ruimte. De PCA bevestigde dit inderdaad, maar de klei en calciet gerelateerde processen bepalen nog steeds het grootste gedeelte van de variabiliteit in bodemsamenstelling. De antropogene invloed op de bodemsamenstelling verklaart slechts een klein deel van de totale variabiliteit. Dit benadrukt dat deze antropogene processen slechts een klein aantal anorganische elementen beïnvloeden en maar in beperkte mate de al aanwezige, natuurlijke variabiliteit vergroten.

Het groeperen van de locaties met behulp van FCMC, gebruik makend van die elementen die door de PCA werden geïdentificeerd als meest karakteristiek voor de verschillende processen, resulteerde in een optimaal 4-cluster model. Hiermee werd het gebied hoofdzakelijk opgedeeld in kalkrijke klei, kalkarme klei, matig kalkhoudende siltige klei en kalkarme silt. Er zijn geen clusters van monsters gevonden die kunnen worden onderscheiden op specifiek antropogene aanrijking. Dit suggereert dat deze menselijke processen, waarvan de invloed duidelijk aanwezig is in de Zeeuwse bodems, niet gerelateerd zijn aan specifieke

regio's maar eerder diffuus verspreid zijn over de provincie.

Hoofdstuk 7. Beoordeling van de regionale DDT concentraties in de bodems van Zeeland

Dit hoofdstuk tracht te voorzien in een regionale beoordeling van de DDT-problematiek in de Provincie Zeeland betreffende huidige resterende DDT concentraties, variabiliteit en afbraak. De data zijn samengevoegd uit een aantal verschillende bodem informatie systemen (soil information systems, SIS). Vergelijking van de concentraties met wettelijke normen en gegevens uit andere gebieden maken het mogelijk de mate van vervuiling te beoordelen. Dit laatste geeft inzicht of de concentraties slechts een onderdeel zijn van de alomtegenwoordige achtergrond of deze overstijgen. Ook de geografische verdeling van de restant DDT waarden laat mogelijk patronen zien die kunnen duiden op bepaalde processen. Daarnaast kan het integreren van de DDT data helpen om de waarde van de gecombineerde SIS voor regionaal milieukundig beleid te bepalen, mogelijk leidend tot aanbevelingen voor de (lokale) autoriteiten voor toekomstig onderzoek en monitoring.

De combinatie van de uit de diverse SIS betrokken sub-datasets resulteerde in een database die kon worden gebruikt voor een regionale beoordeling. De verschillen tussen de afzonderlijke datasets zijn echter overduidelijk, ondanks dat ze in het algemeen allemaal gebaseerd zijn op dezelfde bemonsterings- en analyse-voorschriften. Het is moeilijk te bepalen of deze verschillen duiden op reële verschillen in DDT concentraties of dat zijn grotendeels worden veroorzaakt door de te verwachten kleine variaties in bemonsterings- en laboratorium-methoden.

De $\sum DDx$ concentratie (som van de concentraties van alle DDT isomeren en metaboolieten) zoals gevonden in Zeeland zijn vergelijkbaar met die in overeenkomstige agrarische gebieden. Er is daarom geen reden om aan te nemen dat het huidige niveau van DDT contaminatie ernstiger is dan wat verwacht mag worden voor een agrarisch gebied. Vergelijken met niet beïnvloede gebieden is het niveau in Zeeland een factor 2-10 hoger wat erop duidt dat Zeeland beschouwd kan worden als aangerijkt ten opzichte van de wereldwijd aanwezige achtergrond. Alhoewel de waarden dus binnen een normale bandbreedte vallen voor agrarische gebieden, wordt de ongecorrigeerde streefwaarde S in de helft van de gevallen overschreden en de (voor lutum en organisch stof) gecorrigeerde streefwaarde s in nog meer gevallen, wat kan leiden tot wettelijke problemen.

De relatief grote bijdrage van de variabiliteit op lokale schaal aan de totale variabiliteit van de $\sum DDx$ concentraties houdt in dat een ruimtelijk regionaal beeld van beperkte waarde is aangezien het grootste gedeelte van de waargenomen regionale variantie beschouwd kan worden als ruis, veroorzaakt door kleinschalige processen of factoren. Dit volgt zowel uit de variantie-analyse op herhaald uitgevoerde bemonsteringen, als uit het grote "nugget effect" in de berekende semivariantie.

Gegeven de kleinschalige variabiliteit en de afwijkingen tussen de verschillende datasets, zal het uiteindelijke ruimtelijke beeld grote onzekerheden bevatten. Ondanks dat grootscha-

lige kenmerken zichtbaar lijken te zijn, zoals hogere concentraties in het zuiden van Zuid-Beveland die waarschijnlijk worden veroorzaakt door de aanwezigheid van boomgaarden, zullen deze met voorzichtigheid moeten worden beoordeeld. Extreme voorzichtigheid is nodig als de waarden worden vergeleken met de (gecorrigeerde) streefwaarde. Door de grote onnauwkeurigheid veroorzaakt door de diverse foutbronnen heeft zo'n toetsing zeer beperkte waarde.

Het lage niveau van afbraak van DDT, zoals blijkend uit de verhouding tussen metaboliëten en restant-DDT, duiden erop dat de Zeeuwse bodems een bron blijven voor $\sum DDT_x$ voor een langere tijd dan mogelijk verwacht. Aangezien niet wordt aangenomen dat nu nog toepassing van DDT plaatsvindt zal het monitoren van het DDT niveau vooral een inzicht verschaffend karakter kunnen hebben. Het belangrijkste punt dat aandacht behoeft is het terugdringen van de ruis-variabiliteit. Waarschijnlijk kan de grootste reductie behaald worden door het verder standaardiseren van de bemonstering, wat de systematische verschillen tussen de individuele onderzoeken vermindert. De invloed van de kleinschalige variabiliteit kan aangepakt worden door het vergroten van het aantal submonsters of het verhogen van de mosnterdichtheid, maar dit is wellicht minder kosten-effectief. Van belang is dat het beleid, gebaseerd op de huidige of betere data, altijd rekening zal moeten houden met de gegeven onzekerheden als gevolg van de resterende variabiliteit.

Hoofdstuk 8, Synthese

Deze studie heeft laten zien dat door het gebruik van state-of-the-art analytische technieken en bewezen methoden uit de geochemie, veel inzicht verkregen kan worden in onderzoek naar de natuurlijke bodemsamenstelling en hoe deze wordt beïnvloed door menselijke processen. De aanpak voorziet in zowel een methode om een referentie voor natuurlijke bodemwaarden te krijgen als om de menselijke invloed te kwantificeren. Het raamwerk van wettelijke normen voorziet niet in een dergelijke referentie, en is niet in staat om subtiele antropogene invloeden te onderscheiden. Dit raamwerk voldoet voor zover toegepast in het kader van de bodemsanering, maar het gebruik van de Streefwaarde als basisnorm voor het bouwstoffenbesluit en gerelateerde bodemkwaliteitskaarten die het verspreiden van contaminanten tegen moet gaan, ligt minder voor de hand. Daarom suggereer ik dat in dat geval de referentie voor "schone" bodem zou moeten worden afgeleid van een lokale referentie en niet van wettelijke normen. Bij het verkrijgen van zo'n referentie zijn het kwantificeren van de variabiliteit ten gevolge van bemonstering en analytische procedures, en methoden om afwijkingen tussen separate onderzoeken te reduceren of te compenseren, net zo belangrijk als de referentie zelf. Indien multi-element technieken worden gebruikt, zoals de meeste laboratoria al doen, dan zal er niet alleen aandacht moeten worden geschonken aan de prioritaire componenten maar ook aan componenten die mogelijk informatie verschaffen over de natuurlijk geochemische samenstelling en variabiliteit.

Bodemkwaliteit (of beter vervuilingsrisico) in Zeeland lijkt, volgens de Nederlandse wettelijke normen, voornamelijk bepaald te worden door restanten DDT. Helaas resulteert de gemiddeld lage concentratie, grote kleinschalige variabiliteit en complexe analytische me-

thode in een hoge bepalings-onzekerheid met betrekking tot deze component. Voor het verkrijgen van een meer betrouwbaar “georeferenced” beeld van de DDT niveaus zal meer inspanning gericht moeten worden op het verminderen van variantie en afwijking tussen onderzoeken.

De vraag blijft wat “bodemkwaliteit” betekent voor de Nederlandse situatie en welke indicatoren daarvoor gebruikt moeten worden. In dit hoofdstuk is beargumenteerd dat één van de doelen om duurzaam bodembeleid te bereiken een heroverweging en verbreding van het begrip “bodemkwaliteit” zou moeten zijn. Alleen dan is het mogelijk om te bepalen welke, probleem-specifieke, bodemkwaliteitsindicatoren moeten worden beoordeeld. De referentie-data en methoden uit deze studie kunnen een startpunt vormen voor het verkrijgen van zulke indicatoren, uiteindelijk leidend tot duurzaam beheer van de bodem.

Dankwoord

“Goed, dit doen we dus nooit meer”, het was het enige waar ik aan kon denken terwijl ik één voor één de blaadjes van de eerste complete versie van dit proefschrift uit de zacht zoemende printer haalde. Het is een vreemde gewaarwording om datgene waar je maanden aan geschreven hebt opeens vorm te zien krijgen in een bijna eindeloos pak papier.

Onderzoek doen is leuk, een beetje data vinden, er wat met spelen, dan wat verstandige dingen zeggen over de uitkomsten en daar vervolgens eens stevig over discussiëren. Het gaat bijna allemaal vanzelf maar dan komt het schrijven. . . Toch is niets leuker dan de ideeën uit je onderzoek aan de wereld te presenteren. Waarbij ik me meteen besef dat daarmee de discussie pas echt begint.

Ondertussen heeft de printer overuren gedraaid en het eindresultaat ligt nu voor u. Ofschoon mijn naam onder het werk staat is dit niet het resultaat van één persoon. Een Noorse professor heeft mij ooit uitgelegd dat promoveren een redelijke egoïstische bezigheid is. Ofschoon ik het deels met hem eens ben, heb ik me daar niet bij willen neerleggen. Onderzoek doen en promoveren is in mijn ogen geen werk voor een solist maar het heeft input en feedback nodig van meerdere personen. Daarom ben ik heel blij dat een groot aantal mensen heeft bijgedragen door middel van discussies en opmerkingen of simpelweg wilde luisteren hoewel zij soms totaal niet begrepen waar ik het over had.

Als eerste wil natuurlijk mijn co-promotores bedanken, Pauline van Gaans en Simon Vriend. De vele, soms langdurige, onderlinge discussies hebben grotendeels de ideeën uit dit proefschrift gevormd. Daarbij hebben hun vele opmerkingen over de verschillende versies van de manuscripten (soms misschien wat teveel versies) veel bijgedragen aan het eindresultaat. Mijn promotor, Peter Burrough, wil ik met name danken voor de bijdrage in de latse fase van dit proefschrift. Het was heel verhelderend hoe hij als iemand zonder directe geochemische achtergrond naar het manuscript heeft gekeken en dit heeft becommentarieerd.

Natuurlijk heeft ook de Provincie Zeeland, en met name Ad Vos, een heel belangrijke bijdrage geleverd aan dit onderzoek. Met name de input van Ad maakte het mogelijk om op de hoogte te blijven van wat speelde binnen de (Zeeuwse) bodemwereld en om enkele ideeën uit dit proefschrift in de praktijk uit te proberen.

Wat voor paaseieren geldt, geldt ook voor ons: één AiO is geen AiO. Mijn mede AiO's, Grishja, Marc en met name mijn ex-kamergenoot Pieter-Jan ben ik ook dankbaar. Naast het feit dat het altijd leuk is om over elkaars werk te discussiëren zijn dergelijke collega's

ook nuttig om wat minder wetenschappelijke informatie uit te wisselen. Variërend van de kwaliteit van de Spar koffie tot het uiten van de diverse frustraties die altijd in de loop van een promotie op duiken.

Dit onderzoek was er niet geweest zonder data en de data was er niet geweest zonder laboratorium. De diverse medewerkers van het geochemische laboratorium van TNO-NITG en de UU, Gerard, Bertil, Hans, Erik, Dineke, Rob en Kathrin wil ik graag bedanken voor hun bijdrage aan de chemische analyses. Tijdens het onderzoek is temeer gebleken dat een goede band met het laboratorium direct bijdraagt aan de kwaliteit van de data. Daarnaast wil ik Jerfaas de Jong bedanken voor de eerste gegevens voor dit onderzoek en de hulp bij het verzamelen van de gegevens voor hoofdstuk 3. Tijdens de veldwerken voor de surveys 3 en 4 ben ik bijgestaan door mijn “boor-slaven”: Atilla, Joris, Sander, Vincent, Janneke, Helene en Johan. Niets is zo mooi als een ander voor je te zien werken, voor dat beeld ben ik hen dan ook zeker dankbaar.

Vanzelfsprekend mag ik de rest van mijn collega’s op het departement (of afdeling, vakgroep, al naar gelang welke organisatorische vernieuwing men gebruikt) fysische geografie ook niet vergeten, met z’n allen zorgden ze voor een prettige en motiverende werksfeer. Aan het obligate rijtje namen waag ik mij niet, dan vergeet ik er geheid één. Toch maak ik direct een paar uitzonderingen. Eentje voor de burens: Kor, Derk-Jan, de Jopies (Hansakwast en de Sluitersnuiten) en m’n huidige kamergenoot Reinder; hun gesjouw met koffie heeft mij op cruciale momenten wakker gehouden. Eentje voor Edzer, Maarten ZvE, en Thom voor de vele adviezen. En als laatste voor hen die in stilte veel van het onderzoek ondersteund hebben, de mensen van de werkplaats, materiaaluitleen, secretariaat en FG-lab.

De afgelopen jaren heb ik veel ondersteuning gekregen van mijn vrienden, bewust of onbewust hebben zij bijgedragen aan veel goede ideeën voor mijn promotie en voor de periode er na. Mijn ouders ben ik vooral dankbaar voor de steun tijdens mijn studie en ontwikkeling. Zonder hen was ik nooit zover gekomen. Lieve Marije, jou wil ik vooral bedanken voor je geduld wanneer ik me weer had vastgebeten in dit onderzoek en al die fijne momenten als je me wist los te schudden.

Job Spijker

Curriculum Vitae

I was born in IJmuiden, the Netherlands, on January 13, 1973. I graduated secondary school in 1992 and started my professional education at the chemistry faculty of the Hogeschool Alkmaar. During my qualification for chemical engineer I studied the implications of the use of herbicides on the production of drinking water. In 1996 I started my study at the geochemistry department of the faculty of Earth Sciences at the Utrecht University, specialising in environmental geochemistry. For my master graduation I studied the variability of soil composition on various spatial scales, the latest results are presented in chapter 3 of this thesis. From 2000 to 2005 I worked on my Ph.D. at the department of physical geography of the Utrecht University.